MEAN CURVATURE FLOW WITH SURGERY

ROBERT HASLHOFER AND BRUCE KLEINER

ABSTRACT. We give a new proof for the existence of mean curvature flow with surgery of 2-convex hypersurfaces in \mathbb{R}^N , as announced in [HK13a]. Our proof works for all $N \geq 3$, including mean convex surfaces in \mathbb{R}^3 . We also derive a priori estimates for a more general class of flows in a local and flexible setting.

Contents

2
2
2
4
6
9
9
.1
5
6
9
9
22
24
24
26
29

Date: April 30, 2015.

B.K. was supported by NSF grant DMS-1105656.

Appendix A.	One-sided minimization	31
References		33

1. INTRODUCTION

1.1. **Background.** Mean curvature flow, like Ricci flow and harmonic map heat flow, develops singularities, and understanding them is a central problem in geometric analysis. In the last decades several methods have been developed to continue the flow beyond the first singular time: Brakke flow [Bra78], level set flow [ES91, CGG91], and mean curvature flow with surgery [HS09]; these have different advantages and are to some extent complementary. Mean curvature flow with surgery gives more refined control on the topology of the evolving manifold, and can also be used to give smooth approximations to weak solutions [Hea13, Lau13]; moreover, it has the nice feature that the entire discussion takes place in the framework of smooth differential geometry.

Roughly speaking, the idea of surgery is to continue the flow through singularities by cutting along necks, gluing in caps, and continuing the flow of the pieces; components of known geometry and topology are discarded. Huisken and Sinestrari successfully implemented this idea for the mean curvature flow of 2-convex hypersurfaces in \mathbb{R}^N ($N \ge 4$), i.e. for hypersurfaces where the sum of the smallest two principal curvatures is positive. As in the construction of 3d Ricci flow with surgery by Perelman [Per02, Per03] (see also [Ham97, KL08, MT07, CZ06, BBB⁺10]), this required considerable technical virtuosity, and was the culmination of a series of long papers [HS99a, HS99b, HS09].

1.2. **Overview.** The purpose of the present paper is to give a new proof for the existence of mean curvature flow with surgery for 2-convex hypersurfaces in \mathbb{R}^N ($N \geq 3$), as announced in [HK13a]. The main focus is on shortness, brevity and simplicity. Moreover, our proof works for mean convex surfaces in \mathbb{R}^3 , which was left as an open problem after [HS09], and recently solved also by Brendle-Huisken [BH13], see Remark 1.28. Furthermore, anticipating future applications, we derive our a priori estimates in a completely local and very flexible setting.

Our new approach to surgery is motivated by our paper [HK13a], where we gave a streamlined treatment of White's theory of mean convex level set (Brakke) flow [Whi00, Whi03, Whi11], based on the beautiful noncollapsing result of Andrews [And12]: Given any $\alpha > 0$, the condition that each boundary point of a mean convex domain admits interior and exterior balls of radius $\alpha/H(p)$ is preserved under the flow.

We formulate the existence theory for a class of flows that we call $(\alpha, \delta, \mathbb{H})$ -flows. The parameters $\alpha = (\alpha, \beta, \gamma)$ measure certain geometric quantities of the initial domain (α -Andrews, $\lambda_1 + \lambda_2 \geq \beta H$, $H \leq \gamma$), the parameter δ measures the quality of the surgery-necks, and the three curvature-scales $\mathbb{H} = (H_{\text{trig}}, H_{\text{neck}}, H_{\text{th}})$ are used to specify more precisely when and how surgeries are performed, see Definition 1.17.

Our main existence result, Theorem 1.21, proves the existence of an $(\alpha, \delta, \mathbb{H})$ -flow for any 2-convex initial domain with parameters α . It is complemented by a canonical neighborhood theorem, Theorem 1.22, which gives a precise geometric description of the regions of high curvature. Other geometric, topological and analytic conclusions follow immediately, see Corollary 1.25, Corollary 1.26, and Proposition 1.27.

The key for our existence proof are new a priori estimates for mean curvature flow with surgery. We formulate our a priori estimates for a more general class of flows that we call (α, δ) -flows. They are concatenations of smooth α -Andrews flows, where at finitely many times: (1) some surgeries are performed on δ -necks of comparable scales; and/or (2) some connected components are discarded, see Definiton 1.3.

Our three main a priori estimates for (α, δ) -flows are a local curvature estimate (Theorem 1.6), a convexity estimate (Theorem 1.8), and a global curvature estimate (Theorem 1.10). These estimates generalize to (α, δ) -flows our main estimates for α -Andrews flows from [HK13a]. The presence of surgeries makes the proof of the local curvature estimate significantly more delicate and requires many new ideas, see Section 2. In the proof of the convexity estimate and the global curvature estimate, we then follow closely the scheme from our previous paper [HK13a], highlighting the new steps necessitated by surgeries.

Broadly speaking, once the a-priori estimates are established, the canonical neighborhood theorem follows from a short argument by contradiction, see Section 4.1, and the main existence result follows from a relatively short continuity argument, see Section 4.2 (strictly speaking, we also need certain structural results for ancient solutions and the evolution of standard caps, but their proofs are easy; see Section 3).

Finally, we point out that the notion of (α, δ) -flows is considerably more flexible than the one of $(\alpha, \delta, \mathbb{H})$ -flows. To a certain extent this additional flexibility is needed for locality, see Remark 1.5. It also makes the class of flows larger, and thus the a priori estimates stronger. 1.3. A priori estimates. We will now discuss our a priori estimates for (α, δ) -flows. We start by recalling the simpler notion of α -Andrews flows, which was the framework for our estimates in [HK13a].

Definition 1.1 (c.f. [And12] and [HK13a, Def. 1.1]). Let $\alpha > 0$. A smooth α -Andrews flow $\{K_t \subseteq U\}_{t \in I}$ in an open set $U \subseteq \mathbb{R}^N$ over a time interval $I \subseteq \mathbb{R}$ is a smooth family of mean convex domains moving by mean curvature flow, such that for every $p \in \partial K_t$ the two closed balls \bar{B}_{int} and \bar{B}_{ext} that are tangent at p and have radius $\frac{\alpha}{H(p)}$ satisfy $\bar{B}_{int} \cap U \subseteq K_t$ and $\bar{B}_{ext} \cap U \subseteq U \setminus Int(K_t)$, respectively.

Convention 1.2. We now fix a factor $\mu \in [1,\infty)$, quantifying the notion of surgeries at comparable scales, for the class of all flows under consideration. Note however, that we do not fix any absolute scale.

Definition 1.3. An (α, δ) -flow \mathcal{K} is a collection of finitely many smooth α -Andrews flows $\{K_t^i \subseteq U\}_{t \in [t_{i-1}, t_i]}$ $(i = 1, \dots, k; t_0 < \dots < t_k)$ in an open set $U \subseteq \mathbb{R}^N$ (see Definition 1.1), such that:

- (1) for each $i = 1, \ldots, k-1$, the final time slices of some collection of disjoint strong δ -necks¹ are replaced by pairs of standard caps as described in Definition 2.4, giving a domain $K_{t_i}^{\sharp} \subseteq K_{t_i}^i =: K_{t_i}^-$.
- (2) the initial time slice of the next flow, Kⁱ⁺¹_{ti} =: K⁺_{ti}, is obtained from K[♯]_{ti} by discarding some connected components.
 (3) there exists s[♯] = s[♯](K) > 0, which depends on K, such that all
- necks in item (1) have radius $s \in [\mu^{-1/2} s_{\sharp}, \mu^{1/2} s_{\sharp}]$.

Remark 1.4. To avoid confusion, we emphasize that the word 'some' allows for the empty set, i.e. some of the inclusions $K_{t_i}^+ \subseteq K_{t_i}^{\sharp} \subseteq K_{t_i}^-$ could actually be equalities. In other words, there can be some times t_i where effectively only one of the steps (1) or (2) is carried out. Also, the flow can become extinct, i.e. we allow the possibility that $K_{t_i}^{i+1} = \emptyset$.

Remark 1.5. At first sight, the possibility that some connected components can be discarded whenever someone wants might seem arbitrary, but in fact this flexibility is crucial for obtaining local estimates. For example, imagine the situation that $\{L_t \subset \mathbb{R}^N\}_{t \in I}$ is a mean curvature flow with surgery in \mathbb{R}^N (technically an $(\alpha, \delta, \mathbb{H})$ -flow), and that we want to prove local estimates in an open set $U \subset \mathbb{R}^N$, i.e. estimates for the (α, δ) -flow $\{K_t = L_t \cap U \subseteq U\}_{t \in I'}$, on some interval $I' \subseteq I$. It can of course happen that there is a surgery in L_t on a neck outside U that disconnects L_t into two connected components. According to the standard procedure, the component of high curvature, which has

¹More precisely, we only require that their final time-slices are disjoint in U.

known geometry and topology, is then discarded. Viewed from U one cannot see the neck and thus one observes a component being discarded seemingly on a whim.

We will now state our three main a priori estimates for (α, δ) -flows. We say that an (α, δ) -flow is defined in a parabolic ball $P(p, t, r) = B(p, r) \times (t - r^2, t]$ if it is defined in an open set $U \supseteq B(p, r)$ and for all $t' \in (t - r^2, t]$. Our first main estimate gives curvature control on a whole parabolic ball, from a mean curvature bound at a single point.

Theorem 1.6 (Local curvature estimate). There exist $\delta = \delta(\alpha) > 0$, $\rho = \rho(\alpha) > 0$ and $C_{\ell} = C_{\ell}(\alpha) < \infty$ ($\ell = 0, 1, 2, ...$) with the following property. If \mathcal{K} is an (α, δ) -flow ($\delta \leq \overline{\delta}$) in a parabolic ball P(p, t, r)centered at a boundary point $p \in \partial K_t$ with $H(p, t) \leq r^{-1}$, then

(1.7)
$$\sup_{P(p,t,\rho r) \cap \partial \mathcal{K}} |\nabla^{\ell} A| \le C_{\ell} r^{-1-\ell}$$

Second, we have the convexity estimate for (α, δ) -flows.

Theorem 1.8 (Convexity estimate). For all $\varepsilon > 0$, there exist $\bar{\delta} = \bar{\delta}(\alpha) > 0$ and $\eta = \eta(\varepsilon, \alpha) < \infty$ with the following property. If \mathcal{K} is an (α, δ) -flow $(\delta \leq \bar{\delta})$ defined in a parabolic ball $P(p, t, \eta r)$ centered at a boundary point $p \in \partial K_t$ with $H(p, t) \leq r^{-1}$, then

(1.9)
$$\lambda_1(p,t) \ge -\varepsilon r^{-1}.$$

Theorem 1.8 says that a boundary point (p, t) in an (α, δ) -flow has almost positive definite second fundamental form as long as the flow has had a chance to evolve over a portion of space-time which is large compared with the scale given by H(p, t). In particular, it implies that ancient (α, δ) -flows in \mathbb{R}^N have nonnegative second fundamental form.

Third, we have the global curvature estimate for (α, δ) -flows.

Theorem 1.10 (Global curvature estimate). For all $\Lambda < \infty$, there exist $\bar{\delta} = \bar{\delta}(\alpha) > 0$, $\eta = \eta(\alpha, \Lambda) < \infty$ and $C_{\ell} = C_{\ell}(\alpha, \Lambda) < \infty$ ($\ell = 0, 1, 2, ...$) with the following property. If \mathcal{K} is an (α, δ) -flow $(\delta \leq \bar{\delta})$ in a parabolic ball $P(p, t, \eta r)$ centered at a boundary point $p \in \partial K_t$ with $H(p, t) \leq r^{-1}$, then

(1.11)
$$\sup_{P(p,t,\Lambda r)\cap\partial\mathcal{K}'} |\nabla^{\ell}A| \le C_{\ell} r^{-1-\ell},$$

where \mathcal{K}' denotes the (α, δ) -flow in $P(p, t, \Lambda r)$ that is obtained from \mathcal{K} via restricting to $B(p, \Lambda r)$ and discarding the connected components that do not contain p.

Remark 1.12. Theorem 1.10 quickly implies (and immediately follows from) a global convergence result, Corollary 2.30.

Remark 1.13. Only normalizing the curvature at the basepoint one cannot get curvature control for other connected components. An interesting feature, that shows up later in the proof of the canonical neighborhood theorem for $(\alpha, \delta, \mathbb{H})$ -flows (Section 4.1), is that the Andrews condition in combination with the degeneration of the surgery parameters provides a mechanism to clear out other components.

Remark 1.14. Our a priori estimates are local and thus also apply if mean convexity and the Andrews condition only hold locally around a singularity. This seems to be very useful if one wants to implement surgery for generic mean curvature flow, c.f. [CM12, CIM13, CM13].

1.4. Existence theory. We now turn to the discussion of the existence theory. Our construction only depends on a few parameters of the initial domain, which is always assumed to be 2-convex.

Definition 1.15 (Controlled initial condition). Let $\alpha = (\alpha, \beta, \gamma) \in (0, N-2) \times (0, \frac{1}{N-2}) \times (0, \infty)$. A smooth compact domain $K_0 \subset \mathbb{R}^N$ is called an α -controlled initial condition, if it satisfies the α -Andrews condition and the inequalities $\lambda_1 + \lambda_2 \geq \beta H$ and $H \leq \gamma$.

Remark 1.16. Note that every smooth compact 2-convex domain is a controlled initial condition for some parameters $\alpha, \beta, \gamma > 0$.

For us, a mean curvature flow with surgery is a (α, δ) -flow with (α, β, γ) -controlled initial data, subject to the following additional conditions. First, the flow is β -uniformly 2-convex, i.e. $\lambda_1 + \lambda_2 \geq \beta H$. Besides the neck-quality $\delta > 0$, we have three curvature-scales $H_{\text{trig}} > H_{\text{neck}} > H_{\text{th}} > 1$, to which we refer as the trigger-, neck- and thick-curvature. The surgeries are done at times t when the maximum of the mean curvature hits H_{trig} . They are performed on a minimal disjoint collection of solid δ -necks of curvature H_{neck} that separate the trigger part $\{H = H_{\text{trig}}\}$ from the thick part $\{H \leq H_{\text{th}}\}$ in K_t^- , and the high curvature components are discarded. Finally, we impose the condition that surgeries are done more and more precisely, if the surgery-necks happen to be rounder and rounder. We call our flows with surgery $(\alpha, \delta, \mathbb{H})$ -flows, and the precise definition is as follows.

Definition 1.17. An $(\alpha, \delta, \mathbb{H})$ -flow, $\mathbb{H} = (H_{\text{th}}, H_{\text{neck}}, H_{\text{trig}})$, is an (α, δ) flow $\{K_t \subset \mathbb{R}^N\}_{t\geq 0}$ (see Definition 1.3) with $\lambda_1 + \lambda_2 \geq \beta H$, and with $\alpha = (\alpha, \beta, \gamma)$ -controlled initial condition $K_0 \subset \mathbb{R}^N$ (see Definition 1.15) such that

 $\mathbf{6}$

- (1) $H \leq H_{\text{trig}}$ everywhere, and surgery and/or discarding occurs precisely at times t when $H = H_{\text{trig}}$ somewhere.
- (2) The collection of necks in item (1) of Definition 1.3 is a minimal collection of solid δ -necks of curvature H_{neck} which separate the set $\{H = H_{\text{trig}}\}$ from $\{H \leq H_{\text{th}}\}$ in the domain K_t^- .
- (3) K_t^+ is obtained from K_t^{\sharp} by discarding precisely those connected components with $H > H_{\text{th}}$ everywhere. In particular, of each pair of facing surgery caps precisely one is discarded.
- (4) If a strong δ -neck from item (2) also is a strong δ -neck for some $\hat{\delta} < \delta$, then property (4) of Definition 2.4 also holds with $\hat{\delta}$ instead of δ .

Remark 1.18. To avoid confusion, we emphasize again that the collection of necks in item (2) can be empty. Indeed, the simplest example of an $(\alpha, \delta, \mathbb{H})$ -flow is the evolution of a round ball $K_0 = \bar{B}^N$, which just shrinks self-similarly until H reaches H_{trig} , and then is discarded.

Remark 1.19. By comparison every $(\alpha, \delta, \mathbb{H})$ -flow becomes extinct after some finite time T, i.e. satisfies $K_t = \emptyset$ for all t > T.

Remark 1.20. We point out that our definition of $(\alpha, \delta, \mathbb{H})$ -flows does not include a canonical neighborhood property. Instead, we will prove that the canonical neighborhood property is consequence of the global convergence result (Corollary 2.30), see Theorem 1.22.

Our main existence theorem is the following.

Theorem 1.21 (Existence of mean curvature flow with surgery). There are constants $\overline{\delta} = \overline{\delta}(\alpha) > 0$ and $\Theta(\delta) = \Theta(\alpha, \delta) < \infty$ ($\delta \leq \overline{\delta}$) with the following significance. If $\delta \leq \overline{\delta}$ and $\mathbb{H} = (H_{trig}, H_{neck}, H_{th})$ are positive numbers with $H_{trig}/H_{neck}, H_{neck}/H_{th}, H_{neck} \geq \Theta(\delta)$, then there exists an $(\alpha, \delta, \mathbb{H})$ -flow $\{K_t\}_{t \in [0,\infty)}$ for every α -controlled initial condition K_0 .

Theorem 1.21 enables us to evolve any smooth compact 2-convex domain by mean curvature flow with surgery. More generally, it also enables us evolve (finite- or infinite-dimensional) families of α -controlled initial domains simultaneously, with uniform estimates.

Our existence theorem (Theorem 1.21) is complemented by the following canonical neighborhood theorem (we use the terminology of Perelman [Per03], but our canonical neighborhoods are different), which gives precise geometric information about the regions of high curvature.

Theorem 1.22 (Canonical neighborhood theorem). For all $\varepsilon > 0$, there exist $\overline{\delta} = \overline{\delta}(\alpha) > 0$, $H_{can}(\varepsilon) = H_{can}(\alpha, \varepsilon) < \infty$ and $\Theta_{\varepsilon}(\delta) =$ $\Theta_{\varepsilon}(\alpha, \delta) < \infty$ ($\delta \leq \overline{\delta}$) with the following significance. If $\delta \leq \overline{\delta}$ and \mathcal{K} is an $(\alpha, \delta, \mathbb{H})$ -flow with $H_{trig}/H_{neck}, H_{neck}/H_{th} \geq \Theta_{\varepsilon}(\delta)$, then any $(p,t) \in \partial \mathcal{K}$ with $H(p,t) \geq H_{can}(\varepsilon)$ is ε -close to either (a) a β -uniformly 2-convex ancient α -Andrews flow, or (b) the evolution of a standard cap preceded by the evolution of a round cylinder.

Remark 1.23. The structure of uniformly 2-convex ancient α -Andrews flows and the standard solution are discussed in Section 3.

Remark 1.24. In practice, one can first choose $\varepsilon > 0$ small enough, and then $H_{\text{trig}}/H_{\text{neck}}, H_{\text{neck}}/H_{\text{th}}, H_{\text{neck}} \geq \max\{\Theta_{\varepsilon}(\delta), \Theta(\delta)\} \ (\delta \leq \overline{\delta})$ and $H_{\text{th}} \geq H_{\text{can}}(\varepsilon)$, where $\overline{\delta} = \min\{\overline{\delta}_{1,21}, \overline{\delta}_{1,22}\}$, so that both the existence result (Theorem 1.21) and the canonical neighborhood property at any point with $H(p,t) \geq H_{\text{can}}(\varepsilon)$ (Theorem 1.22) are applicable.

In particular, $(\alpha, \delta, \mathbb{H})$ -flows imply a decomposition of the initial domains into balls \overline{D}^N and solid tori $\overline{D}^{N-1} \times S^1$.

Corollary 1.25 (Discarded components). For $\varepsilon > 0$ small enough, for any $(\alpha, \delta, \mathbb{H})$ -flow with $H_{trig}/H_{neck}, H_{neck}/H_{th} \ge \Theta_{\varepsilon}(\delta)$ ($\delta \le \overline{\delta}$) and $H_{th} \ge H_{can}(\varepsilon)$, where $\Theta_{\varepsilon}(\delta)$, $\overline{\delta}$ and $H_{can}(\varepsilon)$ are from Theorem 1.22, all discarded components are diffeomorphic to \overline{D}^N or $\overline{D}^{N-1} \times S^1$.

Corollary 1.26. Any smooth compact 2-convex domain in \mathbb{R}^N is diffeomorphic to a connected sum of finitely many solid tori $\overline{D}^{N-1} \times S^1$.

Finally, by the work of Lauer [Lau13] we see that sequences of flows with surgery converge to the level set flow [ES91, CGG91], if the initial domain is kept fixed and the thick curvatures tend to infinity.

Proposition 1.27 (Convergence to level set flow). There exists $\overline{\delta} > 0$ such that if $\mathcal{K}^j = \{K_t^j\}_{t \in [0,\infty)}$ is a sequence of $(\alpha, \delta_j, \mathbb{H}_j)$ -flows $(\delta_j \leq \overline{\delta})$ starting at a fixed initial domain K_0 , with $H_{th}^j \to \infty$, then \mathcal{K}^j Hausdorff converges in $\mathbb{R}^N \times [0,\infty)$ to \mathcal{K} , the level set flow of K_0 .

Proposition 1.27 enables us to approximate level set flows with α controlled initial conditions by $(\alpha, \delta, \mathbb{H})$ -flows. In fact, this approximation works uniformly for families of α -controlled initial conditions.

Remark 1.28. We have announced our new construction of mean curvature flow with surgery in April 2013, see [HK13a]. In September 2013, Brendle and Huisken posted a very interesting preprint [BH13], where they also solved the case N = 3, by combining the work of Huisken-Sinestrari [HS09], the local curvature estimate from our previous paper [HK13a], and Brendle's estimate for the inscribed radius [Bre13a]. Although Brendle's estimate is not needed at all for our approach to surgery, we have written a note [HK13b] giving a shorter proof of it. Also, Brendle's interesting variant of the monotonicity formula [Bre13b] is somewhat related to Claim 2.17 in the present paper.

Organization of the paper. In Section 2, we prove our a priori estimates for (α, δ) -flows. In Section 3, we prove some results for ancient solutions and standard solutions, that are needed later. In Section 4, we prove the canonical neighborhood theorem and the main existence theorem.

2. A priori estimates for flows with surgeries

Our a priori estimates hold for flows in any open $U \subseteq \mathbb{R}^N$. Working locally necessitates a number of extra technicalities in the definitions and proofs; readers focussing on the global case can assume $U = \mathbb{R}^N$.

2.1. **Basic properties of** (α, δ) -flows. We will now discuss some basic properties of (α, δ) -flows (Definition 1.3). We start by defining what it means to replace a neck by caps, making explicit the constants that are involved; this notion is used in item (1) of Definition 1.3.

Convention 2.1. In the following, $\mu < \infty$, $C_{\ell} < \infty$ ($\ell = 0, 1, 2, ...$) and $\delta'(\delta)$ with δ' decreasing to zero as $\delta \searrow 0$, are fixed but arbitrary. Strictly speaking, the specification of μ , C_{ℓ} and $\delta'(\delta)$ is part of the definition of an (α, δ) -flow. Other constants may depend on μ , C_{ℓ} and $\delta'(\delta)$, but for brevity we suppress that in the notation. Also, $\Gamma \in [10, \infty)$ denotes a constant to be fixed later (Convention 2.11).

Definition 2.2 (Standard cap). A standard cap is a smooth convex domain $K^{\text{st}} \subset \mathbb{R}^N$ that coincides with a solid round half-cylinder of radius 1 outside a ball of radius 10.

Definition 2.3 (Strong δ -neck). We say that an (α, δ) -flow $\mathcal{K} = \{K_t \subseteq U\}_{t \in I}$ has a strong δ -neck with center p and radius s at time $t_0 \in I$, if $\{s^{-1} \cdot (K_{t_0+s^2t}-p)\}_{t \in (-1,0]}$ is δ -close in $C^{\lfloor 1/\delta \rfloor}$ in $B^U_{1/\delta} \times (-1,0]$ to the evolution of a solid round cylinder $\overline{D}^{N-1} \times \mathbb{R}$ with radius 1 at t = 0, where $B^U_{1/\delta} = s^{-1} \cdot ((B(p, s/\delta) \cap U) - p) \subseteq B(0, 1/\delta) \subset \mathbb{R}^N$.

Definition 2.4 (Replacing a δ -neck by standard caps). We say that the final time slice of a strong δ -neck ($\delta \leq \frac{1}{10\Gamma}$) with center p and radius s is replaced by a pair of standard caps, if the pre-surgery domain $K^- \subseteq U$ is replaced by a post-surgery domain $K^{\sharp} \subseteq K^-$ such that:

(1) the modification takes places inside a ball $B = B(p, 5\Gamma s)$.

(2) there are bounds for the second fundamental form and its derivatives:

$$\sup_{\partial K^{\sharp} \cap B} |\nabla^{\ell} A| \le C_{\ell} s^{-1-\ell} \qquad (\ell = 0, 1, 2, \ldots).$$

- (3) if $B \subseteq U$, then for every point $p_{\sharp} \in \partial K^{\sharp} \cap B$ with $\lambda_1(p_{\sharp}) < 0$, there is a point $p_- \in \partial K^- \cap B$ with $\frac{\lambda_1}{H}(p_-) \leq \frac{\lambda_1}{H}(p_{\sharp})$.
- (4) if $B(p, 10\Gamma s) \subseteq U$, then $s^{-1} \cdot (K^{\sharp} p)$ is $\delta'(\delta)$ -close in $B(0, 10\Gamma)$ to a pair of disjoint standard caps, that are at distance Γ from the origin (see Convention 2.1 and Definition 2.2).

Proposition 2.5 (Spatial separation of surgeries). For every $\zeta < \infty$, there are constants $\overline{\delta} = \overline{\delta}(\zeta) > 0$ and $\eta = \eta(\zeta) < \infty$ with the following property. If \mathcal{K} is an (α, δ) -flow $(\delta \leq \overline{\delta})$, p is the center of a surgery neck of radius s, and \mathcal{K} is defined in an open set that contains the ball $B(p, \eta s)$, then are no other surgeries in $B(p, \zeta s)$.

Proof. Let p' be the center of another surgery neck. If the surgery occurs at the same time t' = t, then by disjointness of the collection of δ -necks in $B(p, \eta s)$, we get $d(p, p') \geq \min\{\delta^{-1}, \eta\}s$. If t' > t, then by monotonicity of sets we have $K_{t'} \subseteq K_t$, and we also know that K_t is close to a pair of standard caps near p. Since a δ -neck can only be contained in a standard cap when the center is far away from the tip, we obtain that $d(p, p') \geq \frac{1}{2}\mu^{-1}\min\{\delta^{-1}, \eta\}s$ for η large and δ small enough. Finally, if t' < t, we can apply the same argument with the roles of p and p' reversed. The assertion follows.

Definition 2.6 (Points modified by surgery). We say that an open set *B* contains *points modified by surgery* at time *t* if $(K_t^- \setminus K_t^{\sharp}) \cap B \neq \emptyset$.

Proposition 2.7 (Forward estimate after surgery). For all $\alpha > 0$, there exist $\varepsilon = \varepsilon(\alpha) > 0$, $C = C(\alpha) < \infty$ and $\overline{\delta} = \overline{\delta}(\alpha) > 0$ with the following property. If \mathcal{K} as an (α, δ) -flow $(\delta \leq \overline{\delta})$, and $r \leq s$ and $t_1 < t_2$ are such the flow is defined in $B(p, r) \times [t_1, t_2]$, with some point in B(p, r) modified by a surgery at scale s at time t_1 , but no points in B(p, r) modified by surgeries for $t \in (t_1, t_2]$, then

(2.8)
$$\sup_{\partial K_t \cap B(p,r/2)} |A| \le Cr^{-1} \quad on \quad [t_1, \min\{t_1 + (\varepsilon r)^2, t_2\}].$$

Proof. By Definition 2.4 and the Andrews-condition we have uniformly controlled geometry in B(p, r) at time t_1 (the curvature bounds follow from item (2) for $\partial K_{t_1}^{\sharp} \cap B(p, r)$, and from being part of a strong δ -neck

10

at other points). Thus, the assertion follows from the pseudolocality theorem for the mean curvature flow, see e.g. [CY07, Thm. 7.5].² \Box

Proposition 2.9 (One-sided minimization). There exist $\bar{\delta} > 0$ and $\Gamma_0 < \infty$ with the following property. If \mathcal{K} is an (α, δ) -flow $(\delta \leq \bar{\delta})$ in an open set U, with cap separation parameter $\Gamma \geq \Gamma_0$ and surgeries at scales between $\mu^{-1}s$ and s, and if $\bar{B} \subset U$ is a closed ball with $d(\bar{B}, \mathbb{R}^N \setminus U) \geq 20\Gamma s$, then

$$(2.10) \qquad \qquad |\partial K_{t_1} \cap \bar{B}| \le |\partial K' \cap \bar{B}|$$

for every smooth comparison domain K' that agrees with K_{t_1} outside \overline{B} and satisfies $K_{t_1} \subseteq K' \subseteq K_{t_0}$ for some $t_0 < t_1$.

Proof. This follows using the geometric measure theory argument in [Whi00, Sec. 3] and [Hea13, Sec. 5]; in Appendix A we give a detailed exposition, and also an alternative elementary argument. \Box

Convention 2.11. We now fix a constant $\Gamma \ge \max{\{\Gamma_0, 10\}}$, where Γ_0 is from Proposition 2.9, that is also large enough such that there are constants $\omega < 1$ and $\overline{\delta} > 0$ with

 $(2.12) \qquad \qquad |\partial K^+ \cap B| \le \omega |\partial K^- \cap B|,$

for every surgery on a δ -neck ($\delta \leq \overline{\delta}$) with $B = B(p, 5\Gamma s) \subseteq U$.

2.2. Local curvature estimate. We will now prove Theorem 1.6. Since the proof is somewhat involved, we first give a detailed outline.

Outline of the proof. Arguing by contradiction, we get a sequence of flows on larger and larger parabolic balls where the curvature goes to zero at the basepoint but blows up at some nearby point. We first split off the two easy cases that there are no nearby surgeries (Case 1) or surgeries at macroscopic scales (Case 2), which can be dealt with by applying the local curvature estimate from our previous paper [HK13a, Thm. 1.8] and the forward estimate after surgeries (Proposition 2.7), respectively. The core of the proof is then to rule out surgeries at microscopic scales (Case 3). We do this as follows: At a surgery neck the value of the Huisken density is close to the value of the cylinder. However, since the mean curvature at the basepoint goes to zero, using a halfspace convergence argument (Claim 2.24) and one-sided minimization (Proposition 2.9), we can show that the Huisken density is close to 1 further back in time. Finally, analyzing the contributions from surgeries in different regimes, c.f. (2.18), we prove that the cumulative

²Note that dropping connected components has the good sign in Huisken's monotonicity inequality. Thus, their proof of pseudolocality goes through in our setting.

error in Huisken's monotonicity inequality due to surgeries goes to zero (Claim 2.17), and conclude that microscopic surgeries cannot occur.

Proof of Theorem 1.6. As in [HK13a, Sec. 2], we will first prove (1.7) under the admissibility assumption that some time slice $K_{\bar{t}}$ contains B(p,r). Assume towards a contradiction that there exists a sequence \mathcal{K}^{j} of admissible (α, j^{-1}) -flows in P(0, 0, j) with $H(0, 0) \leq j^{-1}$, but such that

(2.13)
$$\sup_{P(0,0,j^{-2})} |A| \ge j^2.$$

Case 1: there are no points modified by surgeries in $P(0, 0, j^{-1})$, for infinitely many j. In this case, we can apply the local curvature estimate from our previous paper [HK13a, Thm. 1.8, Rmk. 2.12] (see also Remark 2.25 below for an alternative proof). For large j this gives a contradiction with (2.13).

Case 2: there is a point in $P(0, 0, j^{-1})$ modified by a surgery at scale $s_j \geq \mu \varepsilon^{-1} j^{-1}$, for infinitely many j, where ε is the constant from Proposition 2.7. Let t_1^j be the largest $t \in (-j^{-2}, 0]$ such that there is a point in $B(0, \varepsilon^{-1} j^{-1})$ modified by a surgery. Using j^{-1} -closeness to a strong neck we get curvature estimates prior to t_1^j , and using Proposition 2.7 we get curvature estimates after t_1^j . For large j this gives a contradiction with (2.13).

Case 3: there is a point in $P(0, 0, j^{-1})$ modified by a surgery, and all surgeries are at scale $s_j \leq \mu^2 \varepsilon^{-1} j^{-1}$, for large j. We will show that this case actually cannot occur. Suppressing j in the notation, let x_0 be the center of neck coming from a surgery in $P(0, 0, j^{-1})$, at time t and scale s. Let $t_0 = t + \frac{s^2}{2(N-2)}$, and consider the Huisken density

(2.14)
$$\Theta(\tau) = \int_{\partial K_{t_0-\tau}} \theta^{\sigma}_{X_0} dA,$$

based at $X_0 = (x_0, t_0)$. Here, the integrand $\theta_{X_0}^{\sigma}$ is the backwards heat kernel times a suitable cutoff function at scale $\sigma \gg 1$ (as always, we tacitly assume that j is large enough), namely (2.15)

$$\theta_{X_0}^{\sigma}(x,t) = \left(4\pi(t_0-t)\right)^{-(N-1)/2} e^{-\frac{|x-x_0|^2}{4(t_0-t)}} \left(1 - \frac{|x-x_0|^2 + 2(N-1)(t-t_0)}{\sigma^2}\right)_+^3.$$

Note that for small backwards time, say $\tau = s^2$, we have

(2.16)
$$\liminf_{j \to \infty} \Theta(s^2) \ge \Theta_{N-1} > 1,$$

12

where Θ_{N-1} is the density of the cylinder $\overline{D}^{N-1} \times \mathbb{R}$. Recall that by Huisken's monotonicity inequality, see e.g. [Eck04, Prop. 4.17] or [HK13a, App. B], the function $\Theta(\tau)$ is monotone if there are no surgeries (throwing away connected components has the good sign).

Claim 2.17. The cumulative error in Huisken's monotonicity inequality due to surgeries between $\tau = s^2$ and $\tau = 1$ goes to zero as $j \to \infty$.

Proof. Let $\varepsilon > 0$, and write $\sigma_t = \sqrt{\sigma^2 - 2(N-1)(t-t_0)}$. We say that a surgery center x_i at time t_i is in the nonoscillating regime, if

(2.18) (a)
$$\frac{s|x_i - x_0|}{|t_i - t_0|} < \varepsilon$$
 and (b) $||x_i - x_0| - \sigma_{t_i}| > \varepsilon^{-1}s$.

If ε is sufficiently small, then the change of the Huisken density due to any surgery (x_i, t_i) in the nonoscillating regime has the good sign, i.e.

(2.19)
$$\int_{\partial K_{t_i}^{\sharp} \cap B_i} \theta_{X_0}^{\sigma} dA \le \int_{\partial K_{t_i}^- \cap B_i} \theta_{X_0}^{\sigma} dA.$$

where $B_i = B(x_i, 5\Gamma s_i)$. Indeed, this follows from the fact that area decreases by a definite factor under surgery, see (2.12), and the observation that we can make the ratio between $\sup_{x \in B_i} \theta^{\sigma}_{X_0}(x, t_i)$ and $\inf_{x \in B_i} \theta^{\sigma}_{X_0}(x, t_i)$ as close to 1 as we want, by choosing ε small enough.

We next estimate the cumulative error due to the surgeries (x_i, t_i) violating (a). Taking into account Proposition 2.5, it suffices to estimate the sum

(2.20)
$$\sum_{i} \frac{1}{\tau_{i}^{(N-1)/2}} e^{-|x_{i}-x_{0}|^{2}/5\tau_{i}} A_{i},$$

where $\tau_i = t_0 - t_i$ and A_i is the area of the region modified by the surgery. Here, we used that $|x - x_0|^2 \ge \frac{4}{5}|x_i - x_0|^2$ for x in the region around x_i modified by surgery, for j large. To estimate (2.20), we first pull out a factor $e^{-|x_i-x_0|^2/10\tau_i}$. Using Proposition 2.5 again, we observe that (for j large enough) the minimum of $\frac{|x_i-x_0|^2}{10\tau_i} + \frac{N-1}{2}\log \tau_i$ over τ_i under the constraint $\tau_i \le \frac{s|x_i-x_0|}{\varepsilon}$ is attained at $\tau_i = \frac{s|x_i-x_0|}{\varepsilon}$. Thus (2.21)

$$\sum_{i} \frac{1}{\tau_{i}^{(N-1)/2}} e^{-\frac{|x_{i}-x_{0}|^{2}}{5\tau_{i}}} A_{i} \leq \delta_{j} \sum_{i} \left(\frac{\varepsilon}{s|x_{i}-x_{0}|}\right)^{(N-1)/2} e^{-\frac{\varepsilon|x_{i}-x_{0}|}{10s}} A_{i},$$

where $\delta_j := \sup_i e^{-|x_i - x_0|^2/10\tau_i}$ goes to zero as $j \to \infty$, since

(2.22)
$$\frac{|x_i - x_0|^2}{\tau_i} = \frac{s|x_i - x_0|}{\tau_i} \frac{|x_i - x_0|}{s} \ge \varepsilon \zeta_j \to \infty,$$

again by Proposition 2.5. Since the regions modified by surgeries are separated by a large multiple of s by Proposition 2.5, and have area comparable to s^{N-1} by Definition 2.4, the sum on the right hand side of (2.21) can be uniformly estimated by a multiple of

(2.23)
$$\int_{s}^{\infty} \frac{1}{(sR)^{(N-1)/2}} e^{-R/s} R^{N-2} dR = \int_{1}^{\infty} u^{(N-3)/2} e^{-u} du < \infty.$$

Thus, the cumulative error due to surgeries violating (a) goes to zero as $j \to \infty$.

Finally, observing that $\theta_{X_0}^{\sigma}(x,t) \leq s^3$ in the relevant region, the cumulative error due to the surgeries violating (b) can be estimated by a multiple of $s^3(\frac{\sigma}{s})^N s^{N-1} \to 0$ $(j \to \infty)$. This proves the claim. \Box

Claim 2.24. After rotating coordinates, the sequence \mathcal{K}^j converges in the pointed Hausdorff topology on $\mathbb{R}^N \times (-\infty, 0]$ to the static halfspace $\{x_N \leq 0\} \times (-\infty, 0]$, and similarly for the complements.

Proof. We rotate coordinates such that the outward unit normal of K_0^j at (0,0) is e_N . Since $H(0,0) \leq j^{-1}$, the Andrews condition implies in particular that every compact subset of the lower half plane $\{x_N < 0\}$ is contained in K_0^j for j large enough. The proof is now almost verbatim as the one of [HK13a, Claim 2.4], taking into account the following caveat. In our previous proof we used comparison from the interior with the mean curvature evolution of a ball \bar{B}_R^d . In general, interior comparison can fail for (α, δ) -flows, since there can be jumps due to surgeries and thrown away components. However, in our situation the radius of the comparison ball is much larger than the surgery scale $s_j \to 0$. Thus, the comparison ball must be disjoint from any region modified by surgeries, since any such region is contained in a long and thin neck. Since the comparison ball contains the origin, it cannot be contained in a discarded component either.

Finishing the discussion of Case 3, it follows from the halfspace convergence (Claim 2.24) and the one-sided minimization (Proposition 2.9) that at $\tau = 1$ the value of the Huisken density is close to 1; together with the almost monotonicity (Claim 2.17) this contradicts (2.16).

Finally, the admissibility assumption can be removed as in [HK13a, App. D],³ and replacing ρ by $\rho/2$, the bounds for the derivatives of A follow from standard interior estimates.

³Readers focussing on the admissible version of the estimate can skip this step.

Remark 2.25. A simple nonsurgical variant of the argument in Case 3 gives an alternative proof of [HK13a, Thm. 1.8, Rmk. 2.12]. Namely, given a contradictory sequence \mathcal{K}^j as at the beginning of the proof of Theorem 1.6, select points $X_j \in \partial \mathcal{K}^j \cap P(0, 0, 1/2)$ such that $Q_j = |A|(X_j) \geq j^2$ and $\sup_{P(X_j, jQ_j^{-1})} |A| \leq 2Q_j$. Then, the Huisken density Θ_{X_j} based at X_j satisfies $\liminf_{j\to\infty} \Theta_{X_j}(Q_j^{-2}) > 1$. Arguing as in Case 3, for $\tau = 1$ we are as close as we want to a halfspace; together with the one-sided minimization, this contradicts monotonicity.

2.3. Convexity estimate. Using the local curvature estimate (Theorem 1.6), we can now prove the convexity estimate (Theorem 1.8). The idea is, as in [HK13a, Proof of Thm. 1.8], to consider a contradictory sequence and to pass to a local limit such that $\frac{\lambda_1}{H}$ attains a negative minimum, contradicting the strict maximum principle. To make this idea work in the presence of surgeries, we have to choose the sequence very carefully, using in particular item (3) of Definition 2.4.

Proof. Fix α , and let $\overline{\delta} = \overline{\delta}(\alpha) > 0$ small enough to justify the application of Theorem 1.6 and of the properties of Definition 2.4 in the argument below. The α -Andrews condition implies that for every $\varepsilon \geq \frac{1}{\alpha}$, we can find an $\eta < \infty$ such that the assertion holds. Let $\varepsilon_0 \leq \frac{1}{\alpha}$ be the infimum of the ε 's for which this is possible, and suppose $\varepsilon_0 > 0$.

It follows that there is a sequence $\{\mathcal{K}^j\}$ of (α, δ_j) -flows, $\delta_j \leq \overline{\delta}$, in P(0,0,j) such that $H(0,0) \leq 1$, but $\lambda_1(0,0) \rightarrow -\varepsilon_0$ as $j \rightarrow \infty$. By the choice of ε_0 , it follows that $H(0,0) \rightarrow 1$ as $j \rightarrow \infty$, since otherwise we could parabolically scale our sequence and get a new sequence where $H(0,0) \leq 1$, but $\lambda_1(0,0)$ tends to something strictly smaller than $-\varepsilon_0$.

Let $\rho = \rho(\alpha)$ be the quantity from the local curvature estimate (Theorem 1.6). Then there are uniform bounds on A and its space-time derivatives in $P(0, 0, \rho/2)$. Suppose there is no r > 0 such that the flow is unmodified by surgeries in P(0, 0, r) after passing to a subsequence. In view of the bounds on spacetime derivatives of curvature, we may assume, after translating and parabolic rescaling (by a factors tending to 1 as $j \to \infty$), that t = 0 is a surgery time, and that (0, 0) lies in $\partial K_0^{\sharp} \cap B(p, 5\Gamma s)$, c.f. Definition 2.4. The radius of the surgery neck is comparable to one, again by Definition 2.4. Thus, by item (3) of Definition 2.4, after passing to some point at controlled distance in the presurgery manifold, and parabolically rescaling by factors of controlled size, we may assume that (0, 0) lies in the presurgery manifold ∂K_0^- , H(0, 0) = 1, and $\lambda_1(0, 0) \to -\varepsilon_0$. After modifying the sequence in this way, the argument can now be concluded as in [HK13a, Proof of Thm. 16

1.8]. Namely, using Proposition 2.5 and Theorem 1.6 we get a smooth mean curvature flow \mathcal{K}^{∞} in some parabolic ball P(0,0,r) such that the ratio $\frac{\lambda_1}{H}$ attains a negative minimum $-\varepsilon_0$ at (0,0); this contradicts the strict maximum principle.

2.4. Global curvature estimate. The global convergence theorem from our previous paper ([HK13a, Thm. 1.12]) was based on the local curvature estimate and the convexity estimate. Having established the local curvature estimate and the convexity estimate for (α, δ) -flows (Theorem 1.6 and Theorem 1.8), we will now show that our previous global convergence argument goes through with minor adjustments.

Proof of Theorem 1.10. We choose $\bar{\delta} = \bar{\delta}(\alpha) > 0$ small enough such that the estimates from the previous sections apply. Suppose towards a contradiction, that there is a sequence \mathcal{K}^j of (α, δ_j) -flows $(\delta_j \leq \bar{\delta})$ in $P(0, 0, \eta_j)$, with $\eta_j \to \infty$ and $H(0, 0) \leq 1$, such that

(2.26)
$$\lim_{j \to \infty} \sup_{P(0,0,\Lambda) \cap \partial \mathcal{K}'^j} |A| = \infty,$$

for some $\Lambda < \infty$, where \mathcal{K}'^{j} denotes the (α, δ_{j}) -flow whose time slices are given by the connected component of $K_{t} \cap B(0, \Lambda)$ containing 0.

We can assume that there is some $R < \infty$ such that P(0, 0, R) contains surgeries of \mathcal{K}^j for large j, since otherwise [HK13a, Thm. 1.12, Rmk. 3.5] gives a contradiction with (2.26). Also, it must be the case that the surgery scales $s_j = s_{\sharp}(\mathcal{K}^j)$ (see Definition 1.3) satisfy

(2.27)
$$\limsup_{j \to \infty} s_j < \infty$$

since otherwise Proposition 2.7 forwards in time and the strong δ_j -neck assumption backwards in time, c.f. Case 2 of the proof of Theorem 1.6, gives curvature bounds contradicting again (2.26).

After these preliminary reductions and observations, the proof is now verbatim as in [HK13a, Proof of Thm. 1.12], apart from some obvious changes in wording, like replacing α -Andrews flow by (α, δ) -flow, and from three minor modifications which we will carefully discuss now.

Modification 1: Instead of the nonsurgical version of the local curvature estimate [HK13a, Thm 1.8] and the convexity estimate [HK13a, Thm 1.10] we of course use the versions for (α, δ) -flows established in the present paper, Theorem 1.6 and Theorem 1.8, respectively.

Modification 2: The final paragraph of [HK13a, Proof of Thm. 1.12, Step 2] needs to be expanded, since the (α, δ) -flow $\hat{\mathcal{K}}^{\infty}$ might contain surgeries in $P(q_1, 0, r)$. If some neighborhood of q_1 is unmodified by surgeries at t = 0 (Definition 2.6), our previous argument applies. Otherwise, recall that $\hat{\mathcal{K}}^{\infty}$ arises as smooth limit of (α, δ) -flows $\hat{\mathcal{K}}^{j}$. After passing to a subsequence, we may assume that the surgery scales $s_{\sharp}(\hat{\mathcal{K}}^{j})$ converge to a limit s, which must be comparable to $H^{-1}(q_{1}, 0)$, by Theorem 1.6 and Definition 2.4. Let q'_{1} be a point on the radial segment in the cone X_{1} connecting q_{1} and the tip, such that $H(q'_{1}, 0) \gg$ $H(q_{1}, 0)$. Since all surgeries are done at comparable scales (Definition 1.3), Theorem 1.6 implies that for some r' > 0, the intersection $X_{1} \cap$ $B(q'_{1}, r')$ can be extended to a smooth (α, δ) -flow $\hat{\mathcal{K}}^{\infty}$ without surgeries in $P(q'_{1}, 0, r')$, and our previous argument applies.

Modification 3: In [HK13a, Proof of Thm. 1.12, Step 7] comparison with large enough spheres containing the origin is still legitimate, thanks to (2.27) and $0 \in X_{R,t}^{j}$, c.f. the proof of Claim 2.24. Thus, as in [HK13a, (3.4)] we obtain the estimate

(2.28)
$$\lim_{j \to \infty} \left(\sup_{\partial X^j_{R,t}} H \right) \le f(R,t)$$

for some continuous function f (at surgery times the estimate holds both for the pre- and post-surgery domain); this contradicts (2.26).

Finally, the curvature bounds for the derivatives of the second fundamental form follow from standard interior estimates. $\hfill \Box$

As mentioned in the introduction, the global curvature estimate (Theorem 1.10) enables us to smoothly pass to global limits. It can happen that the limit contains infinitely many surgeries, but we do get a bound for the number of surgeries contained in any compact set.

Definition 2.29 (Generalized (α, δ) -flow). A generalized (α, δ) -flow is a family of closed sets in \mathbb{R}^N that is an (α, δ) -flow when restricted to any open set $U \subset \mathbb{R}^N$ with compact closure.

Corollary 2.30 (Global convergence). There exists $\overline{\delta} = \overline{\delta}(\alpha) > 0$ with the following property. If \mathcal{K}^j is a sequence of (α, δ_j) -flows, $\delta_j \leq \overline{\delta}$, in $P(p_j, t_j, \eta_j H^{-1}(p_j, t_j))$ with $\eta_j \to \infty$, then, after passing to a subsequence, the (α, δ) -flows $\hat{\mathcal{K}}^j$ that are obtained from \mathcal{K}^j by parabolic rescaling $(p, t) \mapsto (H(p_j, t_j)(p - p_j), H^2(p_j, t_j)(t - t_j))$, restricting to $B(0, \Lambda_j)$ for a suitable sequence $\Lambda_j \to \infty$, and discarding the connected components that don't contain the origin, converge smoothly and globally to a limit $\mathcal{K}^{\infty} = \{K_t^{\infty} \subset \mathbb{R}^N\}_{t \in (-\infty, 0]}$, which is a generalized (α, δ) flow with convex time slices. In Corollary 2.30 the precise meaning of convergence is as follows.

Definition 2.31 (Smooth convergence). Let \mathcal{K}^j be a sequence of (α, δ_j) flows, $\delta_j \leq \overline{\delta}$, with connected time slices, normalized such that H(0,0) =1, and defined in $P(0,0,\Lambda_j)$ with $\Lambda_j \to \infty$, and let $\mathcal{K}^{\infty} = \{K_t^{\infty} \subset \mathbb{R}^N\}_{t \in (-\infty,0]}$ be a generalized (α, δ) -flow (Definition 2.29). We say that \mathcal{K}^j converges to \mathcal{K}^{∞} smoothly and globally, if \mathcal{K}^j converges smoothly to \mathcal{K}^{∞} away from the regions modified by surgeries, and if near every point $p \in K_{t_{\infty}}^{\infty,-}$ that that is modified by a surgery at time t_{∞} the following condition is satisfied. There exists a sequence of surgery times $t_j \to t_{\infty}$ in \mathcal{K}^j such that if we consider the forward and backward portions $\mathcal{K}^j_+ = \{K_{t+t_j}^j - p\}_{t \ge 0^+}$ and $\mathcal{K}^j_- = \{K_{t+t_j}^j - p\}_{t \le 0^-}$, and likewise $\mathcal{K}^{\infty}_{\pm}$, then \mathcal{K}^j_{\pm} converges smoothly to $\mathcal{K}^{\infty}_{\pm}$ in a forward respectively backward parabolic neighborhood $P_{\pm}(0, 0, \varepsilon)$, for some $\varepsilon > 0$.

Proof of Corollary 2.30. Let $\bar{\delta} = \bar{\delta}(\alpha) > 0$ small enough such that the previous estimates apply. Let \mathcal{K}^j be a sequence of (α, δ_j) -flows $(\delta_j \leq \bar{\delta})$ in $P(0, 0, \eta_j)$ $(\eta_j \to \infty)$ with H(0, 0) = 1. Choose $\Lambda_j \to \infty$ slowly enough such that the conclusion of Theorem 1.10 holds for the flow \mathcal{K}'^j , that is obtained from \mathcal{K}^j by restricting to $B(0, \Lambda_j)$ and discarding the connected components that don't contain the origin. We want to find a subsequence of \mathcal{K}'^j that converges smoothly and globally.

We can assume that there is some $R < \infty$ such that P(0, 0, R) contains surgeries of \mathcal{K}^{j} for large j, since otherwise [HK13a, Thm. 1.12, Rmk. 3.5] allows us to pass to a smooth limit. Also, it must be the case that the surgery scales $s_{j} = s_{\sharp}(\mathcal{K}^{j})$ (see Definition 1.3) satisfy (2.27) since otherwise Proposition 2.7 gives a contradiction with H(0, 0) = 1.

For each positive integer k, by Theorem 1.10, inequality (2.27), and Proposition 2.5, the parabolic ball P(0, 0, k) contains at most some controlled number N_k^j of surgeries of \mathcal{K}'^j , and their necks are of controlled size. After passing to a subsequence, we can assume that N_k^j equals some fixed number N_k , that the surgery times converge to some limiting surgery times given by a set $\mathcal{T}_k \subset (-\infty, 0]$ with at most N_k elements, and that the (pre and post) surgery time slices converge smoothly. Let $\mathcal{Q} \subset (-\infty, 0]$ be a countable dense set that is disjoint from $\cup_k \mathcal{T}_k$. Arguing as in [HK13a, Proof of Thm. 1.12, Step 7], after passing to a subsequence there are convex sets K_t^{∞} such that the domains $X_{R,t}^j$ converge smoothly to $K_t^{\infty} \cap B(0, R)$ as $j \to \infty$, for all $R < \infty$ and all $t \in \mathcal{Q}$. Finally, putting everything together, namely the convergence at a dense set of times, the convergence at the surgery times, Theorem 1.10, Proposition 2.5 and the Andrews-condition, it follows that there exists a generalized (α, δ) -flow \mathcal{K}^{∞} such that $\mathcal{K}^{\prime j} \to \mathcal{K}^{\infty}$ smoothly and globally, where the meaning of convergence is as in Definition 2.31. \Box

3. Ancient solutions and standard solutions

We will now prove some structural results for uniformly 2-convex ancient α -Andrews flows, and results for the evolution of standard caps. These results will be used later in the blowup analysis in Section 4.

3.1. Structure of uniformly 2-convex ancient α -Andrews flows. In this section, we consider smooth ancient α -Andrews flows $\{K_t \subset \mathbb{R}^N\}_{t \in (-\infty,T)}$, that are β -uniformly 2-convex, i.e. $\lambda_1 + \lambda_2 \geq \beta H$ for some fixed $\beta > 0$. Examples to keep in mind are the cylinder, the bowl soliton [AW94], the sphere and the Angenent ovals [Whi03, HH13].

We recall that ancient α -Andrews flows are always convex [HK13a, Cor. 2.15], and in fact automatically smooth until they become extinct [HK13a, Thm. 1.14]. We will now discuss two structural results that are more specific to the uniformly 2-convex case. First, non ε -neck points are at controlled distance from one another, unless the time slice is compact and the points approximately realize the diameter.

Proposition 3.1 (Non-neck points). For all $\varepsilon_1, \varepsilon_2 > 0$ there exists $\underline{R} = \underline{R}(\varepsilon_1, \varepsilon_2, \alpha, \beta) < \infty$, such that if \mathcal{K} is a β -uniformly 2-convex ancient α -Andrews flow and $p_1, p_2 \in \partial K_t$ are not strong ε_1 -neck points, then at least one of the following holds:

(1) $\max_i H(p_i)d(p_1, p_2) < \underline{R}.$ (2) diam $K_t \le (1 + \varepsilon_2)d(p_1, p_2).$

Second, as suggested by the bowl soliton, non-compact non-cylindrical solutions have a single cylindrical end in a precise quantitative sense.

Proposition 3.2 (Quantitative one-endedness). For all $\varepsilon_1, \varepsilon_2 > 0$ there exists $\underline{R} = \underline{R}(\varepsilon_1, \varepsilon_2, \alpha, \beta) < \infty$, such that if \mathcal{K} is a β -uniformly 2-convex non-compact ancient α -Andrews flow and $p \in \partial K_t$ is not a strong ε_1 -neck point, then for any $R > \underline{R}$, there exists a strong ε_2 -neck point $q \in \partial K_t \cap S(p, RH^{-1}(p, t))$ such that the intersection $K_t \cap S(p, RH^{-1}(p, t))$ is contained in the $\varepsilon_2 H^{-1}(q, t)$ neighborhood of a cross-sectional disc of the solid ε_2 -neck at q.

For the proofs we need the following two lemmas.

Lemma 3.3. If \mathcal{K} is a uniformly 2-convex ancient α -Andrews flow that becomes extinct at a finite time $T < \infty$, then the final time slice K_T is a convex set of dimension at most 1.⁴

Proof of Lemma 3.3. This follows immediately by combining [HK13a, Thm. 1.14] and [HK13a, Refinement of Thm. 1.15].⁵ \Box

Lemma 3.4. Suppose $\{\mathcal{K}^j\}$ is a sequence of β -uniformly 2-convex ancient α -Andrews flows, for which the time-slices K_0^j Hausdorff-converge to a 1-dimensional convex set K^{∞} containing p as an interior point. Then for any $\delta > 0$, any sequence of points $p_j \in \partial K_0^j$ with $p_j \to p$ consists of strong δ -neck points for large j.

Remark 3.5. We point out that K^{∞} can be strictly smaller than the final time slice K_0^{∞} of the space-time Hausdorff limit $\mathcal{K}^j \to \mathcal{K}^{\infty}$. For example, if \mathcal{K}^j is a blowdown sequence of the bowl soliton centered at the tip (0,0), then K^{∞} is a half-line, but K_0^{∞} is the whole line (since the limit \mathcal{K}^{∞} is a shrinking cylinder that becomes extinct at time 0).

Proof of Lemma 3.4. We have $H(p_j) \to \infty$ as $j \to \infty$, for otherwise K^{∞} would be a domain with nonempty interior. Choose $x^{\pm} \in K^{\infty}$ such that p lies in the interior of the segment $\overline{x^-x^+}$, and choose $x_j^{\pm} \in K_{t_j}^j$ such that $x_j^{\pm} \to x^{\pm}$. The segments $\overline{x_j^- p_j}$ and $\overline{p_j x_j^+}$ are contained in $K_{t_j}^j$ by convexity, and the angle between them converges to π . Parabolically rescaling to normalize $H(p_j)$ and passing to a limit [HK13a, Thm. 1.12], we obtain an ancient uniformly 2-convex α -Andrews flow \mathcal{K} that contains a line. Using [HK13a, Lemma 3.14] it follows that \mathcal{K} is a round cylindrical flow. Thus p_j is a strong δ -neck point for large j.

Proof of Proposition 3.1. Assume towards a contradiction that there are a sequence $\{\mathcal{K}^j\}$ of β -uniformly 2-convex ancient α -Andrews flows, and sequences of points $p_1^j, p_2^j \in K_0^j$ which are not strong ε_1 -neck points, such that $\max_i H(p_i^j)d(p_1^j, p_2^j) \to \infty$, and $\dim K_0 > (1 + \varepsilon_2)d(p_1^j, p_2^j)$ for all j. Rescaling so that $d(p_1^j, p_2^j) = 1$ we get $\max_i H(p_i^j) \to \infty$. After passing to a subsequence, we may assume that $p_i^j \to p_i^\infty$ for i = 1, 2, and that $\hat{\mathcal{K}}^j$ converges, by [HK13a, Thm 1.14], to an ancient α -Andrews flow $\hat{\mathcal{K}}^\infty$ which goes extinct at T = 0, where diam $\hat{\mathcal{K}}^\infty \ge 1 + \varepsilon_2$. By Lemma 3.3, at least one of the points p_1^∞, p_2^∞ must be an interior point. Thus, Lemma 3.4 gives a contradiction to the assumption that $\{p_1^j, p_2^j\}$ are not strong ε_1 -neck points.

20

⁴Conjecturally, K_T is either a point or the entire real line, c.f. [Whi03].

⁵It is of course possible to phrase this argument entirely in the smooth setting.

Proof of Proposition 3.2. Assume towards a contradiction that there are a sequence $\{\mathcal{K}^j\}$ of β -uniformly 2-convex non-compact ancient α -Andrews flows, and a sequence $R_j \to \infty$, such that $(0,0) \in \partial K_0^j$ is not a strong ε_1 -neck point, H(0,0) = 1, and $\partial K_0^j \cap S(0,R_j)$ does not contain any point with the asserted property.

By non-compactness and convexity there is a ray in K_0^j starting at the origin. Let x_j be the point where the ray intersects $S(0, R_j)$, and let q_j be a point in $K_0^j \cap S(0, R_j)$ with maximal distance from x_j ; thus we have $q_j \in \partial K_0^j$. We claim that $\frac{d(x_j, q_j)}{R_j} \to 0$; indeed, this follows by rescaling by R_j^{-1} and passing to a limit [HK13a, Thm 1.14], and using the fact that the limit K^{∞} of the time zero slices is non-compact, convex, and 1-dimensional (since it is contained in K_0^{∞} , the time zero slice of the space-time limit \mathcal{K}^{∞} , which is at most 1-dimensional by Lemma 3.3). Using Lemma 3.4, we see that 0 must be an endpoint, and thus that K^{∞} is a ray starting at 0.

After shifting x_j to the origin, parabolically rescaling by $d(x_j, q_j)^{-1}$, and passing to a subsequence, we obtain a new sequence $\hat{\mathcal{K}}^j$ which strongly Hausdorff converges to an ancient α -Andrews flow $\hat{\mathcal{K}}^\infty$. Note that \hat{K}_0^∞ must actually be smooth, since a convex set of dimension 1 cannot contain 2 perpendicular segments. Since $\hat{\mathcal{K}}^\infty$ contains a line and is uniformly 2-convex, it must be a round cylindrical flow, c.f. the proof of Proposition 3.1. Moreover, $(S(0, R_j), x_j)$ converges after rescaling by $d(x_j, q_j)^{-1}$ to a hyperplane orthogonal to the axis of the cylinder, and we get a contradiction.

Remark 3.6. The proof of Proposition 3.2 also works if the assumption that the flow is noncompact is replaced by the assumption that there exists a point in K_t with distance from p at least $(1 + \varepsilon)R$.

For later use (namely for the proof of Claim 4.6), we also prove that the ratio between intrinsic and extrinsic distance is controlled.

Proposition 3.7 (Intrinsic distance). For every $R < \infty$, there is an $L = L(R, \alpha) < \infty$ such that for every ancient α -Andrews flow \mathcal{K} and every point $(p,t) \in \partial K_t$, any point $x \in \partial K_t \cap B(p, RH^{-1}(p,t))$ can be joined to p by a path in ∂K_t of length at most $LH^{-1}(p,t)$.

Proof. If not, there is a sequence \mathcal{K}^j of ancient α -Andrews flows with H(0,0) = 1, and a sequence of points $x_j \in \partial K_0^j \cap B(0,R)$ that cannot be joined to 0 by a path in ∂K_0^j of length at most j. By [HK13a, Thm. 1.12] we can pass to a subsequential limit \mathcal{K}^∞ with convex (in particular

connected) time slices. Since K_0^{∞} cannot be a slab, the boundary ∂K_0^{∞} is also connected, which gives a contradiction for j large enough. \Box

3.2. The standard surgery solution. We will first consider standard caps K^{st} as in Definition 2.2, that are α -Andrews and β -uniformly 2-convex for some values $\alpha = \alpha(K^{\text{st}}), \beta = \beta(K^{\text{st}}) > 0$. Afterwards, we will construct a particular model of K^{st} that is suitable for surgeries.

Proposition 3.8. Let K^{st} be a standard cap as in Definition 2.2, with $\alpha = \alpha(K^{st}), \beta = \beta(K^{st}) > 0$. There is a unique mean curvature flow $\{K_t\}_{t \in [0,1/2(N-2))}$ starting at K^{st} . It has the following properties:

- (1) It is α -Andrews, convex, and β -uniformly 2-convex.
- (2) There are continuous increasing functions $\underline{H}, \overline{H} : [0, \frac{1}{2(N-2)}) \to \mathbb{R}_+$ with $\underline{H}(t) \to \infty$ as $t \to \frac{1}{2(N-2)}$ such that $\underline{H}(t) \leq H(p,t) \leq \overline{H}(t)$ for all $p \in \partial K_t$ and $t \in [0, \frac{1}{2(N-2)})$.
- (3) For every $\varepsilon > 0$ and $\tau < \frac{1}{2(N-2)}$ there exists an $R = R(\varepsilon, \tau) < \infty$ such that outside B(0, R) the flow $\{K_t\}_{t \in [0, \tau]}$ is ε -close to the evolution of the solid round unit cylinder $\bar{D}^{N-1} \times \mathbb{R}$.
- (4) For every $\varepsilon > 0$, there exists a $\tau = \tau(\varepsilon) < \frac{1}{2(N-2)}$ such that every point $(p,t) \in \partial K_t$ with $t \ge \tau$ is ε -close to a β -uniformly 2-convex ancient α -Andrews flow.

Proof. The argument is closely related to the one for Ricci flow [Per03, KL08, MT07], so we will give a brief, but complete, treatment.

Existence and uniqueness of a smooth solution on a maximal time interval [0, T) with bounded curvature on compact subintervals follows from standard theory, see e.g. [EH91, Thm. 4.2], [CY07, Thm. 1.1]. Moreover, since $A = A_j^i$ evolves by $\partial_t A = \Delta A + |A|^2 A$, it follows from the tensor-maximum principle, see e.g. [CCG⁺08, Thm. 12.34], that convexity and β -uniform 2-convexity are preserved along the flow.

Assume (3) fails for some $\varepsilon > 0$, $\tau < T$. Since we have uniform curvature bounds on $[0, \tau]$, we can pass to smooth limits. Thus, as a pointed limit around a suitable sequence of points going to infinity we get a mean curvature flow with bounded curvature that starts from a round cylinder, but is not ε -close to the standard evolution of the round cylinder; this contradicts uniqueness of the evolution of the cylinder.

Next, we recall from [ALM13, Thm. 2] that the quantities \underline{Z} , \overline{Z} introduced there, satisfy the evolution inequalities

(3.9) $\partial_t \underline{Z} \ge \Delta \underline{Z} + |A|^2 \underline{Z}, \qquad \partial_t \overline{Z} \le \Delta \overline{Z} + |A|^2 \overline{Z}$

in the viscosity sense. Arguing as in [ALM13, Cor. 3], it follows that the α -Andrews condition is preserved;⁶ this completes the proof of (1).

Assume T is strictly less than $\frac{1}{2(N-2)}$. Then, since we have shown that (3) holds for $\tau < T$, by the local curvature estimate [HK13a, Thm. 1.8, Rem. 2.10], we obtain uniform bounds on the curvature near infinity; hence the curvature has to blow up inside some compact set. Select a sequence of points (p_j, t_j) with $Q_j = H(p_j, t_j) \to \infty$ and $H \leq 2Q_j$ on $P(p_j, t_j, jQ_j^{-1})$. By convexity and since the curvature stays bounded outside some compact set, the time slices K_{t_j} contain some cones based at p_j with a definite lower bound on the cone angle. Rescaling by Q_j^{-1} and passing to a limit, we obtain an ancient α -Andrews flow with nonvanishing asymptotic volume ratio; this contradicts [HK13a, Rem. 1.20]. Thus $T = \frac{1}{2(N-2)}$.

Since the solution is contained in a shrinking cylinder that becomes extinct at $T = \frac{1}{2(N-2)}$, the Andrews condition implies that the curvature must blow up everywhere as $t \to \frac{1}{2(N-2)}$, i.e. we get (2).

Assertion (4) follows from (2) and the global convergence theorem [HK13a, Thm. 1.12].

Suppose $\{K'_t\}_{t\in[0,T')}$ is any smooth mean curvature flow starting at K^{st} , without any a priori bound on curvature. Let $t_1 \leq \min\{T', \frac{1}{2(N-2)}\}$ be the supremum of the numbers t such that \mathcal{K}' coincides with \mathcal{K} constructed above. If $t_1 < \min\{T', \frac{1}{2(N-2)}\}$, then $\hat{K}_{t_1} = K_{t_1}$, and by pseudolocality [CY07, Thm. 7.5] the curvature of \mathcal{K}' is bounded uniformly for a time interval $[t_1, t_2)$ for some $t_2 > t_1$, contradicting the uniqueness theorem. Thus we obtain a unique solution without the bounded curvature assumption.

Proposition 3.10 (Existence of a suitable standard cap). Given any $\alpha = (\alpha, \beta, \gamma) \in (0, N-2) \times (0, \frac{1}{N-2}) \times (0, \infty)$, there exist $K^{st} \subset \mathbb{R}^N$, $\alpha^{st} > \alpha$, $\beta^{st} > \beta$, and $\overline{\delta} > 0$ (all depending on α) with the following properties:

- (1) K^{st} is a standard cap (see Definition 2.2)
- (2) K^{st} is α^{st} -Andrews and β^{st} -uniformly 2-convex.
- (3) If $\delta \leq \delta$, then it is possible to do surgery such that all properties in Definition 2.4 hold, and such that in addition the α -Andrews condition and the β -uniform 2-convexity are preserved.

⁶Since we already know that the solution is asymptotically cylindrical, there is no subtle part at all about localizing the maximum principle.

Proof. We can take K^{st} a smooth convex domain that is a small perturbation of a solid half cylinder with a half ball attached. This obviously satisfies (1) and (2). Let $\{N_t\}$ be a strong δ -neck with optimal quality δ small enough. Since the neck is strong by assumption, by standard interior estimates we get uniform bounds for all derivatives of the curvatures. It is then clear, that we can cut along the final time slice of the neck and glue in two copies of K^{st} such that item (1), (2) and (4) of Definition 2.4 hold, and such that the α -Andrews condition and the β -uniform 2-convexity are preserved also. The only somewhat nonobvious point is item (3) of Definition 2.4, but as observed by Hamilton [Ham97] (see also [KL08, Sec. 72] or [HS09, p. 155]) this can be ensured by bending the cylinder slightly inwards.

4. EXISTENCE OF MEAN CURVATURE FLOW WITH SURGERY

We keep the parameters $\alpha = (\alpha, \beta, \gamma)$ fixed for this section.

4.1. The canonical neighborhood theorem. In the proof of our main existence result, Theorem 1.21, we will consider sequences of flows where the \mathbb{H} -parameters degenerate suitably. We will now prove the following crucial self-improvement phenomenon: If the surgeries are done on necks, where we a priori only know that they have at least some small but fixed quality $\overline{\delta}$, then the degeneration of the other parameters actually forces them to be more and more precise. In fact, this is just a reformulation of the canonical neighborhood theorem.

Theorem 4.1 (Self-improvement of necks). There exists a constant $\overline{\delta} = \overline{\delta}(\alpha) > 0$ with the following property. If \mathcal{K}^j is a sequence of $(\alpha, \delta_j, \mathbb{H}_j)$ -flows $(\delta_j \leq \overline{\delta})$ with $H^j_{trig}/H^j_{neck}, H^j_{neck}/H^j_{th} \to \infty$, and if $(p_j, t_j) \in \partial \mathcal{K}^j$ is a sequence of points with $H(p_j, t_j) \to \infty$, then after parabolic rescaling to normalize H at (p_j, t_j) and passing to a subsequence we have smooth convergence to either (a) a β -uniformly 2-convex ancient α -Andrews flow, or (b) the evolution of a standard surgery cap preceded by the evolution of a round cylindrical flow.⁷

Remark 4.2. Note that Theorem 4.1 is equivalent to Theorem 1.22.

Outline of the proof. The proof amounts to classifying the limits, whose existence is guaranteed by the global convergence result (Corollary 2.30). If the limit doesn't contain surgeries, then it must be a β -uniformly 2-convex ancient α -Andrews flow. If the limit contains a

24

⁷In case (b), the meaning of smooth convergence is as in Definition 2.31; also, the notion of ε -closeness in Theorem 1.22 should be interpreted accordingly.

surgery, then we argue, using in particular part (2) of Definition 1.17, the assumption that the curvature ratios degenerate, and the global curvature estimate (Theorem 1.10), that the limit must contain a line. It is then easy to conclude that there is in fact only one surgery, and that the limit must have the structure as claimed. Finally, we observe that potential other connected components get cleared out.

Proof. Let $\bar{\delta} = \bar{\delta}(\alpha) > 0$ small enough such that the estimates from the previous sections apply. Since the initial domain has principal curvatures bounded by γ/α , the curvature remains bounded for a definite amount of time. Thus, the rescaled flows $\hat{\mathcal{K}}^j$ are defined on parabolic balls $P(0,0,\eta_j)$ with $\eta_j \to \infty$ as $j \to \infty$. By Corollary 2.30, after passing to a subsequence and discarding connected components that don't contain the origin, $\hat{\mathcal{K}}^j \to \hat{\mathcal{K}}$ smoothly and globally, where the limit $\hat{\mathcal{K}}$ is a generalized $(\alpha, \bar{\delta})$ -flow with convex – and thus in particular connected – time slices.

If $\hat{\mathcal{K}}$ doesn't contain surgeries, then it is a β -uniformly 2-convex ancient α -Andrews flow. Otherwise, let $T \in (-\infty, 0]$ be a surgery time and let $\hat{N} \subset \hat{K}_T^-$ be a surgery neck of quality $\bar{\delta}$ sitting in the backward time slice.

Claim 4.3. $\hat{K}_T^- \setminus \hat{N}$ has two unbounded components.

Proof of Claim 4.3. Note that \hat{N} is the limit of some solid $\bar{\delta}$ -necks \hat{N}^j in the approximators $\hat{\mathcal{K}}^j$. By part (2) of Definition 1.17, we can find a curve γ_j in the approximator connecting $\{H = \hat{H}_{\text{trig}}\}$ and $\{H \leq \hat{H}_{\text{th}}\}$, such that it passes through \hat{N}^j but avoids all other $\bar{\delta}$ -necks of the disjoint collection. We can assume that the curve γ_j enters and leaves \hat{N}^j exactly once (if this isn't already the case, we can look at the earliest entry point and the latest exit point and change γ_j for intermediate times to a curve within \hat{N}^j). Note furthermore that γ_j must intersect each of the two boundary discs of the cylinder exactly once (since otherwise we could modify it within \hat{N}^j into a curve avoiding all necks completely, contradicting the minimal separation property). Let x_j be the center of \hat{N}^j . By the global curvature estimate (Theorem 1.10), for every $\Lambda < \infty$ we have

(4.4)
$$\hat{H}(x)/\hat{H}_{neck} \le C(\Lambda) < \infty$$
 whenever $d(x, x_j) \le \Lambda \hat{H}_{neck}^{-1}$

By the Andrews condition and the local curvature estimate (Theorem 1.6), we also have a lower bound

(4.5)
$$\hat{H}(x)/\hat{H}_{\text{neck}} \ge c(\Lambda) > 0$$
 whenever $d(x, x_j) \le \Lambda \hat{H}_{\text{neck}}^{-1}$.

Since $H_{\text{trig}}^j/H_{\text{neck}}^j, H_{\text{neck}}^j/H_{\text{th}}^j \to \infty$, given any $\Lambda < \infty$, for j large enough the curve γ_j must start and end outside $B(x_j, \Lambda \hat{H}_{\text{neck}}^{-1})$. Thus, $\hat{K}_T^- \setminus \hat{N}$ has at least two unbounded components. Since \hat{K}_T^- is connected, $\hat{K}_T^- \setminus \hat{N}$ must have exactly two components. \Box

Since \hat{K}_T^- has two ends (see Claim 4.3), it contains a line, and by monotonicity of sets all prior time slices contain this line also. At each fixed time the convex set splits off an \mathbb{R} -factor, and thus there cannot be any other surgeries. It follows that $\hat{\mathcal{K}}$ is a round cylindrical flow for t < T (c.f. the proof of Lemma 3.4). Similarly, by the uniqueness of the standard solution (see Proposition 3.8), $\hat{\mathcal{K}}$ must be the evolution of the standard cap for t > T.

Finally, arguing as in [HK13a, Proof of Cor. 2.15] it follows that potential other connected components are cleared out, i.e. $\hat{\mathcal{K}}^j \to \hat{\mathcal{K}}$ smoothly and globally without the need of discarding connected components that don't contain the origin.

4.2. Existence of $(\alpha, \delta, \mathbb{H})$ -flows. We can now prove our main existence theorem for mean curvature flow with surgery, Theorem 1.21.

Outline of the proof. We assume towards a contradiction that we have a sequence \mathcal{K}^j of flows with degenerating \mathbb{H} -parameters that can be defined only on some finite maximal time intervals $[0, T_j]$. For j large, to obtain a contradiction, we want to argue that we can perform surgery and thus continue the flow beyond T_j . This amounts to finding suitable collection of δ -necks. To this end, we first prove Claim 4.6 which shows that the thick and the trigger part in $K_{T_j}^j$ can be separated by a union of balls centered at boundary points with $H(p) = H_{\text{neck}}$ and radius comparable to H_{neck}^{-1} . We then consider a minimal collection of balls with the separation property and prove that their centers are actually centers of strong $\hat{\delta}$ -necks for any $\hat{\delta}$, see Claim 4.7. It is then easy to conclude the proof.

Proof of Theorem 1.21. Let $\bar{\delta} = \bar{\delta}(\alpha) > 0$ small enough such that all previous estimates and also the argument in the last line of the present proof apply. By the maximum principle, the α -Andrews condition and β -uniform 2-convexity are preserved along smooth mean curvature flow [And12, Hui84]. We assume towards a contradiction that for some $\delta \leq \bar{\delta}$ there is no constant $\Theta(\delta) < \infty$ such that the assertion of the theorem holds. Then there is a sequence \mathcal{K}^j of $(\alpha, \delta, \mathbb{H}_j)$ -flows with

 $H_{\text{trig}}^{j}/H_{\text{neck}}^{j}, H_{\text{neck}}^{j}/H_{\text{th}}^{j}, H_{\text{neck}}^{j} \to \infty$, that can only be defined on a finite maximal time interval I_{j} . If some I_{j} were a half-open interval $[0, T_{j})$, then the fact that $H \leq H_{\text{trig}}^{j}$ would allow us to pass to a limit as $t \to T_{j}$, so that $[0, T_{j})$ is not maximal; therefore $I_{j} = [0, T_{j}]$ for some $T_{j} < \infty$. Moreover, it must be the case that we cannot find a minimal collection of strong δ -necks in $K_{T_{j}}^{j}$ as required by the definition of an $(\alpha, \delta, \mathbb{H}_{j})$ -flow (Definition 1.17), since otherwise we could cut along them (Proposition 3.10) and run smooth MCF for a short time, contradicting the maximality of T_{j} .⁸ Therefore our goal is to produce such a collection for large j, to obtain a contradiction.

Let \mathcal{I}_j be the set of points $p \in \partial K^j_{T_j}$ with $H(p) > H^j_{\text{neck}}$, and let \mathcal{J}_j be the set of points $p \in \partial K^j_{T_j}$ with $H(p) = H^j_{\text{neck}}$.

Claim 4.6 (Separation property). There is a constant $C = C(\alpha) \in (2N, \infty)$, such that the union $V_j = \bigcup_{p \in \mathcal{J}_j} B(p, CH^{-1}(p))$, for j large enough, separates $\{H = H^j_{trig}\}$ from $\{H \leq H^j_{th}\}$ in the domain $K^j_{T_i}$.

Proof of Claim 4.6. We suppress j in the notation. Fix $C \in (2N, \infty)$ to be determined later, and let $U = \bigcup_{p \in \mathcal{I}} B(p, 2NH^{-1}(p))$. To establish the claim, it suffices to prove that $U \setminus V$ is open and closed in $K_T \setminus V$. Indeed, for j large enough we have (c.f. (4.4) and (4.5)) that $\{H = H_{\text{trig}}\} \subseteq U \setminus V$ and $\{H \leq H_{\text{th}}\} \subseteq K_T \setminus (U \cup V)$. Thus, once we know that $U \setminus V \subseteq K_T \setminus V$ is open and closed, it follows that the sets $\{H = H_{\text{trig}}\}$ and $\{H \leq H_{\text{th}}\}$ lie in two different components of $K_T \setminus V$.

Starting with the obvious part, the set U is open since it is a union of open sets. Thus, $U \setminus V = U \cap (K_T \setminus V)$ is open in $K_T \setminus V$.

Suppose now $x \in \overline{U \setminus V}$ lies in the closure of $U \setminus V \subseteq K_T \setminus V$. We want to show that $x \in U \setminus V$. There are sequences $\{x_k\} \subset U$, $\{p_k\} \subset \{H > H_{\text{neck}}\}$ such that $x_k \to x$ and $x_k \in B(p_k, 2NH^{-1}(p_k))$. Passing to subsequences, we may assume that $p_k \to p$. Then $H(p) \geq H_{\text{neck}}$. If $H(p) = H_{\text{neck}}$, then $x \in B(p, CH^{-1}(p)) \subseteq V$ since C > 2N; a contradiction to the assumption that $x \in \overline{U \setminus V} \subseteq K_T \setminus V$. Hence we have $H(p) > H_{\text{neck}}$. Let $y \in \partial K_T$ be a point nearest x. Then $d(y, x) \leq (N-1)H^{-1}(y)$, in particular $x \in B(y, 2NH^{-1}(y))$. If $H(y) > H_{\text{neck}}$, we obtain that $x \in U$ and thus that $x \in U \setminus V$, what we wanted to show. If $H(y) = H_{\text{neck}}$, we obtain $x \in V$; a contradiction.

⁸In particular, if $\{H(\cdot, T_j) \leq H_{\text{th}}^j\} = \emptyset$ everything is discarded at $t = T_j$, and thus the flow can be continued forever as empty flow, contradicting maximality.

Finally, let us rule out the remaining case $H(y) < H_{\text{neck}}$. Since y is a boundary point nearest to x, we have $d(y,p) \leq 2d(x,p) \leq 4NH^{-1}(p)$. Thus, by Proposition 3.7, Proposition 3.8 and Theorem 4.1 there is a $C_1 = C_1(\alpha) < \infty$ such that, for j large enough, y and p lie in the same connected component of $B(p, C_1H^{-1}(p)) \cap \partial K_T$. Since $H(y) < H_{\text{neck}} < H(p)$, there is a $z \in B(p, C_1H^{-1}(p)) \cap \partial K_T$ with $H(z) = H_{\text{neck}}$. Note that $d(z,x) \leq d(z,p) + d(p,x) \leq (C_1 + 2N)H^{-1}(p)$, and thus $H(z)d(z,x) \leq (C_1 + 2N)\frac{H(z)}{H(p)} \leq C$, for some $C = C(\alpha) < \infty$ by Theorem 1.10. This shows $x \in B(z, CH^{-1}(z)) \subseteq V$; a contradiction.

Let $\hat{\mathcal{J}}_j \subseteq \mathcal{J}_j$ be a minimal (and thus finite) subset such that the union of balls $\cup_{p \in \hat{\mathcal{J}}_j} B(p, CH^{-1}(p))$ has the separation property.

Claim 4.7 (Strong neck point property). Given any $\hat{\delta} > 0$, for j large enough all points in $\hat{\mathcal{J}}_i$ are strong $\hat{\delta}$ -neck points.

Proof of Claim 4.7. Suppose that for all j one of the points $p_j \in \mathcal{J}_j$ is not a strong $\hat{\delta}$ -neck. By Theorem 4.1, parabolically rescaling to make $H(p_i, T_i) = 1$ and passing to a subsequence $\hat{\mathcal{K}}^j$, we get a limit ancient (α, δ) -flow $\hat{\mathcal{K}}$ with basepoint $(p, t) = (0, 0) \in \partial \hat{\mathcal{K}}$ that is either (a) a β -uniformly 2-convex ancient α -solution, or (b) an evolution of the standard surgery cap preceded by a round shrinking cylinder. In case (a), since the balls separate and the curvature ratios tend to infinity, the limit $\hat{\mathcal{K}}$ is clearly non-compact. Consider a $\hat{\delta}$ -neck \hat{N} at a point q at t = 0 provided by Proposition 3.2. Here we chose q far enough away to ensure that \hat{N} and B(p, 2C) are disjoint. This neck \hat{N} is the limit of some $\hat{\delta}$ -necks \hat{N}^{j} in the approximators. As in the proof of Claim 4.3, let γ_i be a curve in the approximators that connects the trigger and the thick part and passes through $B(p_i, C)$, but avoids all other balls from the minimal collection of separating balls. Always assuming j is large enough, we can short circuit γ_j inside \hat{N}^j . After this modification, γ_i misses $B(p_i, C)$. It also still misses all other balls from our minimal collection, unless \hat{N}^{j} itself intersects a ball $B' = B(p'_{i}, C)$ for some $p'_i \in \mathcal{J}_i \setminus \{p_i\}$. This however is impossible, because in this case one of the complementary components would have uniformly bounded diameter. Thus, we get a contradiction with the separation property of \mathcal{J}_i . Finally, in case (b), by Proposition 3.8 we get a contradiction similar to the one in case (a). By Claim 4.7, for large j, every $p \in \hat{\mathcal{J}}_j$ is a strong δ -neck point. These necks are disjoint for large j, since otherwise again by Claim 4.7 two intersecting δ -necks would lie in a single $\hat{\delta}$ -neck for $\hat{\delta} \ll \delta$, which is impossible by minimality of \mathcal{J}_j . Thus, we have a minimal collection of disjoint strong δ -necks with the separation property (since $C(\alpha) > 2N$ and $\bar{\delta}$ is small enough); this gives the desired contradiction. \Box

4.3. Further properties. Finally, for convenience of the reader we explain how Corollaries 1.25, 1.26 and Proposition 1.27 are obtained as easy consequences (in fact, the proof of Proposition 1.27 could have been given right after stating the axioms of an $(\alpha, \delta, \mathbb{H})$ -flow).

Proof of Corollary 1.25. Fix $\varepsilon_1 = \varepsilon_1(N) \ll 1$ such that the gluing argument below works (Claim 4.8). Fix $\bar{\varepsilon} \ll \min\{\varepsilon_1, \underline{R}_1^{-1}, \underline{R}_2^{-1}\}$, where $\underline{R}_i = \underline{R}(\varepsilon_1/2, \varepsilon_1/2, \alpha, \beta)$ are the constants from Proposition 3.1 and Proposition 3.2, respectively, and let $\underline{R}_3 = \underline{R}_3(\bar{\varepsilon}, \tau(\bar{\varepsilon}))$ be the constant from Proposition 3.8. We will prove that the corollary holds for any $\varepsilon \ll \min\{\bar{\varepsilon}, \underline{R}_3^{-1}\}$.

Let C be a discarded component, and consider the (possibly empty) set $\mathcal{I} \subseteq \partial C$ of ε_1 -neck points.

Claim 4.8. There is a set $N, \mathcal{I} \subseteq N \subseteq C$, such that for each $q \in N$ there is a point $p \in \mathcal{I}$ with $d(p,q) \leq 100H^{-1}(p)$, and such that either $N = C \cong \overline{D}^{N-1} \times S^1$ or $N \subsetneq C$ and each component of N has the topology of $\overline{D}^{N-1} \times I$, for some open interval I.

Proof of Claim 4.8. Let $\mathcal{I}' \subseteq \partial C$ be a maximal collection of ε_1 -neck points such that for any pair $p, q \in \mathcal{I}'$ the separation between them is at least $50\min\{H^{-1}(p), H^{-1}(q)\}$. Note that each $p \in \mathcal{I}'$, being an ε_1 -neck point for ε_1 small, comes with a cylindrical neighborhood of length much longer than $50H^{-1}(p)$. Following this cylinder in each direction, we see that there are 0, 1 or 2 neighboring points $p' \in \mathcal{I}'$, where by neighboring point we mean a point of \mathcal{I}' closest to p in a direction of the cylinder and at distance at most $110H^{-1}(p)$. Let C_p be the open core of a neck at p going say 75% of the distance towards each neighboring point (and going say $10H^{-1}(p)$ in directions without neighboring point), and let $N = \bigcup_{p \in \mathcal{I}'} C_p$. By construction, the set N is built by gluing together necks with intersection multiplicities at most 2 and substantial overlap. Since ε_1 is small enough, it follows that the connected components of N must be either $\bar{D}^{N-1} \times S^1$ or $\bar{D}^{N-1} \times I$. Since C is connected, this implies the claim. If N = C, then C has the topology of $\overline{D}^{N-1} \times S^1$, and we are done. If $N = \emptyset$ then, always assuming ε is sufficiently small, it follows from Theorem 1.22, Proposition 3.1 and Proposition 3.8, that C is modeled on a compact ancient α -solution of controlled geometry, and thus that is has the topology of \overline{D}^N . Assume now $\emptyset \subseteq N \subseteq C$, and let $N' \subseteq N$ be a component of maximal diameter.

We first consider the case diam $N' \leq 10\underline{R}H^{-1}(p)$ for some ε_1 -neck point $p \in \mathcal{I} \cap N'$, where $\underline{R} = \max\{\underline{R}_1, \underline{R}_2\}$. If the ε -model at p provided by Theorem 1.22 were the cylinder, then the connected component of \mathcal{I} containing p would have diameter larger than $\frac{1}{\varepsilon}H^{-1}(p)$, contradicting the assumption that diam $N' \ll \frac{1}{\varepsilon}H^{-1}(p)$ and $\mathcal{I} \subseteq N$. If the ε -model at p were a noncompact α -solution or a compact α -solution with diameter much larger than $\underline{R}H^{-1}(p)$, then we could apply Proposition 3.2 (see also Remark 3.6) at a point $p' \in \partial N' \cap \partial C$, which by definition is not in \mathcal{I} and has curvature comparable to p by Theorem 1.10, and would again get, c.f. Claim 4.8, a much longer ε_1 -neck at controlled distance, contradicting the assumption that N' is maximal. If the ε model at p were a standard solution, Proposition 3.8 would give a similar contradiction. Thus, the ε -model at p must be a compact α solution of controlled size, and thus C has the topology of \overline{D}^N .

Let us now consider the remaining case that diam $N' > 10\underline{R}H^{-1}(p)$ for all ε_1 -neck points $p \in \mathcal{I} \cap N'$. Since for each $q \in N$ there is a point $p \in \mathcal{I}$ with $d(p,q) \leq 100H^{-1}(p)$ and since ε_1 is small, it follows that

(4.9)
$$\operatorname{diam} N' > 9\underline{R}H^{-1}(p)$$

actually holds for all points $p \in N' \cap \partial C$. Select points $p_{\pm} \in \partial N' \cap \partial C$ on the boundary circles of the neck N'. By definition we have $p_{\pm} \notin \mathcal{I}$. We now apply Theorem 1.22, Proposition 3.2 and Proposition 3.8 with center p_{\pm} . If we are not in case (3) of Proposition 3.8, then we get caps C_{\pm} with cylindrical collars of length say $R = 2\underline{R}$. By (4.9) the caps are disjoint, and since $R = 2\underline{R}$ their collars intersect N' with substantial overlap. Thus, the conclusion is that C has the topology of a ball \overline{D}^N . Finally, if we are in case (3) of Proposition 3.8, then we see that actually diam $N' \gg \max{\{\underline{R}_1, \underline{R}_2, \underline{R}_3\}} H^{-1}(p)$ and thus the above argument applies.

Proof of Corollary 1.26. Let K_0 be a smooth compact 2-convex domain in \mathbb{R}^N . By Remark 1.16 it is α -controlled for some α . We choose constants as in Remark 1.24. By Theorem 1.21 there exists an $(\alpha, \delta, \mathbb{H})$ flow $\mathcal{K} = \{K_t\}_{t \in [0,\infty)}$ starting at K_0 . Since \mathcal{K} becomes extinct in finite time (Remark 1.19), by Corollary 1.25 it provides a decomposition of K_0 into a finite connected sum of solid tori $\bar{D}^{N-1} \times S^1$, as claimed. \Box

Proof of Proposition 1.27. It suffices to prove the following claim:

Claim 4.10 (c.f. [Lau13, Prop. 2.2]). For every r > 0 there is a $j_0 < \infty$, such that if $j \ge j_0$, t is a surgery time of \mathcal{K}^j , and $B(p,r) \subseteq K_t^{j,-}$, then $B(p,r) \subseteq K_t^{j,+}$.

Proof. Since $B(p,r) \subseteq K_t^{j,-}$ clearly doesn't fit into a very thin and long neck, c.f. Definition 2.4, we must have $B(p,r) \subseteq K_t^{j,\sharp}$, provided $\bar{\delta} << 1$ and j is sufficiently large. If B(p,r) was contained in a discarded component, then we could find a boundary point with $H \leq (N-1)r^{-1}$. However, by Definition 1.17 the discarded components have $H > H_{\rm th}^j$ everywhere. Thus, $B(p,r) \subseteq K_t^{j,+}$ for j large enough. \Box

As explained in [Lau13], the corollary now follows easily.

Remark 4.11. The same argument applies to more general (α, δ) -flows $\mathcal{K}^j = \{K_t^j \subset \mathbb{R}^N\}_{t \in [0,\infty)}$ with fixed α -controlled initial condition K_0 , assuming only that $s_{\sharp}(\mathcal{K}^j) \to 0, \ \delta \leq \overline{\delta} \ll 1$, and that the minimum of the mean curvature of the discarded components goes to infinity.

APPENDIX A. ONE-SIDED MINIMIZATION

Detailed proof of Proposition 2.9. For convenience of the reader, we now explain the details of the argument of White [Whi00, Sec. 3] and Head [Hea13, Sec. 5]. Let K_{t_0}, K_{t_1} and \overline{B} be as in the statement of the proposition. If t_1 is a surgery time, we first consider the case that K_{t_1} is interpreted as $K_{t_1}^-$. By standard weak compactness and weak lower semicontinuity of perimeter, there is a set $X \subset \mathbb{R}^N$ that minimizes the perimeter (in the following ∂ stands for the reduced boundary)

(A.1)
$$\operatorname{Per}_{\bar{B}}(X) = |\partial X \cap \bar{B}|$$

among all sets of locally finite perimeter satisfying

(A.2)
$$K_{t_1} \subseteq X \subseteq K_{t_0}$$
 and $X \setminus \overline{B} = K_{t_1} \setminus \overline{B}$.

We consider the support of X, which is the closed set

(A.3)
$$\operatorname{spt}(X) = \{x \in \mathbb{R}^N \mid |B(x,r) \cap X| > 0 \text{ for all } r > 0\}.$$

We have to show that $\operatorname{spt}(X) \subseteq K_{t_1}$. Let $\overline{t} \in [t_0, t_1]$ be the supremum of the times $t \in [t_0, t_1]$ such that $\operatorname{spt}(X) \subseteq K_t^-$. Since $\bigcap_{t < \overline{t}} K_t^- = K_{\overline{t}}^-$, we have $\operatorname{spt}(X) \subseteq K_{\overline{t}}^-$. Thus we are done if $\overline{t} = t_1$, so we assume towards a contradiction that $\overline{t} < t_1$. The idea is now to consider suitable mean convex domains $D \subset \mathbb{R}^N$ with $\operatorname{spt}(X) \subseteq D$ (or at least such that $\operatorname{spt}(X) \cap B(x,r) \subseteq D \cap B(x,r)$) and to obtain a contradiction with the 'maximum principle' at any point $x \in \operatorname{spt}(X)$ where ∂D and ∂X touch. To make this rigorous we use the following standard fact from geometric measure theory:

Lemma A.4. Suppose $D \subset \mathbb{R}^N$ is a domain with smooth boundary, $x \in \partial D \setminus K_{t_1}^-$ is a boundary point at which the mean curvature of ∂D is strictly positive, and that for some r > 0 we have the inclusions

 $\operatorname{spt}(X) \cap B(x,r) \subseteq D \cap B(x,r) \subseteq \overline{B} \cap B(x,r).$

Then $x \notin \operatorname{spt}(X)$.

Proof. If $x \in \operatorname{spt}(X)$, then since the open set $B(x,r) \setminus D$ lies in $\mathbb{R}^N \setminus \operatorname{spt}(X)$, it follows that x belongs to the support of the perimeter measure $\operatorname{Per}_{\bar{B}}(X)$. Using a slight deformation of ∂D , one can construct a foliation of a neighborhood of x by mean convex hypersurfaces, such that one of the leaves has strict one-sided exterior contact with D at x. Using a strictly area decreasing deformation one gets a contradiction to the fact that X is minimizing. \Box

For every $x \in \partial \overline{B} \setminus K_{t_1}^-$, we may apply Lemma A.4 with $D = \overline{B}$ to conclude that $x \notin \operatorname{spt}(X)$. Similarly, for every $x \in \partial K_{\overline{t}}^- \setminus \partial K_{t_1}^-$, we may apply the lemma with $D = K_{\overline{t}}^-$, to conclude that $x \notin \operatorname{spt}(X)$. If \overline{t} is not a surgery time, this contradicts the maximality of $\overline{t} < t_1$.

If \bar{t} is a surgery time, consider the post-surgery domain $K^{\sharp}_{\bar{t}} \subseteq K^{-}_{\bar{t}}$ (see Definition 1.3). Suppose p is the center of a surgery neck at time tand scale s such that $B(p, 5\Gamma s) \cap (K_{\overline{t}} \setminus K_{\overline{t}}^{\sharp}) \cap \overline{B} \neq \emptyset$. Suppose $\operatorname{spt}(X) \cap$ $B(p, 10s) \neq \emptyset$. Since $\operatorname{spt}(X) \subseteq K_{\overline{t}}^-$, when Γ is sufficiently large, we may find a mean convex surface (e.g. a slight deformation of the catenoid) that has 1-sided contact with $\operatorname{spt}(X)$ at some point $x \in B(p, \Gamma s) \cap (K_{\overline{t}})$ $K^{\sharp}_{\overline{t}}$). By assumption we have $x \in \overline{B}$, and as argued above, we have $x \in B$. Then Lemma A.4 gives a contradiction. Therefore $spt(X) \cap$ $B(p, 10s) = \emptyset$. By one-sided contact with halfspaces (Lemma A.4) and almost convexity of the caps, we get that $K_{\bar{t}}^{\sharp}$ must be contained in thin neighborhoods of the caps. Since these thin neighborhoods are foliated by mean convex sets, namely the level sets of the distance function, we conclude that $\operatorname{spt}(X) \subseteq K_{\overline{t}}^{\sharp}$, again by Lemma A.4. It follows from the minimality of X that $\operatorname{spt}(X) \cap C = \emptyset$ for any connected component C of $K_{\overline{t}}^{\sharp}$ which is thrown away to form $K_{\overline{t}}^{+}$. Thus $\operatorname{spt}(X) \subseteq K_{\overline{t}}^{+}$. Finally, since $\bar{t} < t_1$, by Lemma A.4 we get that $\operatorname{spt}(X) \cap \partial K_{\bar{t}}^+ = \emptyset$. Since

32

 $K_t \to K_{\bar{t}}^+$ as $t \to \bar{t}^+$, it follows that $\operatorname{spt}(X) \subseteq K_t^-$ for some $t > \bar{t}$. This contradicts the definition of \bar{t} , and we are done.

A similar argument applies if K_{t_1} is interpreted as $K_{t_1}^{\sharp}$ or $K_{t_1}^+$, except that in the case that $\overline{t} = t_1$, one repeats the above argument once more to see that $\operatorname{spt}(X) \subseteq K_{t_1}^{\sharp}$ respectively $\operatorname{spt}(X) \subseteq K_{t_1}^+$. \Box

Remark A.5 (Elementary argument). We now sketch an alternative and more elementary argument. By [HK13a, Rmk 2.6] it is enough to construct a vector field X with div $X \ge 0$, $|X| \le 1$ and X equal to the outward unit normal on ∂K_t . In the case without surgeries and discarded components, such a vector field is given by the outward unit normal of the mean convex foliation. In our case, the vector field given by the outward unit normal might not be defined everywhere, so we have to extend it. We can take care of the discarded pieces by defining X as the negative gradient of the distance to the boundary.⁹ We can take care of the surgeries, by interpolating between the vector field of the cylinder and the vector field of the cap, using a slowly varying cutoff function along the axis of the cylinder. Due to the formula

(A.6)
$$\operatorname{div}(\varphi_1 X_1 + \varphi_2 X_2)$$

= $\varphi_1 \operatorname{div}(X_1) + \varphi_2 \operatorname{div}(X_2) + \langle \nabla \varphi_1, X_1 \rangle + \langle \nabla \varphi_2, X_2 \rangle$,

we can do this preserving positive divergence, provided the cap separation parameter Γ is large enough.

References

[ALM13] B. Andrews, M. Langford, and J. McCoy. Noncollapsing in fully nonlinear curvature flows. Ann. Inst. H. Poincaré ANL, 30(1):23-32, 2013. [And12] B. Andrews. Noncollapsing in mean-convex mean curvature flow. Geom. Topol., 16(3):1413–1418, 2012. [AW94] S. Altschuler and L. Wu. Translating surfaces of the non-parametric mean curvature flow with prescribed contact angle. Calc. Var. PDE, 2(1):101-111, 1994.[BBB+10] L. Bessières, G. Besson, M. Boileau, S. Maillot, and J. Porti. Geometrisation of 3-manifolds. EMS, Zürich, 2010. S. Brendle and G. Huisken. Mean curvature flow with surgery of mean [BH13] convex surfaces in \mathbb{R}^3 . arXiv:1309.1461, 2013. K. Brakke. The motion of a surface by its mean curvature, volume 20 of [Bra78] Mathematical Notes. Princeton University Press, Princeton, N.J., 1978. [Bre13a] S. Brendle. A sharp bound for the inscribed radius under mean curvature flow. arXiv:1309.1459, 2013.

⁹Since the boundary is mean convex, $\operatorname{div} X \ge 0$ in the barrier sense, and thus also in the distributional sense, which is good enough for the calibration argument.

- [Bre13b] S. Brendle. A monotonicity formula for mean curvature flow with surgery. arXiv:1312.0262, 2013.
- [CCG⁺08] B. Chow, S.C. Chu, D. Glickenstein, C. Guenther, J. Isenberg, T. Ivey, D. Knopf, P. Lu, F. Luo, and L. Ni. *The Ricci flow: techniques and applications. Part II.* AMS, Providence, 2008.
- [CGG91] Y.G. Chen, Y. Giga, and S. Goto. Uniqueness and existence of viscosity solutions of generalized mean curvature flow equations. J. Differential Geom., 33(3):749–786, 1991.
- [CIM13] T. Colding, T. Ilmanen, and W. Minicozzi. Rigidity of generic singularities of mean curvature flow. arXiv:1304.6356, 2013.
- [CM12] T. Colding and W. Minicozzi. Generic mean curvature flow I; generic singularities. Ann. of Math. (2), 175(2):755–833, 2012.
- [CM13] T. Colding and W. Minicozzi. Uniqueness of blowups and Lojasiewicz inequalities. arXiv:1312.4046, 2013.
- [CY07] B.-L. Chen and L. Yin. Uniqueness and pseudolocality theorems of the mean curvature flow. Comm. Anal. Geom., 15(3):435–490, 2007.
- [CZ06] H.-D. Cao and X.-P. Zhu. A complete proof of the Poincaré and geometrization conjectures—application of the Hamilton-Perelman theory of the Ricci flow. Asian J. Math., 10(2):165–492, 2006.
- [Eck04] K. Ecker. *Regularity theory for mean curvature flow*. Birkhäuser, Boston, 2004.
- [EH91] K. Ecker and G. Huisken. Interior estimates for hypersurfaces moving by mean curvature. *Invent. Math.*, 105(3):547–569, 1991.
- [ES91] L. Evans and J. Spruck. Motion of level sets by mean curvature. I. J. Differential Geom., 33(3):635–681, 1991.
- [Ham97] R. Hamilton. Four-manifolds with positive isotropic curvature. Comm. Anal. Geom., 5(1):1–92, 1997.
- [Hea13] J. Head. On the mean curvature evolution of two-convex hypersurfaces. J. Differential Geom., 94(2):241–266, 2013.
- [HH13] R. Haslhofer and O. Hershkovits. Ancient solutions of the mean curvature flow. arXiv:1308.4095, 2013.
- [HK13a] R. Haslhofer and B. Kleiner. Mean curvature flow of mean convex hypersurfaces. arXiv:1304.0926, 2013.
- [HK13b] R. Haslhofer and B. Kleiner. On Brendle's estimate for the inscribed radius under mean curvature flow. arXiv:1309.3231, 2013.
- [HS99a] G. Huisken and C. Sinestrari. Mean curvature flow singularities for mean convex surfaces. *Calc. Var. PDE*, 8(1):1–14, 1999.
- [HS99b] G. Huisken and C. Sinestrari. Convexity estimates for mean curvature flow and singularities of mean convex surfaces. Acta Math., 183(1):45– 70, 1999.
- [HS09] G. Huisken and C. Sinestrari. Mean curvature flow with surgeries of two-convex hypersurfaces. *Invent. Math.*, 175(1):137–221, 2009.
- [Hui84] G. Huisken. Flow by mean curvature of convex surfaces into spheres. J. Differential Geom., 20(1):237–266, 1984.
- [KL08] B. Kleiner and J. Lott. Notes on Perelman's papers. Geom. Topol., 12(5):2587–2855, 2008.
- [Lau13] J. Lauer. Convergence of mean curvature flows with surgery. Comm. Anal. Geom., 21(2):355–363, 2013.

- [MT07] J. Morgan and G. Tian. *Ricci flow and the Poincaré conjecture*, volume 3 of *Clay Mathematics Monographs*. AMS, Providence, 2007.
- [Per02] G. Perelman. The entropy formula for the Ricci flow and its geometric applications. arXiv:math/0211159, 2002.
- [Per03] G. Perelman. Ricci flow with surgery on three-manifolds. arXiv:math/0303109, 2003.
- [Whi00] B. White. The size of the singular set in mean curvature flow of mean convex sets. J. Amer. Math. Soc., 13(3):665–695, 2000.
- [Whi03] B. White. The nature of singularities in mean curvature flow of meanconvex sets. J. Amer. Math. Soc., 16(1):123–138, 2003.
- [Whi11] B. White. Subsequent singularities in mean-convex mean curvature flow. arXiv:1103.1469, 2011.

COURANT INSTITUTE OF MATHEMATICAL SCIENCES, NEW YORK UNIVERSITY, 251 MERCER STREET, NEW YORK, NY 10012, USA

E-mail: robert.haslhofer@cims.nyu.edu, bkleiner@cims.nyu.edu