

EQUIVARIANT UNIRATIONALITY OF FANO THREEFOLDS

IVAN CHELTSOV, YURI TSCHINKEL, AND ZHIJIA ZHANG

ABSTRACT. We study unirationality of actions of finite groups on Fano threefolds.

1. INTRODUCTION

In this paper, we consider generically free regular actions of finite groups G on smooth projective varieties X over an algebraically closed field k of characteristic zero. Simplest such actions arise from linear representations, i.e., as projectivizations $\mathbb{P}(V)$ of linear representations V of G . We are interested in birational properties of G -actions, e.g.,

- **(L)** *linearizability*: equivariant birationality to a linear action,
- **(SL)** *stable linearizability*: linearizability of $X \times \mathbb{P}^m$, with trivial action on the second factor.

These notions should be viewed as equivariant analogs of rationality and stable rationality in geometry over nonclosed fields: there are many similarities but also striking differences between these theories, see, e.g., [27], [28] for an introduction to this circle of problems.

Here, we explore an equivariant version of unirationality:

- **(U)** *G -unirationality*: existence of a dominant equivariant rational map

$$\mathbb{P}(V) \dashrightarrow X,$$

where V is a G -representation.

Clearly,

$$(\mathbf{L}) \Rightarrow (\mathbf{SL}) \Rightarrow (\mathbf{U}).$$

In the definition of **(U)**, one can replace $\mathbb{P}(V)$ by V , a G -representation, i.e., **(U)** is equivalent to *very versality* of the G -action, in the terminology of, e.g., [18].

A necessary condition for G -unirationality is existence of fixed points upon restriction to abelian subgroups:

- **Condition (A)**: for every abelian subgroup $H \subseteq G$ one has

$$X^H \neq \emptyset.$$

This should be viewed as analogous to the existence of rational points, in the framework of birational geometry over nonclosed fields. Note that there do exist linearizable G -actions without G -fixed points, moreover, the existence of G -fixed points is not an equivariantly birational property, for nonabelian G . Thus, some standard unirationality constructions in geometry over nonclosed fields fail to apply in the equivariant setting.

Duncan proved that Condition (A) is also sufficient for del Pezzo surfaces of degree ≥ 3 , with generically free actions [17, Theorem 1.4]. The cases of del Pezzo surfaces of degree 2 and 1 remain open.

We turn to smooth Fano threefolds. These are classified by

- $r \in \{1, \dots, 10\}$ – rank of the Picard group,
- $i \in \{1, 2, 3, 4\}$ – Fano index,
- $d = (-K_X)^3/i^3$ – degree.

Their rationality over algebraically closed fields is *almost* settled. There are still outstanding open problems concerning stable rationality and unirationality, e.g., stable rationality of cubic threefolds and unirationality of general quartic threefolds. Rationality properties of geometrically rational Fano threefolds have received a lot of attention, see [20], [21], [31], [30]. There is also a wealth of results concerning (stable) linearizability of group actions on (possibly singular) Fano threefolds, see, e.g., [33], [11], [10], [8], [23], [5], [13].

In this paper, we focus on equivariant unirationality of Fano threefolds of index ≥ 2 ; these should be viewed as 3-dimensional analogs of del Pezzo surfaces. As in the work of Duncan on del Pezzo surfaces [17], Fano threefolds of *small* degree are currently out of reach – we do not even know the unirationality of any smooth Fano threefold of index 2 and degree 1 (all other Fano threefolds of index ≥ 2 are known to be unirational, over algebraically closed fields). Our principal results are:

- A generically free G -action on a smooth quadric threefold is stably linearizable if and only if it satisfies Condition (A), Theorem 4.1.
- A generically free G -action on a smooth cubic threefold satisfying Condition (A) is G -unirational, with the possible exception of $G = C_9 \rtimes C_3$, acting on the Fermat cubic threefold, and $C_5 \rtimes C_{11}$, $\mathrm{PSL}_2(\mathbb{F}_{11})$, acting on the Klein cubic threefold, and a 1-parameter family of cubics with an \mathfrak{A}_5 -action, Theorem 5.1.
- A generically free action on a cubic threefold with isolated singularities, satisfying Condition (A), is G -unirational, Theorem 6.1.

This class contains cubic threefolds with cohomological obstructions to stable linearizability, see [10], [8].

- A generically free G -action on a smooth complete intersection of two quadrics $X \subset \mathbb{P}^5$ is G -unirational if and only if it satisfies Condition **(A)**, which turns out to be equivalent to $X^G \neq \emptyset$, Theorem 7.1.
- A generically free G -action on the (unique) smooth del Pezzo threefold V_5 of degree 5 is linearizable except for $G = \mathfrak{A}_5$, which is not linearizable, by [9], but stably linearizable, by [6, Example 7], see Remark 7.2.

Here is the roadmap of the paper: in Section 2 we recall notions of G -birational geometry, with a focus on G -unirationality. In Section 3 we present several general unirationality constructions, applicable in the equivariant context. In Section 4 we specialize to quadric threefolds. Then we proceed to treat smooth cubics in Section 5, singular cubics in Section 6, and intersections of two quadrics in Section 7.

Acknowledgments: We are grateful to Andrew Kresch and Brendan Hassett for their interest and suggestions. The first author was partially supported by the Leverhulme Trust grant RPG-2021-229 and by EPSRC grant EP/Y033485/1. He also thanks CIRM, Luminy, for its hospitality and for providing a perfect work environment. The second author was partially supported by NSF grant 2301983.

2. GENERALITIES

Throughout, we work over an algebraically closed field k of characteristic zero; all varieties are assumed to be irreducible over k . A G -variety is a projective variety with a generically free regular G -action; we will also consider situations when the G -action is *not* generically free. The G -fixed point locus of a G -variety X will be denoted by X^G . For smooth projective G -varieties X and *abelian* subgroups $H \subseteq G$, existence of H -fixed points is a birational property; this is not the case for nonabelian groups! Note that cyclic actions on rationally connected G -varieties always have fixed points, but this need not be the case for actions of abelian groups, e.g., the C_2^2 -action on \mathbb{P}^1 .

If $\tilde{X} \dashrightarrow X$ is a dominant morphism of G -varieties and \tilde{X} has G -fixed points, then so does X . Since *linear* actions of abelian groups always have fixed points, a necessary condition for G -unirationality of an action of a finite group G on a smooth projective variety X is Condition **(A)**: existence of fixed points upon restriction to every abelian subgroup.

Linearizability and stable linearizability of generically free actions of finite groups on rational surfaces and rational Fano threefolds have been considered in, e.g., [23], [6], [11], [8], [24]. The related notion of G -unirationality, or, equivalently very versality, plays an important role in the study of *essential dimension*, see, e.g., [18], [34]. However, only few concrete examples of G -unirational varieties failing (stable) linearizability are known, see [17] for a systematic treatment of del Pezzo surfaces of degree ≥ 3 .

The bridge between equivariant geometry and geometry over nonclosed fields is provided via *torsors*: For a field extension K/k and a G -torsor T over K , let ${}^T X$ be the T -twist of X over K . A key result is:

Theorem 2.1. [18, Theorem 1.1] *Let X be a G -variety over k . Then*

- *the G -action on X is stably linearizable if and only if for every field extension K/k and every G -torsor T over K , the twist ${}^T X$ is stably rational over K ,*
- *the G -action on X is unirational if and only if every twist ${}^T X$ as above is unirational over K .*

We will apply this in subsequent sections to quadric and cubic hypersurfaces.

A related notion that plays a role in equivariant geometry is the *Amitsur* group $\text{Am}(X, G)$, defined in [3, Section 6] as the image of G -invariant divisor classes

$$\text{Pic}(X)^G \xrightarrow{\delta_2} \text{H}^2(G, k^\times),$$

where δ_2 is the functorial homomorphism arising from the Leray spectral sequence, see, e.g., [26, Section 3]. By [3, Theorem 6.1], this is a G -birational invariant, which vanishes for linear actions and as soon as $X^G \neq \emptyset$. The vanishing of $\text{Am}(X, G)$ implies that every G -invariant divisor class in $\text{Pic}(X)$ is represented by a G -linearized line bundle on X .

All possibilities for $\text{Am}(X, G)$ for G -surfaces X have been determined in [3, Proposition 6.7]. In particular, this group is trivial for del Pezzo surfaces of degree ≤ 6 .

Lemma 2.2. *Let X be a G -variety with nontrivial Amitsur invariant. Then X is not G -unirational.*

Proof. Replacing $\mathbb{P}(V)$ by a G -variety in the same equivariant birational class, we are reduced to the assertion, that an equivariant morphism $Y \rightarrow X$ of smooth projective G -varieties gives rise to an inclusion

$$\text{Am}(X, G) \subseteq \text{Am}(Y, G).$$

This is clear from the functoriality of the map δ_2 in the basic exact sequence

$$0 \rightarrow \mathrm{Hom}(G, k^\times) \rightarrow \mathrm{Pic}(X, G) \rightarrow \mathrm{Pic}(X)^G \xrightarrow{\delta_2} \mathrm{H}^2(G, k^\times),$$

where $\mathrm{Pic}(X, G)$ is the group of isomorphism classes of G -linearized line bundles on X , see, e.g., [29, Section 3]. \square

Example 2.3. Let $X = \mathbb{P}^1$ with the action of the Klein four group $\bar{G} = C_2^2$; it is fixed point free. The Amitsur invariant $\mathrm{Am}(X, \bar{G}) = \mathbb{Z}/2$ is nontrivial, and the \bar{G} -action is not stably linearizable. However, X admits a regular action of the dihedral group $G = \mathfrak{D}_4$ (of order 8), with generic stabilizer C_2 . As a variety with G -action, X is clearly G -unirational: we have a dominant G -equivariant map $V \dashrightarrow X$, where V is an irreducible 2-dimensional representation of G . Of course, every abelian subgroup of G has fixed points on X .

Recall the definition of *Bogomolov multiplier*:

$$\mathrm{B}_0(G) := \mathrm{Ker}(\mathrm{H}^2(G, k^\times) \rightarrow \bigoplus_A \mathrm{H}^2(A, k^\times)),$$

where A runs over all abelian subgroups of G . In fact, it suffices to consider *bicyclic* subgroups [4], and to treat Sylow subgroups, one prime at a time. There is an extensive literature on this invariant, see, e.g., [29, Section 3]. In Example 2.3, we have $\mathrm{B}_0(G) = 0$. More generally, Bogomolov multiplier vanishes when

- G is a p -group of order $|G| \leq p^4$,
- G is an extension of a cyclic group by an abelian group [4, Lemma 4.9], [29, Lemma 3.1],
- G is a split extension of a bicyclic group by an abelian group [29, Lemma 3.2].

For a smooth projective G -variety X , Condition **(A)** implies that

$$\mathrm{Am}(X, G) \subseteq \mathrm{B}_0(G).$$

Lemma 2.4. *Let G be a finite group with $\mathrm{B}_0(G) = 0$. Assume that G acts regularly on \mathbb{P}^n . This G -action is unirational if and only if it satisfies Condition **(A)**.*

Proof. Under our assumption, one has $\mathrm{Am}(\mathbb{P}^n, G) \subseteq \mathrm{B}_0(G) = 0$. \square

Here we do not assume that the G -action on \mathbb{P}^n is generically free. Note that for G with $\mathrm{B}_0(G) \neq 0$, there exist regular generically free actions on projective spaces, which satisfy Condition **(A)** but are *not* linear, and not G -unirational.

Lemma 2.5. *Let*

$$X := \mathbb{P}_B(\mathcal{E})$$

be the projectivization of a vector bundle $\mathcal{E} \rightarrow B$, where B is a smooth projective variety. Assume that X carries a generically free regular G -action such that the canonical projection

$$\pi : X \rightarrow B$$

is G -equivariant. Assume that

- (1) $H^1(G, \text{Pic}(B)) = 0$,
- (2) X satisfies Condition **(A)**, and
- (3) $B_0(G) = 0$.

Then the G -action lifts to \mathcal{E} .

Proof. Consider the sequence of G -modules

$$0 \rightarrow \text{Pic}(B) \xrightarrow{\pi^*} \text{Pic}(X) \rightarrow \mathbb{Z} \rightarrow 1.$$

Note that we do not assume that the G -action on B is generically free.

From the long exact sequence

$$0 \rightarrow \text{Pic}(B)^G \xrightarrow{\pi^*} \text{Pic}(X)^G \rightarrow \mathbb{Z} \rightarrow H^1(G, \text{Pic}(B))$$

and assumption (1), we obtain that there is a G -invariant class in $\text{Pic}(X)$ surjecting onto the class of the relative $\mathcal{O}(1)$. By assumption (2), every line bundle on X admits a linearization upon restriction to abelian subgroups, i.e., $\text{Am}(X, G) \subseteq B_0(G)$. By assumption (3), that latter group is trivial, which implies that every invariant line bundle on X is G -linearizable, in particular, the G -action lifts to \mathcal{E} . \square

3. UNIRATIONALITY CONSTRUCTIONS

General results linking G -unirationality of actions on quadric and cubic hypersurfaces to geometry over nonclosed fields are in [18, Section 10]. We complement these considerations by providing explicit equivariant unirationality constructions in many cases; this allows us to essentially settle unirationality of G -actions on quadric and cubic threefolds, in subsequent sections.

We start with obvious observations:

- G -unirationality is compatible with taking products;
- existence of a dominant G -equivariant rational map $X \dashrightarrow Y$ from a G -unirational X implies G -unirationality of Y .

The following is an analog of a standard result in the theory of quadrics over nonclosed fields, see, e.g., [19, Chapter 4].

Proposition 3.1. *Let G be a finite group and V a representation of G giving rise to a generically free G -action on a smooth quadric hypersurface $X \subset \mathbb{P}(V)$, of dimension ≥ 2 . Assume that there is a G -invariant irreducible subvariety $S \subset X$, which is G -unirational. Then X is G -unirational.*

Here, we allow S to have a nontrivial generic stabilizer, i.e., we require that there is a dominant rational map $\mathbb{P}(W) \dashrightarrow S$, for some G -representation W .

Proof. By assumption, there is a dominant G -equivariant rational map $\mathbb{P}(W) \dashrightarrow S$. Consider the rational map

$$S \times \mathbb{P}(V) \dashrightarrow X,$$

sending a pair of points $(s, p) \in S \times \mathbb{P}(V)$ to the second intersection point of X with the line $\ell(s, p)$ through s and p . This is clearly dominant and G -equivariant, ensuring the G -unirationality of X . \square

We continue with general G -unirationality constructions for cubic hypersurfaces of dimensions ≥ 2 . Over nonclosed fields K of characteristic zero unirationality of smooth cubic hypersurfaces with $X(K) \neq \emptyset$ is classical; generalizations to singular cubic hypersurfaces that are not cones can be found in [25].

Proposition 3.2. *Let V be a representation of a finite group G . Let $X \subset \mathbb{P}(V)$ be a G -invariant irreducible cubic hypersurface which is not a cone. Assume that the G -action on X is generically free and that $X^G \neq \emptyset$. Then X is G -unirational.*

Proof. This is essentially [18, Corollary 10.6], extended to singular cubic hypersurfaces. In particular, if G fixes a singular point, then the G -action is linearizable. If G fixes a smooth point, then the same argument in [18, Corollary 10.6] applies. \square

Proposition 3.3. *Let V be a representation of a finite group G . Let $X \subset \mathbb{P}(V)$ be a G -invariant cubic hypersurface which is not a cone. Assume that the G -action on X is generically free and that G has an index-2 subgroup H such that X is H -unirational. Then X is G -unirational.*

Proof. This is essentially [17, Theorem 3.2], see also [17, Remark 3.3]: G -unirationality is equivalent to K -unirationality of all twists of X over K (via G -torsors), and for cubic hypersurfaces, unirationality over quadratic extensions of K is equivalent to unirationality over K . \square

Corollary 3.4. *Let V be a representation of a finite group G . Let $X \subset \mathbb{P}(V)$ be a G -invariant cubic hypersurface which is not a cone. Assume*

that the G -action on X is generically free and that the set of singular points of X contains a G -orbit of length one or two. Then the G -action on X is G -unirational.

Proof. In the first case, projection from the G -fixed singular point (of multiplicity two) shows that the action is linearizable. In the second case, let $\mathfrak{p}_1, \mathfrak{p}_2$ be the singular points, switched by G . Then there exists an index-2 (normal) subgroup $H \subset G$ fixing \mathfrak{p}_1 and \mathfrak{p}_2 , in particular, X is H -linearizable. Applying Proposition 3.3, X is G -unirational. \square

Proposition 3.5. *Let V be a faithful n -dimensional representation of a finite group G with $n \geq 5$. Let $X \subset \mathbb{P}(V)$ be a G -invariant cubic hypersurface which is not a cone. Assume that the G -action on X is generically free and that there is a G -invariant irreducible hyperplane section $S \subset X$, which is G -unirational. Then X is G -unirational.*

Here we do not require that the G -action on S is generically free.

Proof. Let $\mathcal{T} \rightarrow X^\circ$ be the tangent bundle over the smooth locus X° of X , and $\mathcal{T}|_{S^\circ}$ the restriction of \mathcal{T} to the smooth locus of S . Then $\mathcal{T}|_{S^\circ}$ is a rank- $(n-2)$ vector bundle and the G -action lifts to $\mathcal{T}|_{S^\circ}$. By [14, Theorem 4.2] and [15, Theorem 2.5], we know that the intersection of the tangent space at the generic point of S° with X is not a cone. Thus, each point in the fiber above a general point $\mathfrak{p} \in S^\circ$ canonically corresponds to a line in $\mathbb{P}(V)$ that is tangent to X at \mathfrak{p} with multiplicity 2, and thus intersects X at a unique point $\mathfrak{q} \neq \mathfrak{p}$. Sending each point in $\mathcal{T}|_{S^\circ}$ to the corresponding point $\mathfrak{q} \in X$, we obtain a G -equivariant rational map

$$\varphi : \mathcal{T}|_{S^\circ} \dashrightarrow X. \quad (3.1)$$

We show that φ is dominant. Assume that

$$X = \{f_3(x_1, \dots, x_n) = 0\} \quad \text{and} \quad S = X \cap \{x_1 = 0\}.$$

Given a general point $\mathfrak{q} = [q_1 : \dots : q_n] \in X$, we can find general $p_1, \dots, p_{n-1} \in k$ such that

$$\sum_{i=1}^n \frac{\partial f_3}{\partial x_i}(0, p_1, \dots, p_{n-1}) q_i = f_3(0, p_1, \dots, p_{n-1}) = 0.$$

Then the line passing through \mathfrak{q} and $\mathfrak{p} = [0 : p_1 : \dots : p_{n-1}]$ is tangent to X at \mathfrak{p} . It follows that (3.1) is dominant.

It remains to show that $\mathcal{T}|_{S^\circ}$ is G -unirational. When S has no generic stabilizer, the no-name lemma implies that $\mathcal{T}|_{S^\circ}$ is G -equivariantly birational to $S \times \mathbb{A}^{n-2}$ with trivial action on \mathbb{A}^{n-2} . It follows that $\mathcal{T}|_{S^\circ}$

is G -unirational as S is. When S has a generic stabilizer, we apply the no-name lemma to the vector bundle

$$\mathcal{T}|_{S^\circ} \times V \rightarrow S^\circ \times V,$$

which yields a G -equivariant birationality

$$\mathcal{T}|_{S^\circ} \times V \sim_G S \times V \times \mathbb{A}^{n-2},$$

with trivial action on \mathbb{A}^{n-2} . By our assumptions, $\mathcal{T}|_{S^\circ} \times V$ is G -unirational, implying the G -unirationality of $\mathcal{T}|_{S^\circ}$. \square

Remark 3.6. This argument generalizes to other positive-dimensional G -invariant subvarieties S , under appropriate geometric assumptions, see Section 6.

4. QUADRICS

Stable linearizability of certain G -quadric threefolds $X \subset \mathbb{P}^4$ has been considered in [24, Sections 5 and 6]: all but one subgroup G of the Weyl group $W(D_5)$, acting via signed permutations on the diagonal quadric

$$\sum_{i=1}^5 x_i^2 = 0,$$

are stably linearizable, provided Condition **(A)** is satisfied. The only open case was $G = \mathfrak{D}_4$, acting on

$$\mathbb{P}(\chi_1 \oplus \chi_2 \oplus \chi_3 \oplus V_2),$$

where V_2 is the unique faithful 2-dimensional representation, and χ_1, χ_2, χ_3 are distinct characters of G . There are also many actions that do not arise from subgroups of $W(D_5)$, e.g., an action of \mathfrak{D}_8 on the quadric

$$x_1^2 + x_2x_3 + x_4x_5 = 0,$$

arising from an action on

$$\mathbb{P}(\mathbf{1} \oplus V_2 \oplus V_2'),$$

where V_2 is a faithful representation and V_2' a nonfaithful 2-dimensional representation, see [24, Remark 6.6]. The following theorem covers these and other actions, settling the stable linearizability problem for actions of finite groups on smooth quadric threefolds. The key observation is that, when Condition **(A)** is satisfied, there always exists a G -invariant unirational quadric surface or conic in X and Proposition 3.1 applies.

Theorem 4.1. *Let $X \subset \mathbb{P}^4$ be a smooth quadric threefold, with a generically free action of a finite group G . Then the G -action is stably linearizable if and only if it satisfies Condition **(A)**.*

The rest of this section is devoted to the proof of this theorem.

By Theorem 2.1, stable linearizability of the G -action is equivalent to stable rationality of every twist ${}^T X$ over every field extension K/k . For quadrics, (stable) rationality and unirationality are equivalent over any field. Again by Theorem 2.1, this is equivalent to G -unirationality of the action. By [18, Theorem 10.2], G -unirationality is equivalent to G_2 -unirationality, where G_2 is the 2-Sylow subgroup of G . We may also assume that G is nonabelian: abelian group actions on X satisfying Condition **(A)** are linearizable via the projection from a fixed point.

Thus, to show Theorem 4.1, we now assume that G is a nonabelian 2-group and $X \subset \mathbb{P}(V)$, where V is a faithful representation of G . Since G is a 2-group, we have the following cases:

- (1) $V = \chi_1 \oplus \chi_2 \oplus \chi_3 \oplus V_2$, where V_2 is an irreducible G -representation, and χ_j are characters of G for $j = 1, 2, 3$,
- (2) $V = \chi_1 \oplus V_2 \oplus V'_2$, where V_2 and V'_2 are irreducible G -representations,
- (3) $V = \chi_1 \oplus V_4$, where V_4 is an irreducible representation of G .

We proceed with a case-by-case analysis.

Case (1): Up to isomorphism, we may assume that X is given by

$$X = \{x_1^2 + x_2^2 + x_3^2 + x_4x_5 = 0\} \subset \mathbb{P}^4,$$

with G acting via characters on x_1, x_2, x_3 , and via the irreducible representation V_2 on x_4, x_5 . We claim that Condition **(A)** for X implies Condition **(A)** for the G -action on the conic $C \subset \mathbb{P}^2 = \mathbb{P}(\chi_1 \oplus \chi_2 \oplus \chi_3)$, given by $x_4 = x_5 = 0$. Note that the G -action on this \mathbb{P}^2 has a nontrivial generic stabilizer, since G is nonabelian, and the effective action on the conic C may not satisfy Condition **(A)**.

Since V_2 is irreducible, there exist elements of the form

$$\text{diag}(\pm 1, \pm 1, \pm 1, a, a^{-1}) \in G,$$

where a is a 2-power root of unity of order $\text{ord}(a) \geq 4$. Then

$$\varepsilon_1 = \text{diag}(1, 1, 1, -1, -1) \in G.$$

Note that

$$X^{(\varepsilon_1)} = C \cup \mathfrak{p}_1 \cup \mathfrak{p}_2,$$

where

$$\mathfrak{p}_1 = [0 : 0 : 0 : 1 : 0], \quad \mathfrak{p}_2 = [0 : 0 : 0 : 0 : 1].$$

Let $A \subset G$ be an abelian subgroup containing ε_1 , fixing points on X but not on C . Then A fixes $\mathfrak{p}_1, \mathfrak{p}_2$, and acts via C_2^2 on C . It follows that A contains

$$\varepsilon_2 = \text{diag}(-1, 1, 1, a_1, a_1^{-1}), \quad \varepsilon_3 = \text{diag}(1, -1, 1, b_1, b_1^{-1}),$$

for some 2-power roots of unity a_1, b_1 . Depending on a_1 and b_1 , the elements ε_2 and ε_3 generate at least one of the following elements

$$\text{diag}(-1, 1, 1, -1, -1), \quad \text{diag}(1, -1, 1, -1, -1), \quad \text{diag}(1, 1, -1, -1, -1).$$

Without loss of generality, assume that A contains

$$\varepsilon_4 = \text{diag}(-1, 1, 1, -1, -1).$$

By the irreducibility of V_2 , G also contains an element switching \mathfrak{p}_1 and \mathfrak{p}_2 . Up to multiplying by ε_2 and ε_3 , we may assume that G contains

$$\sigma : (x_1, \dots, x_5) \mapsto (x_1, -x_2, x_3, a_2x_5, a_2^{-1}x_4),$$

for some a_2 . The subgroup

$$\langle \varepsilon_1, \varepsilon_4, \sigma \rangle \simeq C_2^3$$

does not fix points on X .

Let $A \subset G$ be an abelian subgroup not containing ε_1 , fixing points on X but not on C . Then $\langle A, \varepsilon_1 \rangle$ is an abelian group whose fixed locus is $\{\mathfrak{p}_1, \mathfrak{p}_2\}$ or empty. In the first case, we repeat the argument above. We conclude that Condition **(A)** for the G -action on X implies Condition **(A)** for the (not generically free) G -action on C .

Next, we observe that G preserves the set $\{\mathfrak{p}_1, \mathfrak{p}_2\}$, which yields an exact sequence

$$0 \rightarrow N \rightarrow G \rightarrow C_2 \rightarrow 0,$$

where N is the maximal subgroup fixing \mathfrak{p}_1 and \mathfrak{p}_2 , acting diagonally on x_1, \dots, x_5 . Thus N is abelian. By [29, Lemma 3.1], $B_0(G) = 0$, since G is an extension of a cyclic group by an abelian group. By Lemma 2.4, C is G -unirational since Condition **(A)** is satisfied on C . By Proposition 3.1, X is G -unirational.

Case (2): We may assume that G acts on

$$X = \{x_1^2 + x_2x_3 + x_4x_5 = 0\},$$

via χ_1 on x_1 , via V_2 on x_2, x_3 , and via V_2' on x_4, x_5 . By the irreducibility of V_2 and V_2' , G acts on $\mathbb{P}(V_2)$ and $\mathbb{P}(V_2')$ via dihedral groups. Let

$$\mathfrak{p}_1 = [0 : 1 : 0 : 0 : 0], \quad \mathfrak{p}_2 = [0 : 0 : 1 : 0 : 0],$$

$$\mathfrak{p}_3 = [0 : 0 : 0 : 1 : 0], \quad \mathfrak{p}_4 = [0 : 0 : 0 : 0 : 1].$$

Since G leaves invariant the sets $\{\mathfrak{p}_1, \mathfrak{p}_2\}$, and $\{\mathfrak{p}_3, \mathfrak{p}_4\}$ respectively, there is an exact sequence

$$0 \rightarrow N \rightarrow G \rightarrow Q \rightarrow 0, \tag{4.1}$$

where $Q = C_2$ or C_2^2 , depending on the induced action on $\{\mathfrak{p}_1, \mathfrak{p}_2, \mathfrak{p}_3, \mathfrak{p}_4\}$, and $N \subset G$ is the maximal subgroup fixing these points; N consists of elements

$$\varepsilon_1 = \text{diag}(1, t_1, t_1^{-1}, t_2, t_2^{-1}), \quad t_1, t_2 \text{ are 2-power roots of unity.} \quad (4.2)$$

The (nontrivial) center of G acts via scalars on V_2, V_2' , i.e., it must contain one of the following

$$\varepsilon_2 = \text{diag}(1, -1, -1, 1, 1), \quad \varepsilon_3 = \text{diag}(1, 1, 1, -1, -1), \quad \varepsilon_2\varepsilon_3.$$

If $Q = C_2^2 = \langle(1, 2), (3, 4)\rangle$, we may assume that G also contains

$$\begin{aligned} m_1 &: (x_1, \dots, x_5) \mapsto (x_1, a_1x_3, a_1^{-1}x_2, a_2x_4, a_2^{-1}x_5), \\ m_2 &: (x_1, \dots, x_5) \mapsto (x_1, b_1x_2, b_1^{-1}x_3, b_2x_5, b_2^{-1}x_4), \end{aligned}$$

where m_1 is a lift of $(1, 2)$ and m_2 is a lift of $(3, 4)$. After a change of variables, we have $a_1 = b_2 = 1$, and a_2 and b_1 are 2-power roots of unity:

$$m_1^2 = \text{diag}(1, 1, 1, a_2^2, 1/a_2^2), \quad m_2^2 = \text{diag}(1, b_1^2, 1/b_1^2, 1, 1).$$

If $a_2^2 \neq 1$ and $b_1^2 \neq 1$, then G contains ε_2 and ε_3 . The abelian group

$$\langle \varepsilon_2, \varepsilon_3, m_1m_2 \rangle \simeq C_2^3 \quad (4.3)$$

does not fix points on X . If $a_2^2 = 1$ and $b_1^2 \neq 1$, then $\varepsilon_2 \in G$, and N contains an element ε_1 as in (4.2) with $t_2 \neq \pm 1$ since otherwise the representation on x_4, x_5 is reducible. If $\text{ord}(b_1) \geq \text{ord}(t_1)$, or $\text{ord}(t_2) \geq \text{ord}(t_1)$, then ε_3 is generated by m_2^2 and ε_1 , and G contains the C_2^3 from (4.3) with no fixed points. If $\text{ord}(t_1) > \text{ord}(b_1), \text{ord}(t_2)$, one can find $r_1, r_2, r_3 \in \mathbb{Z}$ such that

$$\begin{aligned} m_1^{r_1}\varepsilon_1^{r_2} &: (x_1, \dots, x_5) \mapsto (x_1, c_1x_3, c_1^{-1}x_2, -x_4, -x_5), \\ m_2\varepsilon_1^{r_3} &: (x_1, \dots, x_5) \mapsto (x_1, x_2, x_3, c_2x_5, c_2^{-1}x_4) \end{aligned}$$

for some c_1, c_2 . Then

$$\langle m_1^{r_1}\varepsilon_1^{r_2}, m_2\varepsilon_1^{r_3}, \varepsilon_2 \rangle \simeq C_2^3$$

does not fix points on X . By symmetry, the same applies when $b_1^2 = 1, a_2^2 \neq 1$.

We are reduced to the case $a_2^2 = b_1^2 = 1$, with $\langle m_1, m_2 \rangle \simeq C_2^2$. By the irreducibility of V_2 , N contains an ε_1 as in (4.2) of order at least 4. We repeat the argument above to find an abelian group failing Condition **(A)**, after multiplying m_1 and m_2 by ε_1 .

Thus, if the G -action on X satisfies Condition **(A)**, we have

$$Q = C_2 = \langle(1, 2)(3, 4)\rangle$$

and G contains a lift of $(1, 2)(3, 4)$ of the form

$$m_3 : (x_1, \dots, x_5) \mapsto (x_1, d_1x_3, d_1^{-1}x_2, d_2x_5, d_2^{-1}x_4)$$

for some d_1, d_2 . We discuss cases based on the center of G :

- If $\varepsilon_2, \varepsilon_3 \in G$, then

$$\langle \varepsilon_2, \varepsilon_3, m_3 \rangle \simeq C_2^3$$

has no fixed points on X .

- If $\varepsilon_2\varepsilon_3 \in G$, then every abelian subgroup of G fixes a point on the quadric $S = X \cap \{x_1 = 0\}$. Otherwise, we can find an abelian group without fixed points by adjoining $\varepsilon_2\varepsilon_3$.
- If $\varepsilon_2 \in G$ and $\varepsilon_3 \notin G$, then every abelian subgroup $A \subset G$ fixes a point on $C = X \cap \{x_2 = x_3 = 0\}$. Otherwise, $A' = \langle A, \varepsilon_2 \rangle$ fixes $\{\mathfrak{p}_1, \mathfrak{p}_2\}$, and does not surject onto Q , which implies that it also fixes $\mathfrak{p}_3, \mathfrak{p}_4 \in C$, contradiction.
- If $\varepsilon_3 \in G$ and $\varepsilon_2 \notin G$, the same argument shows that every abelian subgroup of G fixes a point on $C = X \cap \{x_4 = x_5 = 0\}$.

Note that we have $B_0(G) = 0$, since G is an extension of C_2 by an abelian group. In the last two cases, the corresponding conic is G -unirational by Lemma 2.4, and X is G -unirational by Proposition 3.1. In the second case, the G -action on the quadric $S = \mathbb{P}^1 \times \mathbb{P}^1$ does not switch the two rulings. By Lemma 2.4, we can lift the G -action on each \mathbb{P}^1 factor to \mathbb{A}^2 , i.e., the G -action on S to $\mathbb{A}^2 \times \mathbb{A}^2 = \mathbb{A}^4$. It follows that S is G -unirational. By Proposition 3.1, X is also G -unirational.

Case (3): We show that Condition **(A)** is never satisfied. Since V_4 is irreducible, there are no fixed points in $\mathbb{P}(V_4)$, and $S := \mathbb{P}(V_4) \cap X$ is smooth. We may assume that

$$X = \{x_1^2 + x_2x_5 - x_3x_4 = 0\} \subset \mathbb{P}^4 = \mathbb{P}(\mathbf{1} \oplus V_4),$$

with G acting trivially on x_1 , and via V_4 on x_2, x_3, x_4, x_5 . Since V_4 is faithful and G preserves X , the center of G is C_2 , generated by

$$\varepsilon_1 = \text{diag}(1, -1, -1, -1, -1).$$

Finite 2-subgroups of $\text{Aut}(S) = \text{Aut}(\mathbb{P}^1 \times \mathbb{P}^1)$ are subgroups of $\mathfrak{D}_n \wr C_2$, for n some power of 2. This yields 4 distinguished points

$$\mathfrak{p}_1 = [0 : 1 : 0 : 0 : 0], \quad \mathfrak{p}_2 = [0 : 0 : 1 : 0 : 0],$$

$$\mathfrak{p}_3 = [0 : 0 : 0 : 1 : 0], \quad \mathfrak{p}_4 = [0 : 0 : 0 : 0 : 1],$$

and an exact sequence

$$0 \rightarrow N \rightarrow G \rightarrow Q \rightarrow 0,$$

where N fixes each \mathfrak{p}_i . Since V_4 is irreducible, the image $Q \subset \mathfrak{S}_4$, acting via permutations on $\{\mathfrak{p}_1, \mathfrak{p}_2, \mathfrak{p}_3, \mathfrak{p}_4\}$, can be C_2^2, C_4 , or \mathfrak{D}_4 . The subgroup

N is generated by elements of the form

$$\text{diag}(1, t_2, t_1, t_1^{-1}, t_2^{-1}), \quad t_1, t_2 \text{ are 2-power roots of unity.}$$

We proceed to analyze each case.

- $Q = C_2^2$, with lifts of $(1, 2)(3, 4)$ and $(1, 3)(2, 4)$:

$$m_1 : (x_1, \dots, x_5) \mapsto (x_1, a_2 x_3, a_1 x_2, -a_1^{-1} x_5, -a_2^{-1} x_4), \quad (4.4)$$

$$m_2 : (x_1, \dots, x_5) \mapsto (x_1, b_2 x_4, b_1 x_5, -b_1^{-1} x_2, -b_2^{-1} x_3),$$

for some a_1, a_2, b_1, b_2 . Put $s_1 = a_1 a_2, s_2 = b_2 / b_1$ so that

$$m_1^2 = \text{diag}(1, s_1, s_1, 1/s_1, 1/s_1),$$

$$m_2^2 = \text{diag}(1, -s_2, -1/s_2, -s_2, -1/s_2).$$

Since G is a 2-group, s_1 and s_2 are 2-power roots of unity. When the orders $\text{ord}(s_1), \text{ord}(s_2) \geq 4$, we can find integers r_1, r_2 such that

$$s_2^{r_2} = \zeta_4, \quad s_1^{r_1} = \zeta_4^3.$$

Setting

$$\varepsilon_3 := \varepsilon_1^{r_2} m_1^{2r_1} m_2^{2r_2} = \text{diag}(1, 1, -1, -1, 1),$$

we have $S^{(\varepsilon_1, \varepsilon_3)} = \{\mathfrak{p}_1, \dots, \mathfrak{p}_4\}$. The abelian group

$$\langle \varepsilon_3, m_1 m_2, \varepsilon_1 \rangle \simeq C_2^3 \quad (4.5)$$

has no fixed points on X . Thus, s_1 or s_2 equals ± 1 . By symmetry, we may assume that $s_2 = \pm 1$. Then the line

$$\mathfrak{l} := \{x_1 = x_2 + x_3 \sqrt{a_2/a_1} = x_4 \pm x_5 \sqrt{a_2/a_1} = 0\} \subset \mathbb{P}(V)$$

is $\langle m_1, m_2 \rangle$ -invariant. By the irreducibility of V_4 , the kernel N contains an element ε_2 not leaving \mathfrak{l} invariant. In particular, ε_2 takes the form

$$\varepsilon_2 = \text{diag}(1, t_2, t_1, t_1^{-1}, t_2^{-1}) \neq \varepsilon_1, \quad (4.6)$$

for some 2-power roots of unity t_1, t_2 such that $t_1 \neq t_2$. If $\text{ord}(t_2/t_1) \geq 4$, we repeat the argument above, replacing m_1 by $\varepsilon_2 m_1$ to find a C_2^3 with no fixed points on X . Thus we are reduced to $t_1 = -t_2$.

Similarly, if $s_1 = \pm 1$, we are reduced to $t_1 = -1/t_2$. In particular, for $s_1, s_2 \in \{\pm 1\}$, we are reduced to $t_1, t_2 \in \{\pm 1\}$. In this case, G contains ε_3 and a C_2^3 with no fixed points as in (4.5).

We consider the case when $s_1 \neq \pm 1$. As is discussed, we may assume $t_1 = -t_2$. If $\text{ord}(t_1) \leq \text{ord}(s_1)$, then $\varepsilon_3 \in \langle \varepsilon_2, m_1^2 \rangle \subset G$, and G contains a C_2^3 with no fixed points. In the opposite case, there is a $d_1 \in \mathbb{Z}$ such that

$$t_1^{2d_1} = (-1)^{d_1} / s_1.$$

Setting $d_2 = (3 - s_2)/2$, we find that

$$\langle m_1 \varepsilon_2^{d_1}, m_2 \varepsilon_2^{d_2}, \varepsilon_1 \rangle \simeq C_2^3$$

has no fixed points on X .

- $Q = \mathfrak{D}_4$, with G containing m_1, m_2 as in (4.4). If G contains

$$\varepsilon_3 = \text{diag}(1, 1, -1, -1, 1),$$

there is a C_2^3 with no fixed points on X . To show this, we are reduced to the case when $s_1 = \pm 1$ or $s_2 = \pm 1$, as above. Consider the lift of $(1, 3, 4, 2) \in \mathfrak{S}_4$:

$$m_3 : (x_1, \dots, x_5) \mapsto (x_1, c_2 x_4, c_1 x_2, -c_1^{-1} x_5, -c_2^{-1} x_3),$$

for some c_1, c_2 . Observe that

$$\begin{aligned} (m_2 m_3)^2 &= \text{diag}(1, c_2^2/b_1^2, 1, 1, b_1^2/c_2^2), \\ (m_1 m_3)^2 &= \text{diag}(1, 1, a_2^2 c_1^2, 1/(a_2^2 c_1^2), 1). \end{aligned} \quad (4.7)$$

We may assume that $c_1 = \pm 1/a_2$ and $c_2 = \pm b_1$, otherwise ε_3 can be generated by elements in (4.7). Assume $c_1 = 1/a_2$ and $c_2 = b_1$ (the other cases are similar). If $s_1 \neq -s_2$, ε_3 is some power of $m_1 m_2 m_3^2$. When $s_1 = -s_2 = \pm 1$, we find an $\langle m_1, m_2, m_3 \rangle$ -invariant line:

$$\mathfrak{l} = \{x_1 = x_2 + \frac{b_1}{a_1} x_5 = x_3 \mp b_1 a_1 x_4 = 0\} \subset V_4.$$

Thus N contains an ε_2 as in (4.6) which does not leave \mathfrak{l} invariant. This implies that $t_1^2, t_2^2 \neq 1$. As above, when $t_1 \neq \pm t_2$ or $t_1 t_2 \neq \pm 1$, we have $\varepsilon_3 \in G$. When $t_1 = \pm t_2$ and $t_1 t_2 = \pm 1$, we know $t_1^2 = t_2^2 = -1$. Then $\varepsilon_3 = (m_1 m_3 \varepsilon_2)^2 \in G$.

- $Q = C_4$, with G containing a lift m_3 of $(1, 3, 4, 2) \in \mathfrak{S}_4$ as in (4.7). Once we have $\varepsilon_3 \in G$, the abelian group

$$\langle \varepsilon_1, \varepsilon_3, m_3^2 \rangle \simeq C_2^3$$

has no fixed points on X . It follows from the irreducibility of V_4 that N contains an element ε_2 as in (4.6). When $\text{ord}(t_1) \neq \text{ord}(t_2)$, we know that ε_3 is a power of ε_2 . When $t_1^2 = t_2^2 = 1$, $\varepsilon_2 = \varepsilon_3$. We are reduced to

$$t_1 = \zeta_{2^n}^{2d_1+1}, \quad t_2 = \zeta_{2^n}^{2d_2+1}, \quad \text{for some } n \geq 2, \quad d_1, d_2 \in \mathbb{Z}.$$

We have

$$\varepsilon_2 m_3 \varepsilon_2^{-1} m_3^{-1} = \text{diag}(1, t_2/t_1, t_1 t_2, 1/(t_1 t_2), t_1/t_2).$$

Observe that $\text{ord}(t_2/t_1) \neq \text{ord}(t_1 t_2)$. Indeed,

$$t_2/t_1 = \zeta_{2^n}^{2(d_2-d_1)}, \quad t_1 t_2 = \zeta_{2^n}^{2(d_1+d_2+1)},$$

and $d_2 - d_1$ has a different parity from $d_1 + d_2 + 1$. Thus, either ε_3 or $\varepsilon_1\varepsilon_3$ is a power of $\varepsilon_2m_3\varepsilon_2^{-1}m_3^{-1}$, and thus is in G .

This completes the proof of Theorem 4.1.

5. G -UNIRATIONALITY OF SMOOTH CUBIC THREEFOLDS

In this section, we show:

Theorem 5.1. *Let $X \subset \mathbb{P}^4$ be a smooth cubic threefold with a generically free regular action of a finite group G . Assume that the action satisfies Condition **(A)**. Then X is G -unirational, with the possible exception of the following actions:*

- $G = C_9 \rtimes C_3$, and X is the Fermat cubic threefold.
- $G = \mathrm{PSL}_2(\mathbb{F}_{11})$, or $G = C_{11} \rtimes C_5 \subset \mathrm{PSL}_2(\mathbb{F}_{11})$, and X is the Klein cubic threefold.
- $G = \mathfrak{A}_5$, acting on $X \subset \mathbb{P}(V_5)$, where V_5 is the irreducible 5-dimensional representation. There is a 1-parameter family of such cubics, this is the pencil generated by the Klein and the Segre cubic threefolds.

For applications of results of Section 3 to cubic threefolds, we need to address G -unirationality of *singular* cubic surfaces, a case not considered in [17]. All combinations of singularities of cubic surfaces are known classically, see [7]; their automorphisms are partially classified in [37].

Proposition 5.2. *Let $S \subset \mathbb{P}^3$ be a normal rational cubic surface, with a generically free regular action of a finite group G . If the G -action satisfies condition **(A)** then it is G -unirational.*

Proof. The case of smooth cubic surfaces was settled in [17]. Examining possible combinations of singularities [37, Table 1] and applying Corollary 3.4, we obtain G -unirationality in all cases, with the possible exception of $3A_1$, $3A_2$, and $4A_1$ singularities. In the last case, the Cayley cubic surface, the action is birational to a linear action, via the standard Cremona involution, see [10, Section 5] and [8, Section 7]. The $3A_2$ -cubic surface is toric, and G -equivariantly birational to a del Pezzo surface of degree 6, thus G -unirational when it satisfies Condition **(A)**, by [17].

Any $3A_1$ -cubic surface S with an action of a subgroup $G \subseteq \mathrm{Aut}(S)$ is G -unirational. Indeed, we may assume the singular points of S are

$$[1 : 0 : 0 : 0], \quad [0 : 1 : 0 : 0], \quad [0 : 0 : 1 : 0].$$

By linear algebra, S is given by

$$\{x_1x_2x_3 + (x_1 + x_2 + ax_3)x_4^2 + x_4^3 = 0\}, \quad (5.1)$$

for some $a \neq 0$, with $\text{Aut}(S) = \mathfrak{S}_3$, generated by

$$\begin{aligned}\sigma_1 &: (x_1, x_2, x_3, x_4) \mapsto (x_2, x_1, x_3, x_4), \\ \sigma_2 &: (x_1, x_2, x_3, x_4) \mapsto (ax_3, x_1, \frac{x_2}{a}, x_4).\end{aligned}$$

Indeed, it is clear that $\sigma_1, \sigma_2 \in \text{Aut}(S)$. From the equation, one sees that there are no nontrivial elements in $\text{Aut}(S)$ fixing all three nodes. Therefore, $\text{Aut}(S) = \langle \sigma_1, \sigma_2 \rangle$. Furthermore, $\text{Aut}(S)$ fixes three smooth points on S , implying that S is $\text{Aut}(S)$ -unirational, by Proposition 3.2. \square

The rest of this section is devoted to a proof of Theorem 5.1. Actions of finite groups on smooth cubic threefolds $X \subset \mathbb{P}^4$ have been classified by Wei and Yu in [38]. More precisely, they find all groups that can act, and give examples of cubic threefolds that realize such actions, [38, Theorem 1.1 and Section 3]. We recall the maximal nonabelian groups, and the unique cubic threefolds admitting such actions, in the notation of [38]:

	Cubic	Automorphism
X_1	$\sum_{i=j}^5 x_j^3 = 0$	$C_3^4 \rtimes \mathfrak{S}_5$
X_2	$3(\sqrt{3} - 1)x_1x_2x_3 + \sum_{i=1}^5 x_j^3 = 0$	$((C_3^2 \rtimes C_3) \rtimes C_4) \times \mathfrak{S}_3$
X_5	$x_1^2x_2 + x_2^2x_3 + x_3^2x_4 + x_4^2x_5 + x_5^2x_1 = 0$	$\text{PSL}_2(\mathbb{F}_{11})$
X_6	$\sum_{i=1}^6 x_i^3 = \sum_{i=1}^5 x_i = 0$	$C_3 \times \mathfrak{S}_5$

TABLE 1

A table of abelian actions is in [38, Appendix B]. Wei and Yu do not provide normal forms for all cubic threefolds admitting an action of a subgroup from the list in Table 1. We refine their analysis as follows. This relies on `magma`, and supplementary computational material can be found in [12].

We compile a list of all *nonabelian* subgroups of the maximal groups appearing in (1). There are 82 such groups.

We compile a list of groups admitting actions on a *unique* smooth cubic threefold, using [38, Table 2] which lists 12 isomorphisms classes of

abelian groups with actions on a unique cubic; our list contains non-abelian groups with a subgroup isomorphic to one of the distinguished abelian groups. There are 41 such nonabelian groups, up to isomorphism; we note that several of them admit nonconjugated actions on the same cubic – there are 50 conjugacy classes of actions. All the actions are realized on one of the cubic threefolds in Table 1.

For nonabelian groups on the list we check existence of fixed points on X . There are 11 conjugacy classes of actions with fixed points. These actions are unirational by Proposition 3.2. For those without fixed points, we check existence of fixed points for all index-2 subgroups. There are 2 additional classes of actions with fixed points by some index-2 subgroup. These are unirational by Proposition 3.3. Then we check Condition **(A)** and eliminate the groups failing it – these are not G -unirational. There are 31 conjugacy classes of such actions. We are left with 6 groups. They are

$$C_3 \times \mathfrak{S}_5, \quad C_3 \times \mathfrak{A}_5, \quad C_3 \times \mathfrak{F}_5, \quad C_9 \rtimes C_3, \\ C_{11} \rtimes C_5, \quad \mathrm{PSL}_2(\mathbb{F}_{11}),$$

where the first 4 groups act on the Fermat cubic, and the last 2 groups act on the Klein cubic. Among these, the first 3 groups leave invariant a hyperplane section, which is isomorphic to the Clebsch cubic surface, with generic stabilizer C_3 , and a residual action of \mathfrak{S}_5 , \mathfrak{A}_5 , and \mathfrak{F}_5 , respectively. These actions on the Clebsch cubic surface are unirational. By Proposition 3.5, the respective action on the Fermat cubic is unirational. We are left with

$$C_9 \rtimes C_3, \quad C_{11} \rtimes C_5, \quad \mathrm{PSL}_2(\mathbb{F}_{11}).$$

The unirationality of these actions remains open.

We turn to the remaining groups G , i.e., those admitting actions on families of cubics. We write down all possible G -representations V of dimension 5 and compute semi-invariants of degree 3. There are 491 families of cubic threefolds. For each family, we check existence of G -fixed points. There are 170 families with a fixed point, on each member. We do the same for all index-2 subgroups of G ; there are 12 additional families with fixed points. These 182 actions are G -unirational, by Propositions 3.2 and 3.3.

Of the remaining actions, there are 296 violating Condition **(A)** for the generic member. In all these cases, we checked that there is an abelian

group $A \subset G$ with no fixed points on *every* smooth member of the family. These actions are not G -unirational. The following example demonstrates how we checked this.

Example 5.3. Consider $G = \mathbf{SmallGroup}(324, 110)$, with an action on $\mathbb{P}^4 = \mathbb{P}(V)$, with $V = \mathbf{1} \oplus \chi \oplus V_3$, generated by

$$\begin{aligned} & \text{diag}(1, \zeta_3, 1, 1, 1) \\ (\mathbf{x}) & \mapsto (x_1, x_2, x_3, \zeta_3 x_5, \zeta_3^2 x_4), \\ (\mathbf{x}) & \mapsto (x_1, x_2, \zeta_3 x_5, x_3, \zeta_3^2 x_4), \end{aligned}$$

and

$$\begin{aligned} (\mathbf{x}) & \mapsto (x_1, x_2, \frac{\zeta_{12}}{3}((\zeta_6 - 2)x_3 + (1 - 2\zeta_6)x_4 + (\zeta_6 + 1)x_5), \\ & \frac{\zeta_4 + \zeta_{12}}{3}(x_3 + x_4 + x_5), \frac{\zeta_{12}}{3}((1 - 2\zeta_6)x_3 + (\zeta_6 - 2)x_4 + (\zeta_6 + 1)x_5)). \end{aligned}$$

The family of smooth cubic threefolds invariant under G is given by

$$a_1 x_1^2 + a_2 x_2^3 + a_3(x_3^3 + x_4^3 + x_5^3 + 3(\zeta_4 - 2\zeta_{12} - 1)x_3 x_4 x_5) = 0, \quad (5.2)$$

for $a_1, a_2, a_3 \in k$. The subgroup $A \subset G$ generated by

$$\begin{aligned} & \text{diag}(1, 1, \zeta_3, \zeta_3, \zeta_3) \\ (\mathbf{x}) & \mapsto (x_1, \zeta_3^2 x_2, x_4, \zeta_3 x_5, \zeta_3^2 x_4) \end{aligned}$$

is isomorphic to C_3^2 . The fixed locus $(\mathbb{P}^4)^A$ consists of 5 points

$$\begin{aligned} & [1 : 0 : 0 : 0 : 0], \quad [0 : 1 : 0 : 0 : 0], \quad [0 : 0 : \zeta_3 : \zeta_3 : 1], \\ & [0 : 0 : 1 : \zeta_3^2 : 1], \quad [0 : 0 : \zeta_3^2 : 1 : 1]. \end{aligned}$$

One observes that no smooth member of the family (5.2) passes through any of these points. Thus, Condition **(A)** fails for each smooth member of the family.

We are left with 6 isomorphism classes of groups

$$\mathfrak{A}_4, \quad \mathfrak{A}_5, \quad \mathfrak{F}_5, \quad \mathfrak{S}_5, \quad C_3 \times \mathfrak{A}_4, \quad C_3 \times \mathfrak{S}_4,$$

giving rise to 13 actions, up to conjugation:

- \mathfrak{A}_5 admits 2 actions, arising from an irreducible V and a reducible $V = \mathbf{1} \oplus V_4$,
- $C_3 \times \mathfrak{S}_4$ admits 2 actions,
- $C_3 \times \mathfrak{A}_4$ admits 6 actions,

and all others have a unique action. All actions, except the one arising from an *irreducible* 5-dimensional representation of \mathfrak{A}_5 , have the property that $X \subset \mathbb{P}(V)$ admits a G -invariant hyperplane section $S \subset X$, from a decomposition of the representation $V = \mathbf{1} \oplus V_4$, for some faithful 4-dimensional representation V_4 of G .

We observe that the cubic surface S is normal and not a cone. Indeed, S has isolated singularities since X is smooth. If S were a cone, then G would fix a point on S . Using equations of invariant cubics, we see that this is impossible when X is smooth. In particular, we find:

- When G is isomorphic to \mathfrak{A}_5 , \mathfrak{F}_5 , or \mathfrak{S}_5 , there is a unique G -invariant hyperplane section S , which is the Clebsch cubic surface, with trivial generic stabilizer. The G -action on S satisfies Condition **(A)** and is unirational, see [16, Theorem 1.1].
- When G is isomorphic to \mathfrak{A}_4 , there are two invariant hyperplanes. Both of them have trivial generic stabilizer, and the G -actions on them satisfy Condition **(A)**.
- When G is isomorphic to $C_3 \times \mathfrak{A}_4$ or $C_3 \times \mathfrak{S}_4$, there are two invariant hyperplanes. One of them has generic stabilizer C_3 and the G/C_3 -action on the corresponding cubic surface S satisfies Condition **(A)**.

To conclude, in each case when G acts via a reducible representation, there exists an G -invariant normal cubic surface $S \subset X$ which is not a cone such that the effective action of G/G' on S satisfies Condition **(A)**, where G' is the generic stabilizer of S . By Proposition 5.2, the G -action on S is unirational. Applying Proposition 3.5, we see that the G -action on X is unirational.

We are left with the \mathfrak{A}_5 -action with the irreducible representation. Unirationality of this action remains open.

Remark 5.4. Assuming [18, Conjecture 10.4 and Theorem 10.5], G -unirationality of X *would follow* from G_3 -unirationality of X , where G_3 is a 3-Sylow subgroup of G . There are six nonabelian 3-groups contained in (1):

$$\mathrm{He}_3, C_9 \rtimes C_3, C_3 \wr C_3, C_3 \times \mathrm{He}_3, (C_3 \times C_9) \rtimes C_3, C_3 \times (C_3 \wr C_3).$$

The following, obtained via `magma` computations, *would* imply that all G -actions on X , with the possible exception of $C_9 \rtimes C_3$, are G -unirational: Let X be a smooth cubic threefold and $G \subset \mathrm{Aut}(X)$ a 3-group. Then:

- the G -action fixes a point on X , or
- the G -action fails condition **(A)**, or
- X is the Fermat cubic threefold and $G = C_9 \rtimes C_3$ is generated by

$$\mathrm{diag}(1, 1, \zeta_3, \zeta_3^2, 1), \quad \mathrm{diag}(1, 1, 1, \zeta_3, \zeta_3^2), \quad (x_1, \zeta_3 x_2, x_4, \zeta_3 x_5, x_3).$$

6. G -UNIRATIONALITY OF SINGULAR CUBIC THREEFOLDS

This section is devoted to the proof of the following theorem:

Theorem 6.1. *Let X be a cubic threefold with isolated singularities which is not a cone, carrying a generically free regular G -action. Assume that the G -action on X satisfies Condition **(A)**. Then X is G -unirational.*

The proof is based on the classification of configurations of singularities and possible group actions in [11], [10], [8]. In most of the cases, we can apply Propositions 3.2, 3.5, and Corollary 3.4. The remaining cases,

- $3A_1$, $G = \mathrm{GL}_2(\mathbb{F}_3)$,
- $3D_4$, $G = (\mathbb{G}_m^2 \times \mathfrak{S}_3) \rtimes \mathfrak{S}_3$,
- $6A_1$, $G = \mathfrak{S}_4, \mathfrak{S}_5$,

require a separate analysis.

The linearizability properties of singular cubic threefolds have been studied in [10] and [8]; there are linearizable actions, as well as actions failing stable linearizability. The classification results of that paper allow us to address G -unirationality.

By Proposition 3.3, cubic threefolds X with a distinguished singularity or with the following singularity types are G -unirational, for all $G \subseteq \mathrm{Aut}(X)$:

$$\begin{aligned} &2A_n, n = 1, \dots, 5, \quad 2D_4, \quad 2A_2 + 2A_1, \quad 2A_3 + 2A_1, \\ &2D_4 + 2A_1, \quad 2A_2 + 3A_1, \quad 2A_3 + 3A_1, \quad 3A_2 + 2A_1, \quad 2D_4 + 3A_1. \end{aligned}$$

We summarize the analysis of the remaining cases:

$3A_n, 3D_4, n = 1, 2, 3$. With `magma`, we verify that in all cases, except

- (1) $3A_1$, $G = \mathrm{GL}_2(\mathbb{F}_3)$,
- (2) $3D_4$, $G = (\mathbb{G}_m^2 \times \mathfrak{S}_3) \rtimes \mathfrak{S}_3$,

the cubic threefold X contains a G -invariant G -unirational hyperplane section $S \subset X$, a (possibly singular) cubic surface. Indeed, we check that S has fixed points upon restriction of the action to abelian subgroups of the quotient of G by the generic stabilizer of S . Proposition 5.2 implies that S is equivariantly unirational for the action of the quotient, and thus for the action of G . It then follows from Proposition 3.5 that X is G -unirational.

Case 1: $X \subset \mathbb{P}(V)$ is the cubic threefold with $3A_1$ -singularities given by

$$x_1x_2x_3 + x_1q_1 + x_2q_2 + x_3q_3 = 0,$$

where

$$\begin{aligned} q_1 &= x_4^2 + x_4x_5 + x_5^2, \\ q_2 &= x_4^2 + \zeta_3bx_4x_5 + \zeta_3^2x_5^2, \\ q_3 &= x_4^2 + \zeta_3^2bx_4x_5 + \zeta_3x_5^2, \end{aligned}$$

for $b^2 = \sqrt{-2}$, and $V = V_2 \oplus V_3$, for irreducible representations V_2 and V_3 of $G = \mathrm{GL}_2(\mathbb{F}_3)$ generated by

$$\begin{aligned} &\mathrm{diag}(1, 1, 1, -1, -1), \\ (x_1, \dots, x_5) &\mapsto (x_2, x_3, x_1, x_4, \zeta_3x_5), \\ (x_1, \dots, x_5) &\mapsto (\zeta_3x_2, \zeta_3^2x_1, x_3, x_4, \zeta_6x_5), \\ (x_1, \dots, x_5) &\mapsto (x_2, \zeta_6^5x_1, \zeta_6x_3, \frac{\zeta_6bx_4 + x_5}{1 - \zeta_3}, \frac{\zeta_6x_4 + bx_5}{1 - \zeta_3}). \end{aligned}$$

The G -invariant line $\mathfrak{l} = \mathbb{P}(V_2) = \{x_1 = x_2 = x_3 = 0\}$ is contained in the smooth locus of X . Note that \mathfrak{l} is G -unirational, but the G -action is not generically free on \mathfrak{l} . One can check that a general point of \mathfrak{l} is not a *star* point in X , i.e., the intersection of the embedded tangent space at this point with X is not a cone. Applying the argument of Proposition 3.5 to the restriction of the tangent bundle \mathcal{T} of X to \mathfrak{l} , we obtain the G -unirationality of X .

Case 2: X is given by

$$x_1x_2x_3 + x_4^3 + x_5^3 = 0 \tag{6.1}$$

and

$$\mathrm{Aut}(X) = N \times \mathfrak{S}_3, \quad N := \mathbb{G}_m^2 \rtimes \mathfrak{S}_3,$$

where N is acting via \mathfrak{S}_3 -permutation and torus actions on x_1, x_2, x_3 , and \mathfrak{S}_3 via the irreducible 2-dimensional representation on x_4, x_5 . We use notation from [8, Proposition 6.6]

$$N = \langle \tau_{a,b}, \sigma_{(123)}, \sigma_{(12)} \rangle, \quad \mathfrak{S}_3 = \langle \eta, \sigma_{(45)} \rangle, \tag{6.2}$$

where

$$\begin{aligned} \tau_{a,b} &: \mathrm{diag}(a, b, (ab)^{-1}, 1, 1), \\ \eta &: \mathrm{diag}(1, 1, 1, \zeta_3, \zeta_3^2), \end{aligned}$$

and σ are permutations of the variables as is labeled. Consider the subgroup

$$H := \langle \tau_{a,b}, \sigma_{(123)} \cdot \eta \rangle \subset \mathrm{Aut}(X). \tag{6.3}$$

By [8, Proposition 6.6], X is H -equivariantly birational to $S \times \mathbb{P}^1$, where S is the del Pezzo surface of degree 6. In particular, for every $G \subset H$ satisfying Condition **(A)**, we obtain G -unirationality of X .

Lemma 6.2. *Let X be the cubic (6.1), $G \subset \text{Aut}(X)$ a finite subgroup. Then X is G -unirational if and only if Condition **(A)** is satisfied.*

Proof. As discussed, Condition **(A)** is necessary for G -unirationality. We now assume that the G -action on X satisfies Condition **(A)** and proceed to show that X is G -unirational.

Observe that there are two distinguished G -orbits of three points

$$P_1 = \{[1 : 0 : 0 : 0 : 0], [0 : 1 : 0 : 0 : 0], [0 : 0 : 1 : 0 : 0]\},$$

$$P_2 = \{[0 : 0 : 0 : 1 : -1], [0 : 0 : 0 : 1 : -\zeta_3], [0 : 0 : 0 : 1 : -\zeta_3^2]\}.$$

Using Propositions 3.2 and 3.3, we are reduced to the case when G acts on both P_1 and P_2 via C_3 . Up to conjugation in $\text{Aut}(X)$, G contains

$$m_1 := \sigma_{(123)} \cdot \eta.$$

The G -action on P_1 induces a homomorphism $G \rightarrow C_3$. Let N denote its kernel so that we have

$$G = \langle N, m_1 \rangle.$$

If $N \subset \mathbb{G}_m^2(k)$, then $G \subset H$, from (6.3), and X is G -unirational.

If $N \not\subset \mathbb{G}_m^2(k)$, then N contains

$$m_2 := \tau_{a,b} \cdot \eta,$$

for some $\tau_{a,b} \in \mathbb{G}_m^2(k)$, and G is generated by m_1 , m_2 , and a subgroup

$$M = \mathbb{G}_m^2(k) \cap G.$$

Up to replacing m_2 by its power, we may assume that $\text{ord}(\tau_{a,b})$ is a power of 3, so that both a and b are 3-power roots of unity. Condition **(A)** implies that G does not contain either of

$$\eta \quad \text{or} \quad \tau_{\zeta_3, \zeta_3} = \text{diag}(\zeta_3, \zeta_3, \zeta_3, 1, 1),$$

since both abelian groups

$$\langle m_1, \eta \rangle \simeq \langle m_1, \tau_{\zeta_3, \zeta_3} \rangle \simeq C_3^2$$

do not fix points on X . In particular, $(a, b) \neq (1, 1)$.

If $|M|$ is not coprime to 3, then M contains either τ_{1, ζ_3} or $\tau_{\zeta_3, 1}$, which is impossible since

$$\tau_{1, \zeta_3} (m_1 \cdot \tau_{1, \zeta_3} \cdot m_1^{-1})^2 = \tau_{\zeta_3, 1} (m_1^2 \cdot \tau_{\zeta_3, 1} \cdot m_1^{-2})^2 = \tau_{\zeta_3, \zeta_3} \notin G.$$

Hence, $|M|$ is coprime to 3, the order of $\tau_{a,b}$ must be 3, and

$$\tau_{a,b} \in \{\tau_{1,\zeta_3}, \tau_{1,\zeta_3^2}, \tau_{\zeta_3,1}, \tau_{\zeta_3^2,1}, \tau_{\zeta_3,\zeta_3}, \tau_{\zeta_3^2,\zeta_3^2}\}.$$

Conjugating $\tau_{a,b}$ by m_1 , we may assume that

- either $\tau_{a,b} = \tau_{\zeta_3,\zeta_3}$ and $m_2 = \tau_{\zeta_3,\zeta_3} \cdot \eta$, or
- $\tau_{a,b} = \tau_{\zeta_3^2,\zeta_3^2}$ and $m_2 = \tau_{\zeta_3^2,\zeta_3^2} \cdot \eta$.

Without loss of generality, we may assume that $m_2 = \tau_{\zeta_3,\zeta_3} \cdot \eta$. Then G is generated by $m_1 = \sigma_{(123)} \cdot \eta$, m_2 , and a subgroup M of order coprime to 3.

Let $S := X \cap \{x_5 = 0\}$, and $Q := \langle m_1, M \rangle$. Then S is a cubic surface with $3A_2$ -singularities, with a generically free action of Q and generic stabilizer m_2 , i.e., $S = X^{\langle m_2 \rangle}$. Also note that $Q \simeq G / \langle m_2 \rangle$. The Q -action on S satisfies Condition **(A)**. Indeed, if $A \subset Q$ is an abelian subgroup that does not fix any points on S , then $\langle m_2, A \rangle \subset G$ is an abelian subgroup that does not fix points in X . By Theorem 6.1, the Q -action, and thus the G -action on S is unirational, which implies that the G -action on X is unirational, by Proposition 3.2. The Q -unirationality of the surface S also follows from the fact that S is Q -birational to \mathbb{P}^2 with a linear action of Q and classification of all possibilities for the group Q given in [16, Theorem 4.7]. \square

$4A_n, n = 1, 2$, in general position. The singular points are in linear general position. From the equations in [10, Section 5] and [8, Section 7], we see that X contains a G -invariant Cayley cubic surface (give by $x_5 = 0$), which is G -linearizable. Proposition 3.5 shows the G -unirationality of X .

$4A_1$, contained in a plane. Let $\Pi \subset X$ be the plane containing the four nodes. Such X are equivariantly birational to a smooth intersection of two quadrics, discussed in Section 7, with their automorphisms and normal forms. The analysis there allows us to check that a general point on Π is not a star point explicitly via `magma`, in each case. The same argument as in Proposition 3.5 shows G -unirationality of X .

$5A_n, n = 1, 2$. Cubics in this case are equivariantly birational to a smooth quadric threefold. The action is either a subgroup of the \mathfrak{S}_5 -permutation action on coordinates, or a subgroup of the irreducible representation from \mathfrak{A}_5 , which is stably linearizable by Theorem 4.1, see also [24, Proposition 6.2].

$6A_1$. The classification of possible actions in [10, Section 7] yields subcases:

(1) **No plane:** $\text{Aut}(X)$ is one of the following

$$C_2, \mathfrak{S}_3, \mathfrak{S}_4, \mathfrak{D}_4, C_2^2, \mathfrak{D}_6, \mathfrak{S}_3^2 \rtimes C_2, \mathfrak{S}_5.$$

All groups failing Condition **(A)** contain the unique subgroup $C_3^2 \subset \mathfrak{S}_3^2 \rtimes C_2$; they are:

$$C_3^2, \quad C_3 \rtimes \mathfrak{S}_3, \quad C_3 \times \mathfrak{S}_3, \quad \mathfrak{S}_3^2, \quad C_3 \rtimes \mathfrak{S}_3.C_2, \quad \mathfrak{S}_3^2 \rtimes C_2.$$

(2) **One plane:** all actions are linearizable, see [10, Section 7].

(3) **Three planes:** $\text{Aut}(X)$ is one of the following

$$C_2^2, C_2^3, C_2 \times \mathfrak{S}_3, C_2 \times \mathfrak{S}_4,$$

where X is given by

$$x_2x_3x_4 + ax_1^3 + x_1^2(b_1x_2 + b_2(x_3 + x_4)) + x_1(x_2^2 + x_3^2 + x_4^2 - x_5^2) = 0$$

for some $a, b_1, b_2 \in k$.

In Case (3), the point $[0 : 0 : 0 : 0 : 1] \in X$ is fixed by any $G \subseteq \text{Aut}(X)$; thus, X is G -unirational, by Proposition 3.2.

We turn to Case (1). Note that the action on X is not stably linearizable if the invariant class group has rank one, by [10, Proposition 7.5]. To prove unirationality, it suffices to consider the case when $\text{Cl}^G(X) = \mathbb{Z}^2$, by passing to a suitable subgroup of index two, using Proposition 3.3.

This G leaves invariant each class of cubic scrolls on X . Recall from [10, Diagram 7.1] that an invariant divisor class of cubic scrolls shows that there exists a G -equivariant small resolution $X^+ \rightarrow X$ such that X^+ is a weak Fano 3-fold that admits a G -equivariant \mathbb{P}^1 -bundle structure

$$p^+ : X^+ \rightarrow \mathbb{P}^2,$$

whose fibers are mapped to lines in the cubic X . Note that

$$X^+ \simeq \mathbb{P}(\mathcal{E}),$$

where \mathcal{E} is a vector bundle of rank 2 on \mathbb{P}^2 , described in [32, Theorem 3.2(7)]. The weak Fano 3-fold X^+ also appeared in [36, Theorem 3.6].

Note that $H^1(G, \text{Pic}(\mathbb{P}^2)) = 0$ and $B_0(G) = 0$, for all G appearing in this case; the smallest G with nontrivial Bogomolov multiplier has order $|G| = p^5$, see, e.g., [35]. By Lemma 2.4, and Condition **(A)** for X , the base \mathbb{P}^2 is G -unirational. Lemma 2.5 then implies that the G -action lifts to \mathcal{E} . Note that the G -action on the base \mathbb{P}^2 may have generic stabilizers. Nevertheless, applying the no-name lemma as in the proof of Proposition 3.5 yields G -equivariant birationality

$$\mathcal{E} \times V \sim_G \mathbb{P}^2 \times V \times \mathbb{A}^2,$$

where V is a faithful G -representation, with trivial action on \mathbb{A}^2 . This implies G -unirationality of X .

8A₁. In all cases, there is a G -invariant Cayley cubic surface $S \subset X$ (given by $x_3 = 0$ in the equations in [10, Proposition 8.2]), with a G -fixed point on S . It follows that S is G -unirational, and we can apply Proposition 3.5.

9A₁. We recall the normal form, see [10, Section 9]:

$$x_1x_2x_3 - x_4x_5x_6 = a(x_1 + x_2 + x_3) + x_4 + x_5 + x_6 = 0$$

for $a^3 \neq 0, -1$, with G a subgroup of \mathfrak{S}_3^2 , when $a^3 \neq 1$, and of $\mathfrak{S}_3^2 \times C_2$, when $a = 1$. There is an invariant smooth cubic surface $S \subset X$, given by

$$x_1 + x_2 + x_3 = 0.$$

When the G -action on X satisfies Condition (A), then so does the G -action on S . Thus S is G -unirational, and we apply Proposition 3.5.

10A₁. We have a dominant rational map $\text{Gr}(2, 6) \rightarrow X$, the Segre cubic threefold, equivariantly for the action of $G = \mathfrak{S}_6 = \text{Aut}(X)$. By [23, Proposition 19], the \mathfrak{S}_6 -action on $\text{Gr}(2, 6)$ is stably linearizable, thus X is G -unirational.

7. INTERSECTIONS OF TWO QUADRICS

In this section, we prove:

Theorem 7.1. *Let X be a smooth complete intersection of two quadrics in \mathbb{P}^5 and $G \subseteq \text{Aut}(X)$. The following are equivalent:*

- the G -action on X is unirational,
- the G -action on X satisfies condition (A), i.e., every abelian subgroup of G fixes a point on X ,
- the G -action fixes a point on X .

Consider $X = X_{2,2} = Q_1 \cap Q_2 \subset \mathbb{P}^5$, a smooth complete intersection of two quadrics. The problem of classification of actions of finite groups G on X has been addressed in [2]. There is an induced action of a quotient of G on the pencil $\lambda_1 Q_1 + \lambda_2 Q_2$, and we write down equations in each case:

- Type I, \mathfrak{S}_4 : $\sum_{j=0}^4 x_j^2 = x_0^2 + \zeta_4 x_1^2 - x_2^2 - \zeta_4 x_3^2 + x_5^2 = 0$,
- Type II, \mathfrak{D}_6 : $\sum_{j=0}^5 x_j^2 = \sum_{j=0}^5 \zeta_6^j x_j^2 = 0$,
- Type III, \mathfrak{S}_3 : $\sum_{j=0}^5 x_j^2 = a(x_0^2 + \zeta_3 x_1^2 + \zeta_3^2 x_2^2) + x_3^2 + \zeta_3 x_4^2 + \zeta_3^2 x_5^2 = 0$,
- Type IV, C_5 : $\sum_{j=0}^5 x_j^2 = \sum_{j=0}^4 \zeta_5^j x_j^2 = 0$,
- Type V, C_2^2 : $\sum_{j=0}^4 x_j^2 = a(x_0^2 - x_1^2) + x_2^2 - x_3^2 + x_5^2 = 0$,
- Type VI, C_2 : $\sum_{j=0}^5 x_j^2 = a(x_0^2 - x_1^2) + b(x_2^2 - x_3^2) + x_4^2 - x_5^2 = 0$.

We proceed with a list maximal subgroups $G \subseteq \text{Aut}(X)$ satisfying Condition **(A)**.

- Type I: $\mathfrak{S}_3, \mathfrak{D}_4, C_2 \times C_8, C_2^2 \rtimes C_4, C_2 \times \mathfrak{A}_4,$
- Type II: $C_6, \mathfrak{S}_3, C_2 \times C_4, \mathfrak{D}_4, C_2^3, C_2^2 \rtimes C_4, \mathfrak{A}_4 \times C_4,$
- Type III: $\mathfrak{S}_3, \mathfrak{D}_4, C_2^3, C_2 \times \mathfrak{A}_4,$
- Type IV: $C_{10}, C_2^3,$
- Type V: $C_2 \times C_4, \mathfrak{D}_4, C_2^3, C_2^2 \rtimes C_4,$
- Type VI: $\mathfrak{D}_4, C_2^3.$

With `magma`, we find that every subgroup of $\text{Aut}(X)$ satisfying Condition **(A)** has a fixed point on X . We may assume that X does not contain a G -invariant line, since otherwise, the G -action is linearizable, by [22, Theorem 24]. Projecting from the G -fixed point, we find that X is G -birational to a cubic threefold, with singularities of type

$$4A_1, \quad \text{or} \quad 2A_3,$$

containing a unique plane, which must be G -invariant. Using `magma` again, we check a general point on the invariant plane is not a star point, in each case. By results of Section 6, all such cubic threefolds are G -unirational. This proves Theorem 7.1.

It would be interesting to classify actions of finite groups on singular intersections of two quadrics $X_{2,2} \subset \mathbb{P}^5$ and solve the linearization problems for this class of actions. Indeed, these naturally appear in related linearization problems:

Example 7.2. Let V_5 be the unique smooth Fano threefold of index 2 and degree 5 (the quintic del Pezzo threefold), and

$$G \subset \text{Aut}(V_5) \simeq \text{PGL}_2(k)$$

a finite subgroup, not isomorphic to \mathfrak{A}_5 . Then the G -action is linearizable. Indeed, for cyclic or dihedral groups, linearization follows from the construction in [1, Section 5.8]. If $G = \mathfrak{A}_4$, this follows from the proof of [9, Lemma 7.8.2]. If $G = \mathfrak{S}_4$, we have the following G -equivariant commutative diagram:

$$\begin{array}{ccccc}
 & \tilde{V}_5 & \overset{\text{-----}}{\longrightarrow} & \tilde{X}_{2,2} & \\
 & \swarrow \alpha & & \swarrow \gamma & \searrow \delta \\
 V_5 & & & X_{2,2} & \mathbb{P}^2 \\
 & \searrow \beta & & & \\
 & & & &
 \end{array}$$

where

- α is the blowup of the unique G -fixed point,
- β is the contraction of the strict transform of 3 lines in V_5 that pass through the G -fixed point,

- $X_{2,2} \subset \mathbb{P}^5$ is a complete intersection of two quadrics that has three ordinary double points,
- γ is a small resolution of $X_{2,2}$,
- the dashed arrow is a composition of three Atiyah flops, and
- δ is a \mathbb{P}^1 -bundle, with generically free G -action on the base \mathbb{P}^2 .

This commutative diagram appears in [32, Theorem 3.2] and [36, Theorem 3.6]. Linearization of the G -action follows by applying the no-name lemma.

REFERENCES

- [1] C. Araujo, A.-M. Castravet, I. Cheltsov, K. Fujita, A.-S. Kaloghiros, J. Martinez-Garcia, C. Shramov, H. Süß, and N. Viswanathan. *The Calabi problem for Fano threefolds*, volume 485 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 2023.
- [2] A. Avilov. Automorphisms of threefolds that can be represented as an intersection of two quadrics. *Mat. Sb.*, 207(3):3–18, 2016.
- [3] J. Blanc, I. Cheltsov, A. Duncan, and Yu. Prokhorov. Finite quasisimple groups acting on rationally connected threefolds. *Math. Proc. Cambridge Philos. Soc.*, 174(3):531–568, 2023.
- [4] F. A. Bogomolov. The Brauer group of quotient spaces of linear representations. *Izv. Akad. Nauk SSSR Ser. Mat.*, 51(3):485–516, 688, 1987.
- [5] Chr. Böhning, H.-Chr. Graf von Bothmer, and Yu. Tschinkel. Equivariant birational geometry of cubic fourfolds and derived categories, 2023. [arXiv:2303.17678](https://arxiv.org/abs/2303.17678).
- [6] Chr. Böhning, H.-Chr. Graf von Bothmer, and Yuri Tschinkel. Equivariant birational types and derived categories. *Math. Nachr.*, 297(11):4333–4355, 2024.
- [7] J. W. Bruce and C. T. C. Wall. On the classification of cubic surfaces. *J. London Math. Soc. (2)*, 19(2):245–256, 1979.
- [8] I. Cheltsov, L. Marquand, Yu. Tschinkel, and Zh. Zhang. Equivariant geometry of singular cubic threefolds II, 2024. [arXiv:2405.02744](https://arxiv.org/abs/2405.02744).
- [9] I. Cheltsov and C. Shramov. *Cremona groups and the icosahedron*. Monographs and Research Notes in Mathematics. CRC Press, Boca Raton, FL, 2016.
- [10] I. Cheltsov, Yu. Tschinkel, and Zh. Zhang. Equivariant geometry of singular cubic threefolds. *Forum Math. Sigma*, 13:Paper No. e9, 2025.
- [11] I. Cheltsov, Yu. Tschinkel, and Zh. Zhang. Equivariant geometry of the Segre cubic and the Burkhardt quartic. *Selecta Math. (N.S.)*, 31(1):Paper No. 7, 36, 2025.
- [12] I. Cheltsov, Yu. Tschinkel, and Zh. Zhang. Supporting materials, 2025. available at: <https://zhijiazhangz.github.io/equini/>.
- [13] T. Cieurca, S. Tanimoto, and Yu. Tschinkel. Intermediate Jacobians and linearizability, 2024. [arxiv:2403.06047](https://arxiv.org/abs/2403.06047).
- [14] F. Cools and M. Coppens. Star points on smooth hypersurfaces. *J. Algebra*, 323(1):261–286, 2010.
- [15] F. Cools and M. Coppens. Singular hypersurfaces possessing infinitely many star points. *Proc. Amer. Math. Soc.*, 139(10):3413–3422, 2011.

- [16] I. V. Dolgachev and V. A. Iskovskikh. Finite subgroups of the plane Cremona group. In *Algebra, arithmetic, and geometry. In honor of Yu. I. Manin on the occasion of his 70th birthday. Vol. I*, pages 443–548. Boston, MA: Birkhäuser, 2009.
- [17] A. Duncan. Equivariant unirationality of del Pezzo surfaces of degree 3 and 4. *Eur. J. Math.*, 2(4):897–916, 2016.
- [18] A. Duncan and Z. Reichstein. Versality of algebraic group actions and rational points on twisted varieties. *J. Algebr. Geom.*, 24(3):499–530, 2015.
- [19] R. Elman, N. Karpenko, and A. Merkurjev. *The algebraic and geometric theory of quadratic forms*, volume 56 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 2008.
- [20] B. Hassett and Yu. Tschinkel. Rationality of complete intersections of two quadrics over nonclosed fields. *Enseign. Math.*, 67(1-2):1–44, 2021. With an appendix by Jean-Louis Colliot-Thélène.
- [21] B. Hassett and Yu. Tschinkel. Rationality of Fano threefolds of degree 18 over non-closed fields. In *Rationality of varieties*, volume 342 of *Progr. Math.*, pages 237–247. Birkhäuser/Springer, Cham, [2021] ©2021.
- [22] B. Hassett and Yu. Tschinkel. Equivariant geometry of odd-dimensional complete intersections of two quadrics. *Pure Appl. Math. Q.*, 18(4):1555–1597, 2022.
- [23] B. Hassett and Yu. Tschinkel. Torsors and stable equivariant birational geometry. *Nagoya Math. J.*, 250:275–297, 2023.
- [24] B. Hassett and Yu. Tschinkel. Equivariant geometry of low-dimensional quadrics, 2024. [arXiv:2411.00226](https://arxiv.org/abs/2411.00226).
- [25] J. Kollár. Unirationality of cubic hypersurfaces. *J. Inst. Math. Jussieu*, 1(3):467–476, 2002.
- [26] A. Kresch and Yu. Tschinkel. Cohomology of finite subgroups of the plane Cremona group, 2022. [arXiv:2203.01876](https://arxiv.org/abs/2203.01876), to appear in *Algebraic Geom. and Physics*.
- [27] A. Kresch and Yu. Tschinkel. Equivariant birational types and Burnside volume. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, 23(2):1013–1052, 2022.
- [28] A. Kresch and Yu. Tschinkel. Equivariant Burnside groups and representation theory. *Selecta Math. (N.S.)*, 28(4):Paper No. 81, 39, 2022.
- [29] A. Kresch and Yu. Tschinkel. Unramified Brauer group of quotient spaces by finite groups. *J. Algebra*, 664:75–100, 2025.
- [30] A. Kuznetsov and Yu. Prokhorov. Rationality of Fano threefolds over non-closed fields. *Am. J. Math.*, 145(2):335–411, 2023.
- [31] A. Kuznetsov and Yu. Prokhorov. Rationality over nonclosed fields of Fano threefolds with higher geometric Picard rank. *J. Inst. Math. Jussieu*, 23(1):207–247, 2024.
- [32] A. Langer. Fano 4-folds with scroll structure. *Nagoya Math. J.*, 150:135–176, 1998.
- [33] N. Lemire, V. L. Popov, and Z. Reichstein. Cayley groups. *J. Amer. Math. Soc.*, 19(4):921–967, 2006.
- [34] A. S. Merkurjev. Essential dimension: a survey. *Transform. Groups*, 18(2):415–481, 2013.
- [35] P. Moravec. Groups of order p^5 and their unramified Brauer groups. *J. Algebra*, 372:420–427, 2012.

- [36] Th. Peternell P. Jahnke. Almost del Pezzo manifolds. *Adv. Geom.*, 8:387–411, 2008.
- [37] Y. Sakamaki. Automorphism groups on normal singular cubic surfaces with no parameters. *Trans. Amer. Math. Soc.*, 362(5):2641–2666, 2010.
- [38] L. Wei and X. Yu. Automorphism groups of smooth cubic threefolds. *J. Math. Soc. Japan*, 72(4):1327–1343, 2020.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF EDINBURGH, UK

Email address: I.Cheltsov@ed.ac.uk

COURANT INSTITUTE, 251 MERCER STREET, NEW YORK, NY 10012, USA

Email address: tschinkel@cims.nyu.edu

SIMONS FOUNDATION, 160 FIFTH AVENUE, NEW YORK, NY 10010, USA

COURANT INSTITUTE, 251 MERCER STREET, NEW YORK, NY 10012, USA

Email address: zhijia.zhang@cims.nyu.edu