

PDE for Finance, Spring 2003 – Homework 6

Distributed 4/21/03, due 5/5/03. *Includes a hint for problem 4.*

1) This problem develops a continuous-time analogue of the simple Bertsimas & Lo model of “Optimal control of execution costs” presented in the Section 7 notes. The state is (w, p) , where w is the number of shares yet to be purchased and p is the current price per share. The control $\alpha(s)$ is the rate at which shares are purchased. The state equation is:

$$\begin{aligned}dw &= -\alpha ds \text{ for } t < s < T, & w(t) &= w_0 \\dp &= \theta\alpha ds + \sigma dz \text{ for } t < s < T, & p(t) &= p_0\end{aligned}$$

where dz is Brownian motion and θ, σ are fixed constants. The goal is to minimize, among (nonanticipating) controls $\alpha(s)$, the expected cost

$$E \left\{ \int_t^T [p(s)\alpha(s) + \theta\alpha^2(s)] ds + [p(T)w(T) + \theta w^2(T)] \right\}.$$

The optimal expected cost is the value function $u(w_0, p_0, t)$.

(a) Show that the HJB equation for u is

$$u_t + H(u_w, u_p, p) + \frac{\sigma^2}{2}u_{pp} = 0$$

for $t < T$, with Hamiltonian

$$H(u_w, u_p, p) = -\frac{1}{4\theta}(p + \theta u_p - u_w)^2.$$

The final value is of course

$$u(w, p, T) = pw + \theta w^2.$$

(b) Look for a solution of the form $u(w, p, t) = pw + g(t)w^2$. Show that g solves

$$\dot{g} = \frac{1}{4\theta}(\theta - 2g)^2$$

for $t < T$, with $g(T) = \theta$. Notice that u does not depend on σ , i.e. setting $\sigma = 0$ gives the same value function.

(c) Solve for g . (Hint: start by rewriting the equation for g , “putting all the g ’s on the left and all the t ’s on the right”.)

(d) Show by direct examination of your solution that the optimal $\alpha(s)$ is constant.

(Food for thought: what happens if one takes the running cost to be $\int_t^T p(s)\alpha(s) ds$ instead of $\int_t^T p(s)\alpha(s) + \theta\alpha^2(s) ds$?)

2) The Section 7 notes discuss work by Bertsimas, Kogan, and Lo involving least-square replication of a European option. The analysis there assumes all trades are *self-financing*, so the value of the portfolio at consecutive times is related by

$$V_j - V_{j-1} = \theta_{j-1}(P_j - P_{j-1}).$$

Let's consider what happens if trades are permitted to be non-self-financing. This means we introduce an additional control g_j , the amount of cash added to (if $g_j > 0$) or removed from (if $g_j < 0$) the portfolio at time j , and the portfolio values now satisfy

$$V_j - V_{j-1} = \theta_{j-1}(P_j - P_{j-1}) + g_{j-1}.$$

It is natural to add a quadratic expression involving the g 's to the objective: now we seek to minimize

$$E \left[(V_N - F(P_N))^2 + \alpha \sum_{j=0}^{N-1} g_j^2 \right]$$

where α is a positive constant. The associated value function is

$$J_i(V, P) = \min_{\theta_i, g_i, \dots, \theta_{N-1}, g_{N-1}} E_{V_i=V, P_i=P} \left[(V_N - F(P_N))^2 + \alpha \sum_{j=i}^{N-1} g_j^2 \right].$$

The claim enunciated in the Section 7 notes remains true in this modified setting: J_i can be expressed as a quadratic polynomial

$$J_i(V_i, P_i) = \bar{a}_i(P_i)|V_i - \bar{b}_i(P_i)|^2 + \bar{c}_i(P_i)$$

where \bar{a}_i, \bar{b}_i , and \bar{c}_i are suitably-defined functions which can be constructed inductively. Demonstrate this assertion in the special case $i = N - 1$, and explain how $\bar{a}_{N-1}, \bar{b}_{N-1}, \bar{c}_{N-1}$ are related to the functions $a_{N-1}, b_{N-1}, c_{N-1}$ of the Section 7 notes.

3) Consider scaled Brownian motion with drift, $dy = \mu dt + \sigma dw$, starting at $y(0) = 0$. The solution is of course $y = \mu t + \sigma w(t)$, so its probability distribution at time t is Gaussian with mean μt and variance $\sigma^2 t$. Show that solution $\hat{p}(\xi, t)$ obtained by Fourier transform in the Section 8 notes is consistent with this result.

4) Consider scaled Brownian motion with drift and jumps: $dy = \mu dt + \sigma dw + JdN$, starting at $y(0) = 0$. Assume the jump occurrences are Poisson with rate λ , and the jump magnitudes J are Gaussian with mean 0 and variance δ^2 . Find the probability distribution of the process y at time t . (*Hint*: don't try to use the Fourier transform. Instead observe that you know, for any n , the probability that n jumps will occur before time t ; and after conditioning on the number of jumps, the distribution of y is a Gaussian whose mean and variance are easy to determine. Assemble these ingredients to give the density of y as an infinite sum.)