

CSC 220 Algorithms

Midterm Test A, November 4, 2002

1. Using the method of *QUICKSORT*, find the right ('splitting') position of 35 in the unsorted sequence

35, 90, 88, 6, 49, 34, 79, 2, 68.

What is the total number of comparisons you made?

2. Suppose $f(1) = 1$ and for all $n \geq 2$,

$$f(n) = 3 \left(\sum_{i=1}^{n-1} f(n-i) \right) + 3.$$

What is $f(n)$?

3. Design an algorithm for finding simultaneously the two smallest *and* the two largest elements among 16 distinct numbers using at most 28 comparisons.

4. We have 8 coins that look the same. 6 of them have weight 1, one has weight .99 and one has weight 0.98.

a. Design an algorithm for determining the weights of the coins by a *digital scale*, using at most 4 measurements in the worst case.

b. Is it always possible to identify the 6 good (heavy) coins a two-pan *balance*, using at most 3 measurements?

5. a. Let $f(n), g(n), h(n)$ and $j(n)$ be positive valued functions defined on the set of positive integers, and assume that $f(n) = o(g(n))$ and $h(n) = O(j(n))$. Prove or disprove: $f(n) + h^2(n) = O(g(n) + j^2(n))$.

b. Let $f(n) = \frac{n^2+2n+1}{n+2^{-n}}$ and $g(n) = 100n + 3$ for all $n \geq 0$. Prove or disprove: $f(n) = o(g(n))$.

6. Prove that if there in an algorithm for finding the *second smallest* element in a given set of distinct numbers using at most 100 comparisons, then one can also find in the same time simultaneously the *smallest* and the *second smallest* elements using at most 100 comparisons.

Please explain all of your answers! Good luck! - J.P.