

ALGEBRAIC VARIETIES WITH MANY RATIONAL POINTS

YURI TSCHINKEL

CONTENTS

Introduction	1
1. Geometry background	3
2. Existence of points	20
3. Density of points	28
4. Counting problems	35
5. Counting points via universal torsors	63
6. Height zeta functions	71
References	89

INTRODUCTION

Let $f \in \mathbb{Z}[t, x_1, \dots, x_n]$ be a polynomial with coefficients in the integers. Consider

$$f(t, x_1, \dots, x_n) = 0,$$

as an equation in the unknowns t, x_1, \dots, x_n or as an algebraic family of equations in x_1, \dots, x_n parametrized by t . We are interested in integer solutions: in their existence and distribution. Sometimes the emphasis is on individual equations, e.g.,

$$x^n + y^n = z^n,$$

sometimes we want to understand a *typical* equation, i.e., a general equation in some family. To draw inspiration (and techniques) from different branches of algebra it is necessary to consider solutions with values in other rings and fields, most importantly, finite fields \mathbb{F}_q , finite extensions of \mathbb{Q} , or the function fields $\mathbb{F}_p(t)$ and $\mathbb{C}(t)$. While there is a wealth of *ad hoc* elementary approaches to individual equations,

Date: July 21, 2008.

The second author was supported by the NSF grant 0602333.

and deep theories focussing on their visible or hidden symmetries, our primary approach here will be via geometry.

Basic geometric objects are the affine space \mathbb{A}^n and the projective space $\mathbb{P}^n = (\mathbb{A}^{n+1} \setminus 0)/\mathbb{G}_m$, the quotient by the diagonal action of the multiplicative group. Concretely, affine algebraic varieties $X^{\text{affine}} \subset \mathbb{A}^n$ are defined by systems of polynomial equations with coefficients in some base ring R ; their solutions with values in R , $X^{\text{affine}}(R)$, are called R -integral points. Projective varieties are defined by homogeneous equations, and their R -points are equivalence classes of solutions, with respect to diagonal multiplication by nonzero elements in R . If F is the fraction field of R , then $X^{\text{projective}}(R) = X^{\text{projective}}(F)$, and these points are called F -rational points. The geometric advantages of working with “compact” projective varieties translate to important technical advantages in the study of equations, and the theory of rational points is currently much better developed.

The sets $X(F)$ reflect on the one hand the geometric and algebraic complexity of X (e.g., the dimension of X), and on the other hand the structure of the ground field F (e.g., its topology, analytic structure). It is important to consider the variation of $X(F')$, as F' runs over extensions of F , either algebraic extensions, or completions. It is also important to study projective and birational invariants of X , its birational models, automorphisms, fibration structures, deformations. Each point of view contributes its own set of techniques, and it is the interaction of ideas from a diverse set of mathematical cultures that makes the subject so appealing and vibrant.

The focus in these notes will be on smooth projective varieties X defined over \mathbb{Q} , with *many* \mathbb{Q} -rational points. Main examples are varieties \mathbb{Q} -birational to \mathbb{P}^n and hypersurfaces in \mathbb{P}^n of low degree. We will study the relationship between the global *geometry* of X over \mathbb{C} and the distribution of rational points in Zariski topology and with respect to *heights*. Here are the problems we face:

- Existence of solutions: local obstructions, the Hasse principle, global obstructions;
- Density in various topologies: Zariski density, weak approximation;
- Distribution with respect to heights: bounds on smallest points, asymptotics.

Here is the roadmap of the paper. Section 1 contains a summary of basic terms from complex algebraic geometry: main invariants of algebraic varieties, classification schemes, and examples most relevant to arithmetic in dimension ≥ 2 . Section 2 is devoted to the existence of rational and integral points, including aspects of decidability, effectivity, local and global obstructions. In Section 3 we discuss Lang’s conjecture and its converse, focussing on varieties with nontrivial endomorphisms and fibration structures. Section 4 introduces heights, counting functions, and height zeta functions. We explain conjectures of Batyrev, Manin, Peyre and their refinements. The remaining sections are devoted to geometric and analytic techniques employed in the proof of these conjectures: universal torsors, harmonic analysis on adelic groups, p -adic integration and “estimates”.

Acknowledgments. I am very grateful to V. Batyrev, F. Bogomolov, U. Derenthal, A. Chambert-Loir, J. Franke, J. Harris, B. Hassett, A. Kresch, Y. Manin, E. Peyre, J. Shalika, M. Strauch and R. Takloo-Bighash for the many hours of listening and sharing their ideas. Partial support was provided by National Science Foundation Grants 0554280 and 0602333.

sect:geometry

1. GEOMETRY BACKGROUND

We discuss basic notions and techniques of algebraic geometry that are commonly encountered by number theorists. For most of this section, F is an algebraically closed field of characteristic zero. Geometry over algebraically closed fields of positive characteristic, e.g., algebraic closure of a finite field, differs in several aspects: difficulties arising from inseparable morphisms, “unexpected” maps between algebraic varieties, additional symmetries, lack (at present) of *resolution of singularities*. Geometry over nonclosed fields, especially number fields, introduces new phenomena: varieties may have *forms*, not all geometric constructions *descend* to the ground field, parameter counts do not suffice. In practice, it is “equivariant geometry for finite groups”, with Galois-symmetries acting on all geometric invariants and special loci. The case of surfaces is addressed in detail in [Has08].

sect:pic

1.1. Basic invariants. Let X be an algebraic variety over F . We may assume that X is projective and smooth. We seek to isolate invariants of X that are most relevant for arithmetic investigations.

There are two natural types of invariants: *birational* invariants, i.e., invariants of the function field $F(X)$, and *projective geometry* invariants, i.e., those arising from a concrete representation of X as a subvariety of \mathbb{P}^n . Examples are the *dimension* $\dim(X)$, defined as the transcendence degree $F(X)$ over F , and the *degree* of X in the given projective embedding. For hypersurfaces $X_f \subset \mathbb{P}^n$ the degree is simply the degree of the defining homogeneous polynomial. In general, it is defined via the Hilbert function of the homogeneous ideal, or geometrically, as the number of intersection points with a general hyperplane of codimension $\dim(X)$.

The degree alone is not a sensitive indicator of the complexity of the variety: Veronese embeddings of $\mathbb{P}^1 \hookrightarrow \mathbb{P}^n$ exhibit it as a curve of degree n . In general, we may want to consider all possible projective embeddings of a variety X . Two such embeddings can be “composed” via the Segre embedding $\mathbb{P}^n \times \mathbb{P}^m \rightarrow \mathbb{P}^N$, where $N = nm + n + m$. For example, we have the standard embedding $\mathbb{P}^1 \times \mathbb{P}^1 \hookrightarrow \mathbb{P}^3$, with image a smooth quadric. In this way, projective embeddings of X form a “monoid”; the corresponding abelian group is the *Picard group* $\text{Pic}(X)$. Alternatively, it is the group of isomorphism classes of *line bundles* on X . Cohomologically,

$$\text{Pic}(X) = H_{\text{et}}^1(X, \mathbb{G}_m),$$

where \mathbb{G}_m is the sheaf of invertible functions. Yet another description is

$$\text{Pic}(X) = \text{Div}(X) / (\mathbb{C}(X)^* / \mathbb{C}^*),$$

where $\text{Div}(X)$ is the free abelian group generated by codimension one subvarieties of X , and $\mathbb{C}(X)^*$ is the multiplicative group of rational functions of X , each $f \in \mathbb{C}(X)^*$ giving rise to a *principal* divisor $\text{div}(f)$ (divisor of zeroes and poles of f). Sometimes it is convenient to identify divisors with their classes in $\text{Pic}(X)$. Note that Pic is a *contravariant functor*: a morphism $\tilde{X} \rightarrow X$ induces a homomorphism of abelian groups $\text{Pic}(X) \rightarrow \text{Pic}(\tilde{X})$. There is an exact sequence

$$1 \rightarrow \text{Pic}^0(X) \rightarrow \text{Pic}(X) \rightarrow \text{NS}(X) \rightarrow 1$$

where $\text{Pic}^0(X)$ is the connected component of the identity in $\text{Pic}(X)$ and $\text{NS}(X)$ is the *Néron-Severi* group of X . In most applications in this paper, $\text{Pic}^0(X)$ is trivial.

Given a projective variety $X \subset \mathbb{P}^n$, via an explicit system of homogeneous equations, we can easily write down at least one divisor on X , a hyperplane section L in this embedding. Another divisor, the

divisor of zeroes of a differential form of top degree on X , can also be computed from the equations. Its class $K_X \in \text{Pic}(X)$, i.e., the class of the line bundle $\Omega_X^{\dim(X)}$, is called the *canonical class*. In general, it is not known how to write down effectively divisors whose classes are not proportional to linear combinations of K_X and L . This can be done for some varieties over \mathbb{Q} , e.g., smooth cubic surfaces in $X_3 \subset \mathbb{P}^3$ (see Section 1.9), but is already open for smooth quartics $X_4 \subset \mathbb{P}^4$ (for some partial results in this direction, see Section 1.10).

Elements in $\text{Pic}(X)$ corresponding to projective embeddings generate the *ample cone* $\Lambda_{\text{ample}}(X) \subset \text{Pic}(X)_{\mathbb{R}}$; ample divisors arise as hyperplane sections of X in a projective embedding. The closure of $\Lambda_{\text{ample}}(X)$ in $\text{Pic}(X)_{\mathbb{R}}$ is called the *nef cone*. An *effective divisor* is a sum with nonnegative coefficients of irreducible subvarieties of codimension one. Their classes span the *effective cone* $\Lambda_{\text{eff}}(X)$. Divisors giving rise to embeddings of some Zariski open subset of X form the *big cone*. To summarize we have

$$\Lambda_{\text{ample}}(X) \subseteq \Lambda_{\text{nef}}(X) \subseteq \Lambda_{\text{big}}(X) \subseteq \Lambda_{\text{eff}}(X) \subset \text{Pic}(X)_{\mathbb{R}}.$$

These cones and their combinatorial structure encode important geometric information. For example, for all divisors $D \in \Lambda_{\text{nef}}(X)$ and all curves $C \subset X$, the intersection number $D.C \geq 0$ [Kle66]. Divisors on the boundary of $\Lambda_{\text{ample}}(X)$ give rise to fibration structures on X ; we will discuss this in more detail in Section 1.4.

exam:blowup

Example 1.1.1. Let X be a smooth projective variety, $Y \subset X$ a smooth subvariety and $\pi : \tilde{X} = \text{Bl}_Y(X) \rightarrow X$ the *blowup* of X in Y , i.e., the complement in \tilde{X} to the *exceptional divisor* $E := \pi^{-1}(Y)$ is isomorphic to $X \setminus Y$, and E itself can be identified with the projectivized tangent cone to X at Y . Then

$$\text{Pic}(\tilde{X}) \simeq \text{Pic}(X) \oplus \mathbb{Z}E$$

and

$$K_{\tilde{X}} = \pi^*(K_X) + \mathcal{O}((\text{codim}(Y) - 1)E)$$

(see [Har77, Exercise 8.5]). Note that

$$\pi^*(\Lambda_{\text{eff}}(X)) \subset \Lambda_{\text{eff}}(\tilde{X}),$$

but that pullbacks of ample divisors are not necessarily ample.

exam:hypersurface

Example 1.1.2. Let $X \subset \mathbb{P}^n$ be a hypersurface of dimension ≥ 3 and degree d . Then $\text{Pic}(X) = \text{NS}(X) = \mathbb{Z}L$, generated by the class of the

hyperplane section, and

$$\Lambda_{\text{ample}}(X) = \Lambda_{\text{eff}}(X) = \mathbb{R}_{\geq 0}L.$$

The canonical class is

$$K_X = -(n + 1 - d)L.$$

exam:cubic-cone

Example 1.1.3. If X is a smooth cubic surface over an algebraically closed field, then $\text{Pic}(X) = \mathbb{Z}^7$. The anticanonical class is proportional to the sum of 27 exceptional curves (lines):

$$-K_X = \frac{1}{9}(D_1 + \cdots + D_{27}).$$

The effective cone $\Lambda_{\text{eff}}(X) \subset \text{Pic}(X)_{\mathbb{R}}$ is spanned by the classes of the lines.

On the other hand, the effective cone of a minimal resolution of the singular cubic surface

$$x_0x_3^2 + x_1^2x_3 + x_2^3 = 0$$

is a simplicial cone (in \mathbb{R}^7) [HT04].

exam:solvable

Example 1.1.4. Let G be a connected solvable linear algebraic group, e.g., the additive group $G = \mathbb{G}_a$, an algebraic torus $G = \mathbb{G}_m^d$ or the group of upper-triangular matrices. Let X be an equivariant compactification of G , i.e., the action of G on itself by extends to X . Using equivariant resolution of singularities, if necessary, we may assume that X is smooth projective and that the boundary

$$X \setminus G = D = \cup_{i \in \mathcal{I}} D_i, \quad \text{with } D_i \text{ irreducible,}$$

is a divisor with normal crossings. Every divisor D on X is equivalent to a divisor with support in the boundary since it can be “moved” there by the action of G . Thus $\text{Pic}(X)$ is generated by the components D_i , and the relations are given by functions with zeroes and poles supported in D , i.e., by the characters $\mathfrak{X}^*(G)$. We have an exact sequence

eqn:exact-cone

$$(1.1) \quad 0 \rightarrow \mathfrak{X}^*(G) \rightarrow \oplus_{i \in \mathcal{I}} \mathbb{Z}D_i \xrightarrow{\pi} \text{Pic}(X) \rightarrow 0$$

The cone of effective divisors $\Lambda_{\text{eff}}(X) \subset \text{Pic}(X)_{\mathbb{R}}$ is the image of the simplicial cone $\oplus_{i \in \mathcal{I}} \mathbb{R}_{\geq 0}D_i$ under the projection π . The anticanonical class is

$$-K_X = \oplus_{i \in \mathcal{I}} \kappa_i D_i, \quad \text{with } \kappa_i \geq 1, \text{ for all } i.$$

For unipotent G one has $\kappa_i \geq 2$, for all i [HT99].

For higher-dimensional varieties without extra symmetries, the computation of the ample and effective cones, and of the position of K_X with respect to these cones, is a difficult problem. A sample of recent papers on this subject is: [CS06], [Far06], [FG03], [Cas07], [HT03], [HT02], [GKM02]. However, we have the following fundamental result (see also Section 1.4):

`thm:fg-cones`

Theorem 1.1.5. *Let X be a smooth projective variety with $-K_X \in \Lambda_{\text{ample}}(X)$. Then $\Lambda_{\text{nef}}(X)$ is a finitely generated rational cone. If $-K_X$ is big and nef then $\Lambda_{\text{eff}}(X)$ is finitely generated.*

Finite generation of the nef cone goes back to Mori (see [CKM88] for an introduction). The result concerning $\Lambda_{\text{eff}}(X)$ has been proved in [Bat92] in dimension ≤ 3 , and in higher dimensions in [BCHM06] (see also [Ara05], [Leh08]).

`sect:schemes`

1.2. Classification schemes. In some arithmetic investigations (e.g., Zariski density or rational points) we rely mostly on birational properties of X ; in others (e.g., asymptotics of points of bounded height), we need to work in a fixed projective embedding.

Among *birational* invariants, the most important are those arising from a comparison of X with a projective space:

- (1) *rationality*: there exists a birational isomorphism $X \sim \mathbb{P}^n$, i.e., there is an isomorphism of function fields $F(X) = F(\mathbb{P}^n)$, for some $n \in \mathbb{N}$;
- (2) *unirationality*: there exists a dominant map $\mathbb{P}^n \dashrightarrow X$;
- (3) *rational connectedness*: for general $x_1, x_2 \in X(F)$ there exists a morphism $f : \mathbb{P}^1 \rightarrow X$ such that $x_1, x_2 \in f(\mathbb{P}^1)$.

It is easy to see that

$$(1) \Rightarrow (2) \Rightarrow (3).$$

These properties are equivalent in dimension two, but diverge in higher dimensions. First examples of unirational but not rational threefolds were constructed in [IM71] and [CG72]. The approach of [IM71] was to study of the group $\text{Bir}(X)$ of birational automorphisms of X ; finiteness of $\text{Bir}(X)$, i.e., *birational rigidity*, implies nonrationality. No examples of smooth projective rationally connected but not unirational varieties are known.

Interesting unirational varieties arise as quotients V/G , where $V = \mathbb{A}^n$ is a representation space for a faithful action of a linear algebraic group G . For example, the moduli space $\mathcal{M}_{0,n}$ of n points on \mathbb{P}^1 is birational to $(\mathbb{P}^1)^n/\text{PGL}_2$. Moduli spaces of degree d hypersurfaces $X \subset \mathbb{P}^n$

are naturally isomorphic to $\mathbb{P}(\mathrm{Sym}^d(\mathbb{A}^{n+1}))/\mathrm{PGL}_{n+1}$. Rationality of V/G is known as *Noether's problem*. It has a positive solution for G being the symmetric group \mathfrak{S}_n , the group PGL_2 [Kat82], [BK85], and in many other cases [SB89], [SB88]. Counterexamples for some *finite* G were constructed in [Sal84], [Bog87]; nonrationality is detected by the *unramified Brauer group*, $\mathrm{Br}_{\mathrm{un}}(V/G)$, closely related to the Brauer group of the function field $\mathrm{Br}(F(V/G)) = \mathrm{H}_{\mathrm{et}}^2(F(V/G), \mathbb{G}_m)$.

Now we turn to invariants arising from projective geometry, i.e., ample line bundles on X . For smooth curves C , an important invariant is the *genus* $g(C) := \dim \mathrm{H}^0(X, K_X)$. In higher dimensions, one considers the *Kodaira dimension*

$$\boxed{\text{eqn:kodaira-dim}} \quad (1.2) \quad \kappa(X) := \limsup \frac{\log(\dim \mathrm{H}^0(X, nK_X))}{\log(n)},$$

and the related graded *canonical* section ring

$$\boxed{\text{eqn:graded-kx}} \quad (1.3) \quad R(X, K_X) = \bigoplus_{n \geq 0} \mathrm{H}^0(X, nK_X).$$

A fundamental theorem is that this ring is finitely generated [BCHM06]. The Kodaira dimension is the dimension of the variety $\mathrm{Proj}(R(X, K_X))$, or equivalently, the dimension of the image of X under the map

$$X \rightarrow \mathbb{P}(\mathrm{H}^0(X, nK_X)),$$

for sufficiently large n . For K_X ample one has $\kappa(X) = \dim(X)$.

A very rough classification of smooth algebraic varieties is based on the position of the anticanonical class with respect to the cone of ample divisors. Numerically, this is reflected in the value of the Kodaira dimension. There are three main cases:

- *Fano*: $-K_X$ ample;
- *general type*: K_X ample;
- *intermediate type*: none of the above.

For curves, this classification can be read off from the genus: curves of genus 0 are of Fano type, of genus 1 of intermediate type, and of genus ≥ 2 of general type. Other examples of varieties in each group are:

- Fano: \mathbb{P}^n , smooth degree d hypersurfaces $X_d \subset \mathbb{P}^n$, with $d \leq n$;
- general type: hypersurfaces $X_d \subset \mathbb{P}^n$, with $d \geq n + 2$, moduli spaces of curves of high genus and abelian varieties of high dimension;
- intermediate type: \mathbb{P}^2 blown up in 9 points, abelian varieties, Calabi-Yau varieties.

There are only finitely many families of smooth Fano varieties in each dimension [KMM92]. On the other hand, the universe of varieties of general type is boundless and there are many open classification questions already in dimension 2. The qualitative behavior of rational points on X is closely related to this classification (see Section 3). In our arithmetic applications we will mostly encounter Fano varieties and varieties of intermediate type.

In finer classification schemes such as the *Minimal Model Program* (MMP) it is important to take into account fibration structures and mild singularities (see [KMM87] and [Cam04]). Indeed, recall that $R(X, K_X)$ is finitely generated and put $Y = \text{Proj}(R(X, K_X))$. Then the general fiber of the rational projection

$$X \dashrightarrow Y = \text{Proj}(R(X, K_X)),$$

is a (possibly singular) Fano variety. For example, a surface of Kodaira dimension 1 is birational to a Fano fiber space over a curve of genus ≥ 1 .

Analogously, in many arithmetic questions, the passage to fibrations is inevitable (see Section 4.16). These often arise from the section rings

eqn:rxl

$$(1.4) \quad R(X, L) = \bigoplus_{n \geq 0} H^0(X, nL).$$

Consequently, one considers the *Iitaka dimension*

eqn:iitaka-dim

$$(1.5) \quad \kappa(X, L) := \limsup \frac{\log(\dim H^0(X, nL))}{\log(n)}.$$

Finally, a pair (X, D) , where X is smooth projective and D is a divisor in X , gives rise to another set of invariants: the *log Kodaira dimension* $\kappa(X, K_X + D)$ and the *log canonical ring* $R(X, K_X + D)$, whose finite generation is also known in many cases [BCHM06]. Again, one distinguishes

- *log Fano*: $\kappa(X, -(K_X + D)) = \dim(X)$;
- *log general type*: $\kappa(X, K_X + D) = \dim(X)$;
- *log intermediate type*: none of the above.

This classification has consequences for the study of *integral* points on the open variety $X \setminus D$.

sect:sing

1.3. Singularities. Assume that X is \mathbb{Q} -Cartier, i.e., there exists an integer m such that mK_X is a Cartier divisor. Let \tilde{X} be a normal variety and $f : \tilde{X} \rightarrow X$ a proper birational morphism. Denote by E

the f -exceptional divisor and by e its generic point. Let $g = 0$ be a local equation of E . Locally, we can write

$$f^*(\text{generator of } \mathcal{O}(mK_X)) = g^{md(E)}(dy_1 \wedge \dots \wedge dy_n)^m$$

for some $d(E) \in \mathbb{Q}$ such that $md(E) \in \mathbb{Z}$. If, in addition, $K_{\tilde{X}}$ is a line bundle (e.g., \tilde{X} is smooth), then $mK_{\tilde{X}}$ is linearly equivalent to

$$f^*(mK_X) + \sum_i m \cdot d(E_i)E_i; \quad E_i \text{ exceptional,}$$

and numerically

$$K_{\tilde{X}} \sim f^*(K_X) + \sum_i d(E_i)E_i.$$

The number $d(E)$ is called the *discrepancy* of X at the exceptional divisor E . The discrepancy $\text{discr}(X)$ of X is

$$\text{discr}(X) := \inf\{d(E) \mid \text{all } f, E\}$$

If X is smooth then $\text{discr}(X) = 1$. In general, $\text{discr}(X) \in \{-\infty\} \cup [-1, 1]$.

defin:singularities

Definition 1.3.1. The singularities of X are called

- *terminal* if $\text{discr}(X) > 0$ and
- *canonical* if $\text{discr}(X) \geq 0$.

It is essential to remember that *terminal* = smooth in codimension 2 and that for surfaces, *canonical* means *Du Val* singularities.

Canonical isolated singularities on surfaces can be classified via Dynkin diagrams: Let $f : \tilde{X} \rightarrow X$ be the *minimal* desingularization. Then the submodule in $\text{Pic}(\tilde{X})$ spanned by the classes of exceptional curves (with the restriction of the intersection form) is isomorphic to the root lattice of the corresponding Dynkin diagram (exceptional curves give simple roots).

Canonical singularities don't influence the expected asymptotic for rational points on the complement to all exceptional curves: for (singular) Del Pezzo surfaces X we still expect an asymptotic of points of bounded anticanonical height of the shape $\mathbf{B} \log(\mathbf{B})^{9-d}$, where d is the degree of X , just like in the smooth case (see Section 4.12). This fails when the singularities are worse than canonical.

exam:weight

Example 1.3.2. Let $w = (w_0, \dots, w_n) \in \mathbb{N}^n$, with $\text{gcd}(w_0, \dots, w_n) = 1$ and let

$$X = X(w) = \mathbb{P}(w_0, \dots, w_n)$$

be a *weighted projective space*, i.e., we have a quotient map

$$(\mathbb{A}^{n+1} \setminus 0) \xrightarrow{\mathbb{G}_m} X,$$

where the torus \mathbb{G}_m acts by

$$\lambda \cdot (x_0, \dots, x_{n+1}) \mapsto (\lambda^{w_0} x_0, \dots, \lambda^{w_n} x_n).$$

For $w = (1, \dots, 1)$ it is the usual projective space, e.g., $\mathbb{P}^2 = \mathbb{P}(1, 1, 1)$. The weighted projective plane $\mathbb{P}(1, 1, 2)$ has a canonical singularity and the singularity of $\mathbb{P}(1, 1, m)$, with $m \geq 3$, is worse than canonical.

For a discussion of singularities on general weighted projective spaces and so called *fake* weighted projective spaces see, e.g., [Kas08].

sect:mmp

1.4. Minimal Model Program. Here we recall basic notions from the Minimal Model Program (MMP) (see [CKM88], [KM98], [KMM87], [Mat02] for more details). The starting point is the following fundamental theorem due to Mori [Mor82]:

thm:mori

Theorem 1.4.1. *Let X be a smooth Fano variety of dimension n . Then there exists an integer $d \leq n + 1$ such that through every geometric point x of X there passes a rational curve of $-K_X$ -degree $\leq d$.*

These rational curves move in families. Their specializations are rational curves, which may move again, and again, until one arrives at “rigid” rational curves.

Theorem 1.4.2 (Cone theorem). *Let X be a smooth Fano variety. Then the closure of the cone of (equivalence classes of) effective curves in $H_2(X, \mathbb{R})$ is finitely generated by classes of rational curves.*

The generating rational curves are called *extremal rays*, they correspond to codimension-1 faces of the dual cone of nef divisors. Mori’s Minimal Model Program links the convex geometry of the nef cone $\Lambda_{\text{nef}}(X)$ with natural transformations of X . Pick a divisor D on the face dual to an extremal ray $[C]$. It is not ample anymore, but it still defines a map

$$X \rightarrow \text{Proj}(R(X, D)),$$

which contracts the curve C to a point. The map is one of the following:

- a fibration over a base of smaller dimension, and the restriction of D to a general fiber proportional to the anticanonical class of the fiber, which is a (possibly singular) Fano variety,
- a birational map contracting a divisor,
- a contraction of a subvariety in codimension ≥ 2 (a *small* contraction).

The image could be singular, as in Example 1.3.2, and one of the most difficult issues of MMP was to develop a framework which allows to maneuver between birational models with singularities in a restricted class, while keeping track of the modifications of the Mori cone of curves. In arithmetic applications, for example proofs of the existence of rational points as in, e.g., [CTSSD87a], [CTSSD87b], [CTS89], one relies on the *fibration method and descent*, applied to some auxiliary varieties. Finding the “right” fibration is an art. Mori’s theory gives a systematic approach to these questions.

A variant of Mori’s theory, the *Fujita program*, analyzes fibrations arising from divisors on the boundary of the *effective cone* $\Lambda_{\text{eff}}(X)$. In this case, the restriction of D to a general fiber is a perturbation of (some positive rational multiple of) the anticanonical class of the fiber by a *rigid* effective divisor. This theory turns up in the analysis of height zeta functions in Section 6 (see also Section 4.15).

We will need the notion of the *geometric hypersurface of linear growth*:

$$\boxed{\text{eqn:lin-geom}} \quad (1.6) \quad \Sigma_X^{\text{geom}} := \{L \in \text{NS}(X)_{\mathbb{R}} \mid a(L) = 1\}$$

Let X be smooth projective with $\text{Pic}(X) = \text{NS}(X)$ and a finitely generated effective cone $\Lambda_{\text{eff}}(X)$. For a line bundle L on X define

$$\boxed{\text{eqn:a(L)}} \quad (1.7) \quad a(L) := \min(a \mid aL + K_X \in \Lambda_{\text{eff}}(X)).$$

Let $b(L)$ be the maximal codimension of the face of $\Lambda_{\text{eff}}(X)$ containing $a(L)L + K_X$. In particular,

$$a(-K_X) = 1 \quad \text{and} \quad b(-K_X) = \text{rk Pic}(X).$$

These invariants arise in Manin’s conjecture in Section 4.12 and the analysis of analytic properties of height zeta functions in Section 6.1.

$\boxed{\text{sect:campana}}$

1.5. Campana’s program. Recently, Campana developed a new approach to classification of algebraic varieties with the goal of formulating necessary and sufficient conditions for potential density of rational points. The key notions are: the *core* of an algebraic variety and *special* varieties. Special varieties include Fano varieties and Calabi–Yau varieties. They are conjectured to have a potentially dense set of rational points. This program is explained in detail in [Abr08].

$\boxed{\text{sect:cox}}$

1.6. Cox rings. Again, we assume that X is a smooth projective variety with $\text{Pic}(X) = \text{NS}(X)$. Examples are Fano varieties, equivariant compactifications of algebraic groups and holomorphic symplectic varieties. Fix line bundles L_1, \dots, L_r whose classes generate $\text{Pic}(X)$. For

$\mathbf{n} = (n_1, \dots, n_r) \in \mathbb{Z}^r$ we put

$$L^{\mathbf{n}} := L_1^{n_1} \otimes \dots \otimes L_r^{n_r}.$$

The *Cox* ring is the multigraded section ring

$$\text{Cox}(X) := \bigoplus_{\mathbf{n} \in \mathbb{Z}^r} H^0(X, L^{\mathbf{n}}).$$

The key issue is finite generation of this ring. This has been proved under quite general assumptions in [BCHM06, Corollary 1.1.9]. Assume that $\text{Cox}(X)$ is finitely generated. Then both $\Lambda_{\text{eff}}(X)$ and $\Lambda_{\text{nef}}(X)$ are finitely generated polyhedral (see [HK00, Proposition 2.9]). Other important facts are:

- (1) X is a toric variety if and only if $\text{Cox}(X)$ is a polynomial ring [Cox95], [HK00, Corollary 2.10].
- (2) $\text{Cox}(X)$ is multigraded for $\text{NS}(X)$, in particular, it carries a natural action of the dual torus T_{NS} .

sect:cox

1.7. Universal torsors. We continue to work over an algebraically closed field. Let G be a linear algebraic group and X an algebraic variety. A G -torsor over X is a principal G -bundle $\pi : \mathcal{T}_X \rightarrow X$. Basic examples are GL_n -torsors, they arise from vector bundles over X ; for instance, each line bundle L gives rise to a $\text{GL}_1 = \mathbb{G}_m$ -torsor over X . Up to isomorphism, G -torsors are classified by $H_{\text{et}}^1(X, G)$; line bundles are classified by $H_{\text{et}}^1(X, \mathbb{G}_m) = \text{Pic}(X)$. When G is commutative, $H_{\text{et}}^1(X, G)$ is a group.

Let $G = \mathbb{G}_m^d$ be an algebraic torus and $\mathfrak{X}^*(G) = \mathbb{Z}^d$ its character lattice. A G -torsor over an algebraic variety X is determined, up to isomorphism, by a homomorphism

$$\chi : \mathfrak{X}^*(G) \rightarrow \text{Pic}(X).$$

Assume that $\text{Pic}(X) = \text{NS}(X) = \mathbb{Z}^d$ and that χ is in fact an isomorphism. The arising G -torsors are called *universal*. The introduction of universal torsors is motivated by the fact that over *nonclosed* fields they “untwist” the action of the Galois group on the Picard group of X (see Sections 1.13 and 2.5). The “extra dimensions” and “extra symmetries” provided by the torsor add crucial freedom in the analysis of the geometry and arithmetic of the underlying variety. Examples of applications to rational points will be presented in Sections 2.5 and 5. This explains the surge of interest in explicit equations for universal torsors, the study of their geometry: singularities and fibration structures.

Assume that $\text{Cox}(X)$ is finitely generated. Then $\text{Spec}(\text{Cox}(X))$ contains a universal torsor \mathcal{T}_X of X as an open subset.

[Cox95]
 [STV06], [TVAV08], [SS07], [SS08], book [Sko01]
 [HK00]

sect:hyper

1.8. Hypersurfaces. We now turn from the general theory to specific varieties. Let $X \subset \mathbb{P}^n$ be a smooth hypersurface of degree d . We have already described some of its invariants in Example 1.1.2, at least when $\dim(X) \geq 3$. In particular, in this case $\text{Pic}(X) \simeq \mathbb{Z}$. In dimension two, there are more possibilities.

The most interesting cases are $d = 2, 3$, and 4. A quadric X_2 is isomorphic to $\mathbb{P}^1 \times \mathbb{P}^1$ and has Picard group $\text{Pic}(X_2) \simeq \mathbb{Z} \oplus \mathbb{Z}$. A cubic has Picard group of rank 7. These are examples of *Del Pezzo surfaces* discussed in Section 1.9. They are birational to \mathbb{P}^2 . A smooth quartic $X_4 \subset \mathbb{P}^3$ is an example of a *K3 surface* (see Section 1.10). We have $\text{Pic}(X_4) = \mathbb{Z}^r$, with r between 1 and 20. They are not rational and, in general, do not admit nontrivial fibrations.

Cubic and quartic surfaces have a rich geometric structure, with large “hidden” symmetries. This translates into many intricate arithmetic issues.

sect:del-pezzo

1.9. Del Pezzo surfaces. A smooth projective surface X with ample anticanonical class is called a *Del Pezzo surface*. Standard examples are \mathbb{P}^2 and $\mathbb{P}^1 \times \mathbb{P}^1$. Over algebraically closed ground fields, all other Del Pezzo surfaces X_d are obtained as blowups of \mathbb{P}^2 in $9 - d \leq 8$ points in general position (e.g., no three on a line, no 6 on a conic). This d is the anticanonical *degree* of X_d . Del Pezzo surfaces of low degree admit the following realizations:

- $d = 4$: intersection of two quadrics in \mathbb{P}^4 ;
- $d = 3$: hypersurface of degree 3 in \mathbb{P}^3 ;
- $d = 2$: hypersurface of degree 4 in the weighted projective space $\mathbb{P}^2(1, 1, 1, 2)$ given by

$$w^2 = f_4(x, y, z), \quad \text{with } \deg(f_4) = 4.$$

- $d = 1$: hypersurface of degree 6 in $\mathbb{P}(1, 1, 2, 3)$ given by

$$w^2 = t^3 + f_4(x, y)t + f_6(x, y), \quad \text{with } \deg(f_i) = i.$$

Visually and mathematically most appealing are, perhaps, *cubic surfaces* with $d = 3$. Note that for $d = 1$, the anticanonical linear series has one *base point*, in particular, $X_1(F) \neq \emptyset$, over *any* field F .

Let us compute the geometric invariants of a Del Pezzo surface of degree d , expanding the Example 1.1.3. Since $\text{Pic}(\mathbb{P}^2) = \mathbb{Z}L$, the hyperplane class, we have

$$\text{Pic}(X_d) = \mathbb{Z}L \oplus \mathbb{Z}E_1 \oplus \cdots \oplus \mathbb{Z}E_{9-d},$$

where E_i are the full preimages of the blown-up points. The canonical class is computed as in Example 1.1.1

$$K_{X_d} = -3L + (E_1 + \cdots + E_{9-d}).$$

The intersection pairing defines a quadratic form on $\text{Pic}(X_d)$, with $L^2 = 1$, $L \cdot E_i = 0$, $E_i \cdot E_j = 0$, for $i \neq j$, and $E_j^2 = -1$. Let W_d be the subgroup of $\text{GL}_{10-d}(\mathbb{Z})$ of elements preserving K_{X_d} and the intersection pairing. For d small enough there are several other classes with square -1 :

$$L - (E_i + E_j), \quad 2L - (E_1 + \cdots + E_5), \quad \text{etc.}$$

Those classes whose intersection with K_{X_d} is also -1 are called (classes of) *exceptional curves*, these curves are *lines* in the anticanonical embedding. Their number $r(d)$ can be found in the table below. We have

$$-K_{X_d} = c_d \sum_{j=1}^{r(d)} E_j,$$

the sum over all exceptional curves, where $c_d \in \mathbb{Q}$ can be easily computed, e.g., $c_3 = 1/9$. The effective cone is spanned by the $r(d)$ classes of exceptional curves, and the nef cone is the cone dual to $\Lambda_{\text{eff}}(X_d)$ with respect to the intersection pairing on $\text{Pic}(X_d)$. Put

eqn:alpha-cubic

$$(1.8) \quad \alpha(X_d) := \text{vol}(\Lambda_{\text{nef}}(X_d) \cap \{C \mid (-K_X, C) = 1\}).$$

This “volume” of the nef cone has been computed in [?]:

$9 - d$	1	2	3	4	5	6	7	8
$r(d)$	1	3	6	10	16	27	56	240
$\alpha(X_d)$	1/6	1/24	1/72	1/144	1/180	1/120	1/30	1

Given a Del Pezzo surface over a number field, the equations of the lines can be computed effectively. For example, this is easy to see for the diagonal cubic surface

$$x_0^3 + x_1^3 + x_2^3 + x_3^3 = 0.$$

Writing

$$x_i^3 + x_j^3 = \prod_{r=1}^3 (x_i + \zeta_3^r x_j) = x_\ell^3 + x_k^3 = \prod_{r=1}^3 (x_\ell + \zeta_3^r x_k)0,$$

with $i, j, k, l \in [0, \dots, 3]$, and permuting indices we get all 27 lines. In general, equations for the lines can be obtained by solving the corresponding equations on the Grassmannian of lines.

For $9 - d = 3, 4, 5, \dots, 8$, the group W_d is the *Weyl group* of a *root system*:

$$A_1 \times A_2, A_4, D_5, E_6, E_7, E_8,$$

and the root lattice itself is the orthogonal to K_{X_d} in $\text{Pic}(X_d)$, the so-called *primitive* Picard group. Recent work on Cox rings of Del Pezzo surfaces established a geometric connection to Lie groups with root systems of the above type: a universal torsor over a Del Pezzo surface embeds into a certain flag variety for the simply-connected Lie group with the corresponding root system [Der07], [SS07].

Degenerations of Del Pezzo surfaces are also interesting and important. Typically, they arise as special fibers of fibrations, and their analysis is unavoidable in the theory of *models* over rings such as \mathbb{Z} , or $\mathbb{C}[t]$. There is a classification of singular Del Pezzo surfaces with the help of Dynkin diagrams [BW79]. Models of Del Pezzo surfaces over curves are discussed in [Cor96]. Volumes of nef cones of singular Del Pezzo surfaces are computed in [DJT08].

exam:sing-4

Example 1.9.1 (Degree four). Here are some examples of singular degree four Del Pezzo surfaces $X = \{Q_0 = 0\} \cap \{Q = 0\} \subset \mathbb{P}^4$, where $Q_0 = x_0x_1 + x_2^2$ and Q is one of the following quadratic forms

$4A_1$	$x_3x_4 + x_2^2$
$3A_1$	$x_2(x_1 + x_2) + x_3x_4$
$2A_1 + A_2$	$x_1x_2 + x_3x_4$
$A_1 + A_3$	$x_3^2 + x_4x_2 + x_0^2$
$2A_1 + A_3$	$x_0^2 + x_3x_4$
A_3	$x_3^2 + x_4x_2 + (x_0 + x_1)^2$
D_4	$x_3^2 + x_4x_1 + (x_0 + x_2)^2$
D_4	$x_3^2 + x_4x_1 + x_0^2$
D_5	$x_1x_2 + x_0x_4 + x_3^2$
D_5	$x_1^2 + x_1x_2 + x_0x_4 + x_3^2$

For more details see [DP80].

exam:sing-3

Example 1.9.2 (Cubics). Here are some singular cubic surfaces $X \subset \mathbb{P}^3$, given by the vanishing of the corresponding cubic form:

$4A_1$	$x_0x_1x_2 + x_1x_2x_3 + x_2x_3x_0 + x_3x_0x_1$
$2A_1 + A_2$	$x_0x_1x_2 = x_3^2(x_1 + x_2 + x_3)$
$2A_1 + A_3$	$x_0x_1x_2 = x_3^2(x_1 + x_2)$
$A_1 + 2A_2$	$x_0x_1x_2 = x_1x_3^2 + x_3^3$
$A_1 + A_3$	$x_0x_1x_2 = (x_1 + x_2)(x_3^2 - x_1^2)$
$A_1 + A_4$	$x_0x_1x_2 = x_3^2x_2 + x_3x_1^2$
$A_1 + A_5$	$x_0x_1x_2 = x_1^3 - x_3^2x_2$
$3A_2$	$x_0x_1x_2 = x_3^3$
A_4	$x_0x_1x_2 = x_2^3 - x_3x_1^2 + x_3^2x_2$
A_5	$x_3^3 = x_1^3 + x_0x_3^2 - x_2^2x_3$
D_4	$x_1x_2x_3 = x_0(x_1 + x_2 + x_3)^2$
D_5	$x_0x_1^2 + x_1x_3^2 + x_2^2x_3$
E_6	$x_3^3 = x_1(x_1x_0 + x_2^2)$.

In practice, most geometric questions are easier for smooth surfaces, while most arithmetic questions turn out to be easier in the singular case. For a survey on arithmetic problems on rational surfaces, see Sections 2.4 and 3.4, as well as [MT86].

sect:K3

1.10. K3 surfaces. Let X be a smooth projective surface with trivial canonical class. There are two possibilities: X could be an abelian surface or a K3 surface. In the latter case, X is simply-connected and $h^1(X, \mathcal{O}_X) = 0$. The Picard group $\text{Pic}(X)$ of a K3 surface X is a torsion-free \mathbb{Z} -module of rank ≤ 20 and the intersection form on $\text{Pic}(X)$ is even, i.e., the square of every class is an even integer. K3 surfaces of with polarizations of small degree can be realized as complete intersections in projective space. The most common examples are K3 surfaces of degree 2, given explicitly as double covers $X \rightarrow \mathbb{P}^2$ ramified in a curve of degree 6; or quartic surfaces $X \subset \mathbb{P}^3$.

exam:rank-k3

Example 1.10.1. The Fermat quartic

$$x^4 + y^4 + z^4 + w^4 = 0$$

has Picard rank 20 over $\mathbb{Q}(\sqrt{-1})$. The surface X given by

$$xy^3 + yz^3 + zx^3 + w^4 = 0$$

has $\text{Pic}(X) = \mathbb{Z}^{20}$ over \mathbb{Q} (see [Ino78] for more examples). The surface $w(x^3 + y^3 + z^3 + x^2z + xw^2) = 3x^2y^2 + 4x^2yz + x^2z^2 + xy^2z + xyz^2 + y^2z^2$ has geometric Picard rank 1, i.e., $\text{Pic}(X_{\bar{\mathbb{Q}}}) = \mathbb{Z}$ [vL07].

Other interesting examples arise from abelian surfaces as follows:
Let

$$\begin{aligned} \iota : A &\rightarrow A \\ a &\mapsto -a \end{aligned}$$

be the standard involution. Its fixed points are the 2-torsion points of A . The quotient A/ι has 16 singularities (the images of the fixed points). The minimal resolution of these singularities is a K3 surface, called a *Kummer surface*. There are several other finite group actions on abelian surfaces such that a similar construction results in a K3 surface, a *generalized Kummer surface* (see [Kat87]).

The nef cone of a polarized K3 surface (X, g) admits the following characterization: h is ample if and only if $(h, C) > 0$ for each class C with $(g, C) > 0$ and $(C, C) \geq -2$. The *Torelli theorem* implies an intrinsic description of automorphisms: every automorphism of the lattice $\text{Pic}(X)$ preserving the nef cone arises from an automorphism of X . There is an extensive literature devoted to the classification of arising automorphism groups [Nik81], [Dol08]. These automorphisms give examples of interesting algebraic dynamical systems [McM02], [Can01]. they can be used to propagate rational points and curves [BT00], and to define canonical heights [Sil91], [Kaw08].

sect:3folds

1.11. Threefolds. The classification of smooth Fano threefolds was a major achievement completed in the works of Iskovskikh [Isk79], [IP99a], and Mori–Mukai [MM86]. There are more than 100 families. Among them, for example, cubics $X_3 \subset \mathbb{P}^4$, quartics $X_4 \subset \mathbb{P}^4$ or double covers of $W_2 \rightarrow \mathbb{P}^3$, ramified in a surface of degree 6. Many of these varieties, including the above examples, are not rational. Unirationality of cubics can be seen directly: projecting from a line on X_3 we get a cubic surface fibration, which *splits* after base change. The nonrationality of cubics was proved in [CG72] using *intermediate Jacobians*. Nonrationality of quartics was proved by establishing *birational rigidity*, i.e., showing triviality of the group of birational automorphisms, via an analysis of *maximal* singularities of such maps [IM71]. This technique has been substantially developed within the Minimal Model Program (see [Isk01], [Puk98], [Puk07], [Che05]) Some quartic threefolds are also unirational, e.g., the diagonal, Fermat type, quartic

$$\sum_{i=0}^4 x_i^4 = 0.$$

It is expected that the *general* quartic is *not* unirational. However, it admits an elliptic fibration: fix a line $\ell \in X_4 \subset \mathbb{P}^4$ and consider a plane in \mathbb{P}^4 containing this line, the residual plane curve has degree three and genus 1. A general double cover W_2 does not admit an elliptic or abelian fibration, even birationally [?].

sect:holo

1.12. Holomorphic symplectic varieties. Let X be a smooth projective simply-connected variety. It is called *holomorphic symplectic* if it carries a unique, modulo constants, nondegenerate holomorphic two-form. Typical examples are K3 surfaces X and their Hilbert schemes $X^{[n]}$ of zero-dimensional length- n subschemes. Another example is the variety of lines of a smooth cubic fourfold, it is deformation equivalent to $X^{[2]}$ of a K3 surface [?].

These varieties are interesting for the following reasons:

- the symplectic forms allows to define a *quadratic* form on $\text{Pic}(X)$, the Beauville–Bogomolov form;
- there is a *local* Torelli theorem;
- there is a *conjectural* characterization of the ample cone, at least in dimension 4 [?].

Further, there are examples with (Lagrangian) abelian fibrations over \mathbb{P}^n or with infinite *endomorphisms*, resp. *birational* automorphisms, which are interesting for arithmetic and algebraic dynamics.

sect:nonclosed

1.13. Geometry over nonclosed fields. Issues: descent, effective computation of invariants over the groundfield.

Example 1.13.1. The Picard group of cubic surfaces may be smaller over nonclosed fields: for X given by

$$x_0^3 + x_1^3 + x_2^3 + x_3^3 = 0$$

$\text{Pic}(X_{\mathbb{Q}}) = \mathbb{Z}^4$. It has a basis e_1, e_2, e_3, e_4 . such that $\Lambda_{\text{eff}}(X)$ is spanned by

$$\begin{aligned} &e_2, \quad e_3, \quad 3e_1 - 2e_3 - e_4, \quad 2e_1 - e_2 - e_3 - e_4, \quad e_1 - e_4, \\ &4e_1 - 2e_2 - 2e_3 - e_4, \quad e_1 - e_2, \quad 2e_1 - 2e_2 - e_4, \quad 2e_1 - e_3 \end{aligned}$$

(see [PT01]).

Computing Picard groups
[VAZ], [Zar08],
Effective

Example 1.13.2. Let X be a K3 surface over a number field \mathbb{Q} . Fix a model \mathcal{X} over \mathbb{Z} . For primes p of good reduction we have an injection

$$\mathrm{Pic}(X_{\bar{\mathbb{Q}}}) \hookrightarrow \mathrm{Pic}(X_{\bar{\mathbb{F}}_p}).$$

The rank ρ_p of $\mathrm{Pic}(X_{\bar{\mathbb{F}}_p})$ is always even. In some examples, ρ_p can be computed by counting points over \mathbb{F}_{p^r} , for several r , and by using the Weil conjectures.

This local information can sometimes be used to determine the rank $\rho_{\bar{\mathbb{Q}}}$ of $\mathrm{Pic}(X_{\bar{\mathbb{Q}}})$. Let p, q be distinct primes of good reduction such that $\rho_p, \rho_q \leq 2$ and the discriminants of the lattices $\mathrm{Pic}(X_{\bar{\mathbb{F}}_p}), \mathrm{Pic}(X_{\bar{\mathbb{F}}_q})$ do not differ by a square of a rational number. Then $\rho_{\bar{\mathbb{Q}}} = 1$.

2. EXISTENCE OF POINTS

sect:existence

2.1. Projective spaces and their forms. Let F be a field and \bar{F} an algebraic closure of F . A projective space over F has many rational points: they are dense in Zariski topology and in the adelic topology. Varieties F -birational to a projective space inherit these properties.

Over nonclosed fields F , projective spaces have *forms*, so called *Brauer–Severi* varieties. These are isomorphic to \mathbb{P}^n over \bar{F} but not necessarily over F . They can be classified via the nonabelian cohomology group $H^1(F, \mathrm{Aut}(\mathbb{P}^n))$, where $\mathrm{Aut}(\mathbb{P}^n) = \mathrm{PGL}_{n+1}$ is the group of algebraic automorphisms of \mathbb{P}^n . The basic example is a conic $C \subset \mathbb{P}^2$, e.g.,

eqn:legendre

$$(2.1) \quad ax^2 + by^2 = cz^2, \quad \text{with } a, b, c \in \mathbb{N}.$$

It is easy to verify solvability of this equation modulo p , for $p \nmid abc$ (cf. Theorem ??). Legendre proved that (2.1) has nontrivial solutions in \mathbb{Z} if and only if it has nontrivial solutions modulo p , for all primes p . This is an instance of a local-to-global principle that will be discussed in Section ??.

Checking solvability modulo p is a finite problem which gives a finite procedure to verify solvability in \mathbb{Z} . Actually, Legendre’s proof provides effective bounds for the size of the smallest solution, e.g.,

$$\max(|x|, |y|, |z|) \leq abc,$$

which gives another approach to checking solvability - try all $x, y, z \in \mathbb{N}$ subject to the inequality. If $C(\mathbb{Q}) \neq \emptyset$, then the conic is \mathbb{Q} -isomorphic to \mathbb{P}^1 : draw lines through a \mathbb{Q} -point in C .

In general, forms of \mathbb{P}^n over number fields satisfy the local-to-global principle. Moreover, Brauer–Severi varieties with at least one F -rational point are *split* over F , i.e., isomorphic to \mathbb{P}^n over F . It would be useful

to have a routine (in **Magma**) that would check efficiently whether or not a Brauer–Severi variety of small dimension over \mathbb{Q} , presented by explicit equations, is split, and to find the smallest solution.

sect:hyper

2.2. Hypersurfaces. Algebraically, the simplest examples of varieties are hypersurfaces, defined by a single homogeneous equation $f(\mathbf{x}) = 0$. Many classical diophantine problems reduce to the study of rational points on hypersurfaces.

Theorem 2.2.1 (Chevalley–Warning, Abh. M. Sem. Hamb. (1936)). *Let $X = X_f \subset \mathbb{P}^n$ be a hypersurface over a finite field F given by the equation $f(\mathbf{x}) = 0$. If $\deg(f) \leq n$ then $X(F) \neq \emptyset$.*

Proof. We reproduce a textbook argument [BS66], for $F = \mathbb{F}_p$.

Step 1. Consider the δ -function

$$\sum_{x=1}^{p-1} x^d = \begin{cases} -1 & \text{mod } p \quad \text{if } p-1 \mid d \\ 0 & \text{mod } p \quad \text{if } p-1 \nmid d \end{cases}$$

Step 2. Apply it to a (not necessarily homogeneous) polynomial $\phi \in \mathbb{F}_p[x_0, \dots, x_n]$, with $\deg(\phi) \leq n(p-1)$. Then

$$\sum_{x_0, \dots, x_n} \phi(x_0, \dots, x_n) = 0 \pmod{p}.$$

Indeed, for monomials, we have

$$\sum_{x_0, \dots, x_n} x_1^{d_1} \cdots x_n^{d_n} = \prod (\sum x_j^{d_j}), \quad \text{with } d_0 + \dots + d_n \leq n(p-1).$$

For some j , we have $0 \leq d_j < p-1$.

Step 3. For $\phi(x) = 1 - f(x)^{p-1}$ we have $\deg(\phi) \leq \deg(f) \cdot (p-1)$. Then

$$\mathbf{N}(f) := \#\{x \mid f(x) = 0\} = \sum_{x_0, \dots, x_n} \phi(x) = 0 \pmod{p},$$

since $\deg(f) \leq n$.

Step 4. The equation $f(\mathbf{x}) = 0$ has a trivial solution. It follows that

$$\mathbf{N}(f) > 1 \quad \text{and} \quad X_f(\mathbb{F}_p) \neq \emptyset.$$

□

Jacobi sums ...

Theorem 2.2.2. [Esn03] *If X is a Fano variety over a finite field \mathbb{F}_q then*

$$X(\mathbb{F}_q) \neq \emptyset.$$

Now we pass to the case in which $F = \mathbb{Q}$. Given a form $f \in \mathbb{Z}[x_0, \dots, x_n]$, homogeneous of degree d , we ask how many solutions $\mathbf{x} = (x_0, \dots, x_n) \in \mathbb{Z}^{n+1}$ to the equation $f(\mathbf{x}) = 0$ should we expect? Primitive solutions with $\gcd(x_0, \dots, x_n) = 1$, up to diagonal multiplication with ± 1 , are in bijection to rational points on the hypersurface $X_f \subset \mathbb{P}^n$. We have $|f(\mathbf{x})| = O(B^d)$, for $\|\mathbf{x}\| := \max_j(|x_j|) \leq B$. We may argue that f takes values $0, 1, 2, \dots$ with equal probability, so that the the probability of $f(\mathbf{x}) = 0$ is B^{-d} . There are B^{n+1} “events” with $\|\mathbf{x}\| \leq B$. In conclusion, we expect B^{n+1-d} solutions with $\|\mathbf{x}\| \leq B$. There are three cases:

- $n+1 < d$: as $B \rightarrow \infty$ we should have *fewer and fewer* solutions, and, eventually, none!
- $n+1 = d$: this is the *stable* regime, we get no information in the limit $B \rightarrow \infty$;
- $n+1 > d$: the expected number of solutions grows.

We will see many instances when this heuristic reasoning fails. However, it is reasonable, as a first approximation, at least when $n+1-d \gg 0$.

Diagonal hypersurfaces have attracted the attention of computational number theorists (see <http://euler.free.fr>). A sample is given below:

exam:many

Example 2.2.3.

- There are no rational points (with non-zero coordinates) on the Fano 5-fold $x_0^6 = \sum_{j=1}^6 x_j^6$ with height $\leq 2.6 \cdot 10^6$.
- There are 12 (up to signs and permutations) rational points on $x_0^6 = \sum_{j=1}^7 x_j^6$ of height $\leq 10^5$ (with non-zero coordinates).
- The number of rational points (up to signs, permutations and with non-zero coordinates) on the Fano 5-fold $x_0^6 + x_1^6 = \sum_{j=2}^6 x_j^6$ of height $\leq 10^4$ (resp. $2 \cdot 10^4, 3 \cdot 10^4$) is 12 (resp. 33, 57).

It is visible, that it is difficult to generate solutions when the $n-d$ is small. On the other hand, we have a the following theorem:

thm:birch

Theorem 2.2.4. [Bir62] *If $n \geq (\deg(f) - 1) \cdot 2^{\deg(f)}$, and f is smooth, then X_f satisfies the Hasse principle and the number $\mathbf{N}(f, B)$ of solutions $x = (x_i)$ with $\max(|x_i|) \leq B$ is*

$$\mathbf{N}(f, B) \sim \prod_p \tau_p \cdot \tau_\infty B^{n+1-d}, \quad \text{as } B \rightarrow \infty,$$

where τ_p, τ_∞ are the p -adic, resp. real, densities.

Further developments:

- asymptotic formulas and weak approximation
- better bounds for n for small $\deg(f)$
- works over $\mathbb{F}_q[t]$

Now we assume that $X = X_f$ is a hypersurface over a function field in one variable $F = \mathbb{C}(t)$. We have

Theorem 2.2.5. *If $\deg(f) \leq n$ then $X_f(\mathbb{C}(t)) \neq \emptyset$.*

Proof. It suffices to count parameters: Insert $x_j = x_j(t) \in \mathbb{C}[t]$, of degree e , into

$$f = \sum_J f_J x^J = 0,$$

with $|J| = \deg(f)$. This gives a system of $e \cdot \deg(f) + \text{const}$ equations in $e(n + 1)$ variables. This system is solvable for $e \gg 0$, provided $\deg(f) \leq n$. □

sect:decide

2.3. Decidability. Hilbert's 10th problem.

Matiyasevich (1970), Matiyasevich-Robinson (1975)

Theorem 2.3.1. *Let $f \in \mathbb{Z}[t, z_1, \dots, z_n]$ be polynomial. The set of $t \in \mathbb{Z}$ such that $f(t, \dots, z_n) = 0$ is solvable in \mathbb{Z} is not decidable, i.e., there is no algorithm to decide whether or not a diophantine equation is solvable in integers.*

Theorem 2.3.2. [?] *The set of $t \in \mathbb{Z}$ such that $f_t = 0$ has infinitely many primitive solutions is algorithmically random ¹.*

¹The author's abstract: "One normally thinks that everything that is true is true for a reason. I've found mathematical truths that are true for no reason at all. These mathematical truths are beyond the power of mathematical reasoning because they are accidental and random. Using software written in Mathematica that runs on an IBM RS/6000 workstation, I constructed a perverse 200-page algebraic equation with a parameter t and 17,000 unknowns. For each whole-number value of the parameter t , we ask whether this equation has a finite or an infinite number of whole number solutions. The answers escape the power of mathematical reason because they are completely random and accidental."

Decidability over other rings

- Yes: $\mathbb{Z}[S^{-1}]$, for some sets S of primes, of density 1 (Poonen/Shlapentokh 2003)
- Yes: $\mathbb{F}_p, \mathbb{Q}_p, \mathbb{R}, \mathbb{C}$
- Unknown: \mathcal{O}_K , number fields K (including \mathbb{Q})
- Unknown: $\mathbb{C}(t)$
- No: $\mathbb{F}_q[t], \mathbb{C}(t_1, \dots, t_n)$ with $n \geq 2$

sect:obstruct

2.4. Obstructions. As we have seen in Section 2.3, there is no hope of finding an algorithm which would determine the solvability of a diophantine equation in integers, i.e., there is no algorithm to test for the existence of *integral* points on quasi-projective varieties. The corresponding question for *homogeneous* equations, i.e., for *rational* points, is still open. It is reasonable to expect that at least for some classes of algebraic varieties, for example, for Del Pezzo surfaces, the existence question can be answered. In this section we survey some recent results in this direction.

Let X_B be a scheme over a base scheme B . We are looking for obstructions to the existence of points $X(B)$, i.e., sections of the structure morphism $X \rightarrow B$. Each morphism $B' \rightarrow B$ gives rise to a base-change diagram, and each section $x : B \rightarrow X$ provides a section $x' : B' \rightarrow X_{B'}$.

$$\begin{array}{ccc} X & \longleftarrow & X_{B'} \\ \downarrow & & \downarrow \\ B & \longleftarrow & B' \end{array} \qquad \begin{array}{ccc} X & \longleftarrow & X_{B'} \\ \uparrow x & & \uparrow x' \\ B & \longleftarrow & B' \end{array}$$

This gives rise to a *local* obstruction, since it is sometimes easier to check that $X_{B'}(B') = \emptyset$. In practice, B could be a curve and B' a cover, or an analytic neighborhood of a point on B . In the number-theoretic context, $B = \text{Spec}(F)$ and $B' = \text{Spec}(F_v)$, where v is a valuation of the number field F and F_v the v -adic completion of F . One says that the *local-global principle*, or the *Hasse principle*, holds, if the existence of F -rational points is implied by the existence of v -adic points in all completions.

exam:hasse

Example 2.4.1. The Hasse principle holds for:

- (1) smooth quadrics $X_2 \subset \mathbb{P}^n$;
- (2) Brauer–Severi varieties;
- (3) Del Pezzo surfaces of degree ≥ 5 ;

- (4) Chatelet surfaces $y^2 - az^2 = f(x_0, \dots, x_n)$, where f is an irreducible polynomial of degree ≤ 4 [CTSSD87b];
- (5) hypersurfaces $X_d \subset \mathbb{P}^n$, for $n \gg d$ (see Theorem 4.8.1).

The Hasse principle may fail for cubic curves, e.g.,

$$3x^3 + 4y^3 + 5z^3 = 0.$$

In topology, there is a rich obstruction theory to the existence of sections. An adaptation to algebraic geometry is formulated as follows: Let \mathfrak{C} be a contravariant functor from the category of schemes over a base scheme B to the category of abelian groups. Applying the functor \mathfrak{C} to the diagrams above, we have

$$\begin{array}{ccc} \mathfrak{C}(X) & \longrightarrow & \mathfrak{C}(X_{B'}) \\ \uparrow & & \uparrow \\ \mathfrak{C}(B) & \longrightarrow & \mathfrak{C}(B') \end{array} \qquad \begin{array}{ccc} \mathfrak{C}(X) & \longrightarrow & \mathfrak{C}(X_{B'}) \\ \downarrow x & & \downarrow x' \\ \mathfrak{C}(B) & \longrightarrow & \mathfrak{C}(B') \end{array}$$

If for all sections x' , the image of x' in $\mathfrak{C}(B')$ is nontrivial in the cokernel of the map $\mathfrak{C}(B) \rightarrow \mathfrak{C}(B')$, then we have a problem, i.e., an obstruction to the existence of B -points on X . So far, this is still a version of a local obstruction. However, a *global obstruction* may arise, when we vary B' .

We are interested in the case when $B = \text{Spec}(F)$, for a number field F , with B' ranging over all completions F_v . A global obstruction is possible whenever the map

$$\mathfrak{C}(\text{Spec}(F)) \rightarrow \prod_v \mathfrak{C}(\text{Spec}(F_v))$$

has a nontrivial cokernel. What are sensible choices for \mathfrak{C} ? Basic contravariant functors on schemes are $\mathfrak{C}(-) := H_{et}^i(-, \mathbb{G}_m)$. For $i = 1$, we get the Picard functor, introduced in Section 1.1. However, by Hilbert's theorem 90, $H_{et}^1(\text{Spec}(F), \mathbb{G}_m) = 0$, for all fields F , and this won't generate an obstruction. For $i = 2$, we get the (cohomological) Brauer group $\text{Br}(X) = H_{et}^2(X, \mathbb{G}_m)$, classifying sheaves of central simple algebras over X , modulo equivalence (see [?, Chapter 4]). By class field theory, we have an exact sequence

ean:brauer (2.2) $0 \rightarrow \text{Br}(F) \rightarrow \bigoplus_v \text{Br}(F_v) \xrightarrow{\sum_v \text{inv}_v} \mathbb{Q}/\mathbb{Z} \rightarrow 0,$

where $\text{inv}_v : \text{Br}(F_v) \rightarrow \mathbb{Q}/\mathbb{Z}$ is the *local invariant*. We apply it to the diagram and obtain:

$$\begin{array}{ccccccc}
\mathrm{Br}(X_F) & \longrightarrow & \bigoplus_v \mathrm{Br}(X_{F_v}) & & & & \\
\downarrow x & & \downarrow (x_v)_v & & & & \\
0 \longrightarrow & \mathrm{Br}(F) & \longrightarrow & \bigoplus_v \mathrm{Br}(F_v) & \xrightarrow{\sum_v \mathrm{inv}_v} & \mathbb{Q}/\mathbb{Z} & \longrightarrow 0,
\end{array}$$

Define

eqn:hasse-prin

$$(2.3) \quad X(\mathbb{A}_F)^{\mathrm{Br}} := \bigcap_{A \in \mathrm{Br}(X)} \{(x_v)_v \in X(\mathbb{A}_F) \mid \sum_v \mathrm{inv}(A(x_v)) = 0\}.$$

Let $\overline{X(F)}$ be the closure of $X(F)$ in $X(\mathbb{A}_F)$, in the adelic topology. One says that X satisfies *weak approximation* over F if $\overline{X(F)} = X(\mathbb{A}_F)$. We have

$$X(F) \subset \overline{X(F)} \subset X(\mathbb{A}_F)^{\mathrm{Br}} \subset X(\mathbb{A}_F).$$

From this we derive the *Brauer–Manin* obstruction to the Hasse principle and weak approximation:

- if $X(\mathbb{A}_F) \neq \emptyset$ but $X(\mathbb{A}_F)^{\mathrm{Br}} = \emptyset$ then $X(F) = \emptyset$, i.e., the Hasse principle fails;
- if $X(\mathbb{A}_F) \neq X(\mathbb{A}_F)^{\mathrm{Br}}$ then weak approximation fails.

Del Pezzo surfaces of degree ≥ 5 satisfy the Hasse principle and weak approximation. Arithmetically most interesting are Del Pezzo surfaces of degree 4, 3, and 2: these may fail the Hasse principle:

- deg = 4: $z^2 + w^2 = (x^2 - 2y^2)(3y^2 - x^2)$ [Isk71];
- deg = 3: $5x^3 + 12y^3 + 9z^3 + 10w^3 = 0$ [CG66];
- deg = 2: $w^2 = 2x^4 - 3y^4 - 6z^4$ [KT04].

One says that the Brauer–Manin obstruction to the existence of rational point is the only one, if $X(\mathbb{A}_F)^{\mathrm{Br}} \neq \emptyset$ implies that $X(F) \neq \emptyset$. This holds for:

- (1) abelian varieties over a number field F with *finite* Tate–Shafarevich group [Man71];
- (2) principal homogeneous spaces for a connected linear algebraic group;
- (3) Del Pezzo surfaces of degree ≥ 3 admitting a conic bundle structure defined over the groundfield F .
- (4) conjecturally(!), for all geometrically rational surfaces.

descent, torsor formalism

Specialization: $x \in X(F) \mapsto [\mathcal{T}_x] \in H_{\mathrm{et}}^1(F, T)$, a finite set. This gives a partition:

$$X(F) = \bigcup_{\alpha \in H_{\mathrm{et}}^1(F, T)} \pi_\alpha(\mathcal{T}_\alpha(F)),$$

where \mathcal{T}_α are the *descent* varieties.

Transcendental obstructions: [Har96]

sect:descent

2.5. **Descent.** Abelian descent

Nonabelian descent [Har08], [Sko99], [HS05], [HS02]

Bjorn’s example [Poo08]

For detailed surveys of the theory of the Brauer-Manin obstruction are [Pey05], [Har04].

sect:effect

2.6. **Effectivity.** In light of the discussion in Section 2.3 it is important to know whether or not the Brauer–Manin obstruction can be computed, effectively in terms of the coefficients of the defining equations. There is an extensive literature on computations of the Brauer–Manin obstruction for curves [].

Effective computability of the Brauer-Manin obstruction for all Del Pezzo surfaces over number fields has been proved in [KT08]. The main steps are as follows:

- (1) Computation of the equations of the exceptional curves and of the action of the Galois group of the splitting field G on these curves as in Section 1.13. One obtains the exact sequence of G -modules

$$0 \rightarrow \text{Relations} \rightarrow \oplus \mathbb{Z}E_j \rightarrow \text{Pic}(\overline{X}) \rightarrow 0.$$

- (2) We have

$$\text{Br}(X)/\text{Br}(F) = H^1(G, \text{Pic}(\overline{X})).$$

Using the equations for exceptional curves and functions realizing relations between the curves classes in the Picard group one can compute explicitly Azumaya algebras $\{\mathcal{A}_i\}$ representing the classes of $\text{Br}(X)/\text{Br}(F)$.

- (3) The local points $S(F_v)$ can be effectively decomposed into a *finite* union of subsets such that each \mathcal{A}_i is constant on each of these subsets. This step uses an effective version of the arithmetic Hilbert Nullstellensatz.
- (4) It remains to compute the local invariants.

2.7.

Example 2.7.1. [HBM04] Let $a, b \in \mathbb{Z}$ be coprime such that $a = \pm b \pmod{9}$. Then the cubic surface

$$x_0^3 + 2x_1^3 + ax_2^3 + bx_3^3 = 0$$

has a \mathbb{Q} -rational point.

The proof combines a deep result in the theory of elliptic curves [?], namely, the existence of nontrivial Heegner points on the curve $x_0^3 + 2x_1^3 = py^3$, for any prime $p = 2 \pmod{9}$, with a result in analytic number theory on representations of primes p by the cubic form $ax_2^3 + bx_3^3$.

3. DENSITY OF POINTS

sect:denlang

3.1. Lang’s conjecture. One of the main principles underlying arithmetic geometry is the expectation that the trichotomy in the classification of algebraic varieties via the Kodaira dimension in Section 1.2 has an arithmetic manifestation. The broadly accepted form of this is

conj:lang

Conjecture 3.1.1 (Lang’s conjecture). Let X be a variety of general type, i.e., a smooth projective variety with ample canonical class, defined over a number field F . Then $X(F)$ is not Zariski dense.

What about a converse? The obvious necessary condition for Zariski density of rational points, granted Conjecture ??, is that X does not dominate a variety of general type. This condition is not enough, as was shown in [?]: there exist surfaces which do not dominate curves of general type but which have étale covers dominating curves of general type. By the Chevalley–Weil theorem (see Section ??), these covers would have a dense set of rational points, over some finite extension of the ground field, contradicting Conjecture ??.

The formulation of a converse statement

For a detailed discussion of the geometric considerations related to potential density and Campana’s program see [Abr08].

sect:dense-fixed

3.2. Zariski density over fixed fields. Here we discuss Zariski density of rational points in the “unstable” situation, when the density of points is governed by subtle number-theoretical properties, rather than geometric considerations. We have the following fundamental result:

thm:pot-curve

Theorem 3.2.1. *Let C be a smooth curve of genus $g = g(C)$ over a number field F . Then*

- if $g = 0$ and $C(F) \neq \emptyset$ then $C(F)$ is Zariski dense;
- if $g = 1$ and $C(F) \neq \emptyset$ then $C(F)$ is an abelian group (the Mordell–Weil group) and there is a constant c_F (independent of C) bounding the order of the torsion subgroup $C(F)_{\text{tors}}$ of $C(F)$ [Maz77], [Mer96]; in particular, if there is an F -rational point of infinite order then $C(F)$ is Zariski dense;
- if $g \geq 2$ then $C(F)$ is finite [Fal83], [Fal91].

In fact, assuming the Lang-Vojta conjecture, one obtains *uniform* bounds for $\#C(F)$, if $g(C) \geq 2$ [CHM97b], [CHM97a]. EXPAND...

In higher dimensions we have:

thm:pot-de

Theorem 3.2.2. *Let X be an algebraic variety over a number field F . Assume that $X(F) \neq \emptyset$ and that X is one of the following*

- X is a Del Pezzo surface of degree 2 and has a point on the complement to exceptional curves;
- X is a Del Pezzo surface of degree ≥ 3 ;
- X is a Brauer-Severi variety.

Then $X(F)$ is Zariski dense.

The proof of the first claims can be found in [Man86, p. ??].

rema:de1

Remark 3.2.3. Let X/F be a Del Pezzo surface of degree 1 (it always contains an F -rational point, the base point of the anticanonical linear series) or a conic bundle $X \rightarrow \mathbb{P}^1$, with $X(F) \neq \emptyset$. It is unknown whether or not $X(F)$ is Zariski dense.

Theorem 3.2.4. [Elk88] *Let $X \subset \mathbb{P}^3$ be the quartic K3 surface given by*

eqn:xx

$$(3.1) \quad x_0^4 + x_1^4 + x_2^4 = x_3^4.$$

Then $X(\mathbb{Q})$ is Zariski dense.

The trivial solutions $(1 : 0 : 0 : 1)$ etc are easily seen. The smallest nontrivial solution is ...

Example 3.2.5. [EJ06] Let $X \subset \mathbb{P}^3$ be the quartic given by

$$x^4 + 2y^4 = z^4 + 4t^4.$$

The obvious \mathbb{Q} -rational points are given by $y = t = 0$ and $x = \pm z$. The next smallest solution is

$$1484801^4 + 2 \cdot 1203120^4 = 1169407^4 + 4 \cdot 1157520^4.$$

sect:potential

3.3. Potential density: techniques. Here is a (short) list of possible strategies to propagate points:

- use the group of automorphisms $\text{Aut}(X)$, if it is infinite;
- try to find a dominant map $\tilde{X} \rightarrow X$ where \tilde{X} satisfies potential density (for example, try to prove *unirationality*);
- try to find a fibration structure $X \rightarrow B$ where the fibers satisfy potential density in some *uniform way* (that is, the field extensions needed to insure potential density of the fibers V_b can be uniformly controlled).

In particular, it is important for us keep track of minimal conditions which would insure Zariski density of points on varieties. A fundamental result is

exam:conic-bundles

Example 3.3.1. Let $\pi : X \rightarrow \mathbb{P}^1$ be a conic bundle, defined over a field F . Then rational points on X are potentially dense. Indeed, by Tsen's theorem, π has section $s : \mathbb{P}^1 \rightarrow X$ (which is defined over some finite extension F'/F), each fiber has an F' -rational point and it suffices to apply Theorem 3.2.1. Potential density for conic bundles over higher-dimensional bases is an open problem.

If X is an abelian variety then there exists a finite extension F'/F and a point $P \in X(F')$ such that the cyclic subgroup of $X(F')$ generated by P is Zariski dense.

Example 3.3.2. If $\pi : X \rightarrow \mathbb{P}^1$ is a Jacobian nonisotrivial elliptic fibrations (π admits a section and the j -invariant is nonconstant), then potential density follows from a strong form of the Birch/Swinnerton-Dyer conjecture [GM97], [Man95]. The key problem is to control the variation of the *root number* (the sign of the functional equation of the L -functions of the elliptic curve) (see [GM97]).

On the other hand, rational points on the elliptic fibration are not potentially dense [CTSSD97].

exam:geome-ell

Example 3.3.3. One geometric approach to Zariski density of rational points on (certain) elliptic fibration can be summarized as follows:

Case 1. Let $\pi : X \rightarrow \mathbb{P}^1$ be a Jacobian elliptic fibration and $e : \mathbb{P}^1 \rightarrow X$ its zero-section. Suppose that we have another section s which is nontorsion in the Mordell-Weil group of $X(F(B))$. Then a specialization argument implies that the restriction of the section to infinitely many fibers of π gives a nontorsion point in the Mordell-Weil group of the corresponding fiber (see [Ser90], 11.1). In particular, $X(F)$ is Zariski dense.

Case 2. Suppose that $\pi : X \rightarrow \mathbb{P}^1$ is an elliptic fibration with a *multisection* M (an irreducible curve surjecting onto the base \mathbb{P}^1). After a basechange $X \times_{\mathbb{P}^1} M \rightarrow M$ the elliptic fibration acquires the identity section Id (the image of the diagonal under $M \times_{\mathbb{P}^1} M \rightarrow V \times_{\mathbb{P}^1} M$) and a (rational) section

$$\tau_M := d\text{Id} - \text{Tr}(M \times_{\mathbb{P}^1} M),$$

where d is the degree of $\pi : M \rightarrow \mathbb{P}^1$ and $\text{Tr}(M \times_{\mathbb{P}^1} M)$ is obtained (over the generic point) by summing all the points of $M \times_{\mathbb{P}^1} M$. We will say that M is *nontorsion* if τ_M is nontorsion.

If M is nontorsion and if $M(F)$ is Zariski dense then the same holds for $X(F)$ (see [BT99]).

sect:pot-dense-res

3.4. Potential density for surfaces.

exem:brs-fibr

Example 3.4.1. Let $\pi : X \rightarrow C$ be a fibration with generic fiber a Brauer-Severi variety or a Del Pezzo surface of degree ≥ 2 . Theorem ?? implies that π has a section. Potential density for X follows from Theorem 3.2.2.

rema:ab-fibr

Remark 3.4.2. An argument similar to Example ?? works for abelian fibrations. The difficulty here is to formulate some simple geometric conditions insuring that a (multi)section leads to points which are not only of infinite order in the Mordell-Weil groups of the corresponding fibers, but in fact generate Zariski dense subgroups.

By Theorem 3.2.1, potential density holds for curves of genus $g \leq 1$. It holds for surfaces which become rational after a finite extension of the ground field, e.g., for all Del Pezzo surfaces. The classification theory in dimension 2 gives us the following list of surfaces of intermediate type:

- abelian surfaces;
- bielliptic surfaces;
- Enriques surfaces;
- K3 surfaces.

Potential density for the first two classes follows from Theorem 3.2.2. The classification of Enriques surfaces X implies that either $\text{Aut}(X)$ is infinite or X is dominated by a K3 surface \tilde{X} with $\text{Aut}(\tilde{X})$ infinite [?]. Thus we are reduced to the study of K3 surfaces.

Theorem 3.4.3. [BT00] *Let X be a K3 surface over any field of characteristic zero. If X is elliptic or admits an infinite group of automorphisms then rational points on X are potentially dense.*

Sketch of the proof. The main difficulty is to find sufficiently nondegenerate rational or elliptic multisections of the elliptic fibration $X \rightarrow \mathbb{P}^1$. These are produced using deformation theory. One starts with special K3 surfaces which have rational curves $C_t \subset X_t$ in the desired homology class (for example, Kummer surfaces) and then deforms the pair. Actually, this deformation technique has to be applied to twists of the original elliptic surface. \square

rema:genk3

Remark 3.4.4. General K3 surfaces X have (geometric) Picard number one. In particular, $\text{Aut}(X)$ is finite and there are no elliptic fibrations. Potential density of rational points is an open problem.

Example 3.4.5. A smooth hypersurface $X \subset \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ of bi-degree $(2, 2, 2)$ is a K3 surface with $\text{Aut}(X)$ infinite.

exem:k33

Example 3.4.6. Every smooth quartic surface $S_4 \subset \mathbb{P}^3$ which contains a line is an elliptic K3 surface. Indeed, let M be this line and assume that both S_4 and M are defined over a number field F . Consider the 1-parameter family of planes $\mathbb{P}_t^2 \subset \mathbb{P}^3$ containing M . The residual curve in the intersection $\mathbb{P}_t^2 \cap S_4$ is a plane cubic intersecting M in 3 points. This gives a fibration $\pi : S_4 \rightarrow \mathbb{P}^1$ with a *rational* tri-section M .

To apply the strategy of Section 6.1 we need to insure that M is nontorsion. A sufficient condition, satisfied for generic quartics S_4 , is that the restriction of π to M ramifies in a smooth fiber of $\pi : X \rightarrow \mathbb{P}^1$. Under this condition $X(F)$ is Zariski dense.

LATER

We now collect some results and open problems concerning the arithmetic of K3 surfaces.

Existence:

Swinerton-Dyer/Slater

Zariski density

Theorem 3.4.7. [Elk88] *Let $X \subset \mathbb{P}^3$ be the quartic K3 surface given by*

eqn:xx (3.2)
$$x_0^4 + x_1^4 + x_2^4 = x_3^4.$$

Then $X(\mathbb{Q})$ is Zariski dense.

The trivial solutions $(1 : 0 : 0 : 1)$ etc are easily seen. The smallest nontrivial solution is ... Geometrically, over $\bar{\mathbb{Q}}$, the surface given by (3.2) is a Kummer surface, with many elliptic fibrations.

Theorem 3.4.8. [HT00] *Let $X \subset \mathbb{P}^3$ be a quartic K3 surface containing a line defined over a field F . If X is general, then $X(F)$ is Zariski dense. In all cases, there exists a finite extension F'/F such that $X(F')$ is Zariski dense.*

Theorem 3.4.9. [?] *Let X be any elliptic K3 surface over any field F . Then rational points are potentially dense.*

sect:pot3

3.5. Potential density in dimension ≥ 3 . Potential density holds for unirational varieties. Classification of (smooth) Fano threefolds and the detailed study of occurring families implies unirationality for all but three cases:

- X_4 : quartics in \mathbb{P}^4 ;
- V_1 : double covers of a cone over the Veronese surface in \mathbb{P}^5 famified in a surface of degree 6;
- W_2 : double covers of \mathbb{P}^3 ramified in a surface of degree 6.

We now sketch the proof of potential density for quartics from [HT00], the case of V_1 is treated by similar techniques in [BT99].

The threefold X_4 contains a 1-parameter family of lines. Choose a line M (defined over some extension of the groundfield, if necessary) and consider the 1-parameter family of hyperplanes $\mathbb{P}_t^3 \subset \mathbb{P}^4$ containing M . The generic hyperplane section $S_t := \mathbb{P}_t^3 \cap X_4$ is a quartic surface with a line. Now we would like to argue as in the Example 3.4.6. We need to make sure that M is nontorsion in S_t for a dense set of $t \in \mathbb{P}^1$. This will be the case for general X_4 and M . The analysis of all exceptional cases requires care.

Remark 3.5.1. It would be interesting to have see (nontrivial) examples of Calabi-Yau threefolds with a dense set of rational points.

Theorem 3.5.2. [?] *Let X be a K3 surface over a field F , of degree $2d$. Then rational points on $X^{[n]}$ are potentially dense.*

The proof relies on the existence of an abelian fibration

$$Y := X^{[n]} \rightarrow \mathbb{P}^n,$$

with a nontorsion multisection which has a potentially dense set of rational points. Numerically, such fibrations are predicted by square-zero classes in the Picard group $\text{Pic}(Y)$, with respect to the Beauville–Bogomolov form (see Section 1.12). Geometrically, the fibration is the degree n Jacobian fibration associated to hyperplane sections of X .

Theorem 3.5.3. [AV07] *Let Y be the Fano variety of lines on a general cubic fourfold $X_3 \subset \mathbb{P}^5$ over a field of characteristic zero. Then rational points on Y are potentially dense.*

Sketch of proof. The key tool is a rational endomorphism $\phi: Y \rightarrow Y$ analyzed in [?]: let ℓ on $X_3 \subset \mathbb{P}^5$ be a general line and $\mathbb{P}_\ell^2 \subset \mathbb{P}^5$ the unique plane everywhere tangent to ℓ . Let $[\ell] \in Y$ be the corresponding point and put $\phi([\ell]) := [\ell']$, where ℓ' is the residual line in X_3 .

Generically, one can expect that the orbit $\{\phi^n([\ell])\}_{n \in \mathbb{N}}$ is Zariski dense in Y . This was proved by Amerik and Campana in [AC08], over *uncountable* ground fields. Over countable fields, one faces the difficulty that the countably many exceptional loci could cover all algebraic points of Y . Amerik and Voisin were able to overcome this obstacle over number fields. Rather than proving density of $\{\phi^n([\ell])\}_{n \in \mathbb{N}}$ they find surfaces $\Sigma \subset Y$, birational to abelian surfaces, whose orbits are dense in Y . The main effort goes into showing that one can choose sufficiently general Σ defined over $\bar{\mathbb{Q}}$, provided that Y is sufficiently general and still defined over a number field. In particular, Y has geometric Picard number one. A case by case geometric analysis excludes the possibility that the Zariski closure of $\{\phi^n(\Sigma)\}_{n \in \mathbb{N}}$ is a proper subvariety of F . \square

Theorem 3.5.4. [HT] *Let Y be the variety of lines of a cubic fourfold $X_3 \subset \mathbb{P}^5$ which contains a cubic scroll T . Assume that the hyperplane section of X_3 containing T has exactly 6 double points in linear general position and that X_3 does not contain a plane. If X_3 and T are defined over a field F then $Y(F)$ is Zariski dense.*

rema:uni-para

Remark 3.5.5. In higher dimensions, (smooth) hypersurfaces $X_d \subset \mathbb{P}^n$ of degree d represent a major challenge. The circle method works well when

$$n \gg 2^d$$

while the geometric methods for proving unirationality require at least a super-exponential growth of n (see [?] for a construction of a unirational parametrization).

3.6. Density in analytic topologies. Automorphisms $\phi : X \rightarrow X$ give rise to interesting dynamical systems. Again, K3 surfaces serve as basic examples. The forward orbits $\{\phi^n(x)\}_{n \in \mathbb{N}}$ of real points can generate beautiful pictures in $X(\mathbb{R})$ (see [McM02]).

3.7. Approximation.

Example 3.7.1. [Wit04] Let $E \rightarrow \mathbb{P}^1$ be the elliptic fibration given by $y^2 = x(x - g)(x - h)$ where $g(t) = 3(t - 1)^3(t + 3)$ and $h = g(-t)$. Its minimal proper regular model X is an elliptic K3 surface that fails weak approximation. The obstruction comes from transcendental classes in the Brauer group of X .

4. COUNTING PROBLEMS

sect:counting

Here we consider projective algebraic varieties $X \subset \mathbb{P}^n$ defined over a number field F . We assume that $X(F)$ is Zariski dense. We seek to understand the distribution of rational points with respect to heights.

sect:heights

4.1. Heights. First we assume that $F = \mathbb{Q}$. Then we can define a *height* integral (respectively rational points) on the affine (respectively projective space) as follows

$$\begin{aligned} H_{\text{affine}} : \mathbb{A}^n(\mathbb{Z}) = \mathbb{Z}^n &\rightarrow \mathbb{R}_{\geq 0} \\ x = (x_1, \dots, x_n) &\mapsto \max_j(|x_j|) \\ H : \mathbb{P}^n(\mathbb{Q}) = (\mathbb{Z}_{\text{prim}}^{n+1} \setminus 0)/\pm &\rightarrow \mathbb{R}_{> 0} \\ x = (x_0, \dots, x_n) &\mapsto \max_j(|x_j|). \end{aligned}$$

This induces heights on points of subvarieties of affine or projective spaces. In some problems it is useful to work with equivalent norms, e.g., $\sqrt{\sum x_j^2}$ instead of $\max_j(|x_j|)$. Such choices are referred to as a *change of metrization*. A more conceptual definition of heights and adelic metrizations is given in Section ??

Heights on products, reembeddings, etc.

4.2. Counting functions. For a subvariety $X \subset \mathbb{P}^n$ put

$$N(X, B) := \#\{x \in X(\mathbb{Q}) \mid H(x) \leq B\}.$$

What can be said about

$$N(X, B), \text{ for } B \rightarrow \infty ?$$

Main questions here concern:

- (uniform) upper bounds,

- asymptotic formulas,
- geometric interpretation of the asymptotics.

By the very definition, $\mathbf{N}(X, B)$ depends on the projective embedding of X . For $X = \mathbb{P}^n$ over \mathbb{Q} , with the standard embedding via the line bundle $\mathcal{O}(1)$, we get

$$N(\mathbb{P}^n, B) \sim \frac{1}{\zeta(n+1)} \cdot \tau_\infty \cdot B^{n+1}, \quad B \rightarrow \infty.$$

But we may also consider the Veronese re-embedding

$$\begin{aligned} \mathbb{P}^n &\rightarrow \mathbb{P}^N \\ x &\mapsto x^I, \quad |I| = d, \end{aligned}$$

e.g.,

$$\begin{aligned} \mathbb{P}^1 &\rightarrow \mathbb{P}^2 \\ (x_0 : x_1) &\mapsto (x_0^2 : x_0x_1 : x_1^2) \end{aligned}$$

The image $y_0y_2 = y_1^2$ has $\sim B$ points of height $\leq B$. Similarly, the number of rational points on height $\leq B$ in the $\mathcal{O}(d)$ embedding of \mathbb{P}^n will be $\sim B^{(n+1)/d}$.

More generally, if F/\mathbb{Q} is a finite extension, put

$$\begin{aligned} \mathbb{P}^n(F) &\rightarrow \mathbb{R}_{>0} \\ x &\mapsto \prod_v \max(|x_j|_v). \end{aligned}$$

`thm:schanuel`

Theorem 4.2.1. [?]

`eqn:schanuel`

$$(4.1) \quad N(\mathbb{P}^n(F), B) \sim \frac{h_F R_F (n+1)^{r_1+r_2-1}}{w_F \zeta_F(n+1)} \left(\frac{2^{r_1} (2\pi)^{r_2}}{\sqrt{\text{disc}(F)}} \right)^{n+1} B^{n+1},$$

where

- h_F is the class number of F ;
- R_F the regulator;
- r_1 (resp. r_2) the number of real (resp. pairs of complex) embeddings of F ,
- $\text{disc}(F)$ the discriminant;
- w_F the number of roots of 1 in F ;
- ζ_F the zeta function of F .

With this starting point, one may try to prove asymptotic formulas of similar precision for arbitrary projective algebraic varieties X , at least under some natural geometric conditions. This program was initiated in [FMT89] and it has rapidly grown in recent years.

sect:experiments

4.3. Experiments. Experimental data

Elsenhans/Jahnel

4.4. Upper bounds. Pila (1995)

Theorem 4.4.1. *Let $X \subset \mathbb{P}^n$ be a geometrically irreducible variety, and $\epsilon > 0$. Then*

$$N(X, B) \ll_{\deg(X), \dim(X), \epsilon} B^{\dim(X) + \frac{1}{\deg(X)} + \epsilon}$$

Based on previous work of Bombieri-Pila (1989) - geometry of numbers.

- Pila (1996): Let $X \subset \mathbb{A}^2$ be a geometrically irreducible affine curve. Then

$$\#\{x \in X(\mathbb{Z}) \mid H(x) \leq B\} \ll_{\deg(X)} B^{\frac{1}{\deg(X)}} \log(B)^{2\deg(X)+3}.$$

- Ellenberg/Venkatesh (2005): Let $X \subset \mathbb{P}^2$ be a geometrically irreducible curve of genus ≥ 1 . Then there is a $\delta > 0$ such that

$$N(X, B) \ll_{\deg(X), \delta} B^{\frac{2}{\deg(X)} - \delta}.$$

Browning, Heath-Brown, Salberger (2006)

Theorem 4.4.2. *Let $X \subset \mathbb{P}^n$ be a geometrically irreducible variety, and $\epsilon > 0$. Then*

$$N(X, B) \ll_{\deg(X), \dim(X), \epsilon} \begin{cases} B^{\dim(X) - \frac{3}{4} + \frac{5}{3\sqrt{3}} + \epsilon} & \deg(X) = 3 \\ B^{\dim(X) - \frac{2}{3} + \frac{3}{2\sqrt{\deg(X)}} + \epsilon} & \deg(X) = 4, 5 \\ B^{\dim(X) + \epsilon} & \deg(X) \geq 6 \end{cases}$$

4.5. Lower bounds. Lower bounds for Del surfaces

lem:lower

Lemma 4.5.1. *Let X be a smooth Fano variety over a number field F and $Y := \text{Bl}_Z(X)$ a blowup in a smooth subvariety $Z = Z_F$ of codimension ≥ 2 . If $N(X^\circ(F), -K_X, B) \gg B^1$, for all dense Zariski open $X^\circ \subset X$ then the same holds for Y :*

$$N(Y^\circ(F), -K_Y, B) \gg B^1.$$

Proof.

□

In particular, *split* Del Pezzo surfaces X_d satisfy the lower bound of Conjecture 4.12.1

$$N(X_d(F), -K_{X_d}, B) \gg B^1.$$

Finer lower bounds, in some nonsplit cases have been proved in [?], [?]:

$$N(X_3^\circ(F), \mathbf{B}) \gg \mathbf{B} \log(\mathbf{B})^{r-1},$$

if X_3 is a cubic surface with at least two skew lines defined over F . This gives support to Conjecture 4.12.2. The following theorem provides evidence for Conjecture 4.12.1 in dimension 3.

`theo:manin-3`

Theorem 4.5.2. [?] *Let X be a Fano threefold over a number field F_0 . For every Zariski open subset $X^\circ \subset X$ there exists a finite extension F/F_0 such that*

$$N(X(F), -K_X, \mathbf{B}) \gg \mathbf{B}^1$$

This relies on the classification of Fano threefolds (see [IP99b], [MM82], [MM86]). One case was missing from the classification when [?] was published; the Fano threefold obtained as a blowup of $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ is a curve of tri-degree $(1, 1, 3)$ [MM03]. Lemma ?? proves the expected lower bound in this case as well.

4.6. Finer issues.

- *local or global obstructions:*

$$x_0^2 + x_1^2 + x_2^2 = 0 \text{ or } x_0^3 + 4x_1^3 + 10x_2^3 + 25x_3^3 = 0$$

- *singularities:*

$x_1^2x_2^2 + x_2^2x_3^2 + x_3^2x_1^2 = x_0x_1x_3x_4$ has $\sim B^{3/2}$ points of height $\leq B$, on every Zariski open subset

accumulating subvarieties:

For example, $x_0^3 + x_1^3 + x_2^3 + x_3^3 = 0$ has $\sim B^2$ points on \mathbb{Q} -lines and provably $O(B^{4/3+\epsilon})$ points in the complement [?]. The expectation is $\mathbf{B} \log(\mathbf{B})^3$, over \mathbb{Q} . Similar effects persist in higher dimensions. A quartic $X_4 \subset \mathbb{P}^4$ contains a 1-parameter family of lines, each contributing $\sim B^2$ to the asymptotic, while the expectation is $\sim \mathbf{B}$. Lines on a cubic $X_3 \subset \mathbb{P}^4$ are parametrized by a surface, which is of general type. We expect $\sim \mathbf{B}^2$ points of height $\leq \mathbf{B}$ on the cubic threefold, and on each line. In [?, Theorem 11.10.11] it is shown that

$$N_{\text{lines}}(\mathbf{B}) \sim c \mathbf{B}^2, \quad \text{as } \mathbf{B} \rightarrow \infty,$$

where the count is over F -rational points on lines defined over F , and the constant c is a *converging* sum of leading terms of contributions from each line of the type (4.1). In particular, each line contributes a positive density to the main term. On the other hand, one expects the same asymptotic $\sim \mathbf{B}^2$ on the complement of lines, with the leading term a product of local

densities. How to reconcile this? The forced compromise is to discard such *accumulting* subvarieties and to hope that for some Zariski open subset $U \subset X$, the asymptotic of points of bounded height does reflect the geometry of X .

$X_f \subset \mathbb{P}^n$ - open questions

If X_f is smooth, with $\deg(f) \leq n$ and $n \geq 4$ then

- *Potential density*: there exists a finite extension F/\mathbb{Q} such that $X(F)$ is Zariski dense;
- *Growth*: $N(X_f(F), B) \sim \tau \cdot B^{n+1-d}$;
- *Tamagawa number*: $\tau = \prod_v \tau_v$ - product of local densities.

Hold when $\deg(f) = 2$ and $n \geq 2$, except when $n = 3$ and the discriminant of the quadric is a square: $B^2 \log(B)$.

Consider the variety $X \subset \mathbb{P}^5$ over \mathbb{Q} given by

$$x_0x_1 - x_2x_3 + x_4x_5 = 0.$$

It is visibly a quadric hypersurface and we could apply the circle method as in Section ???. It is also the Grassmannian variety $\text{Gr}(2, 4)$ and an equivariant compactification of \mathbb{G}_a^4 . We could count points using any of the structures.

Cubic surfaces $X \subset \mathbb{P}^3$ - conjectures

- *Global obstruction*: the Brauer-Manin obstruction to the Hasse principle is the only one;
- *Linear Growth*: $N(X(F), B) \sim c \cdot B^1 \log(B)^{r-1}$, where $r = \text{rk Pic}(X)$;
- *Leading coefficient*: $c = \alpha\beta\tau$, where $\alpha \in \mathbb{Q}$, $\beta = |\text{H}^1(\text{Gal}, \text{Pic}(\bar{X}))|$, and

$$\tau = \int_{X(F)} \omega_{\mathcal{K}}$$

a Tamagawa type number.

For X *split* over \mathbb{Q} : $\alpha = 1/(7! \cdot 120)$, $\beta = 1$ and $\tau = \prod_p \tau_p \cdot \tau_\infty$ is the product of local densities, for almost all p :

$$\tau_p = \left(1 + \frac{7}{p} + 1\right) \left(1 - \frac{1}{p}\right)^7.$$

Cubic surfaces $X \subset \mathbb{P}^3$ - results

Upper bound - Heath-Brown (2002):

$$N(X^\circ(F), B) \ll_\epsilon B^{52/27+\epsilon},$$

where X° is the complement to F -lines on X

Salberger (2006): $N(X^\circ(F), B) \ll_\epsilon B^{\sqrt{3}+\epsilon}$

Asymptotic - Derenthal (2005): linear growth + leading coefficient for the E_6 -cubic:

$$x_1x_2^2 + x_2x_0^2 + x_3^3 = 0.$$

conj:layer

Conjecture 4.6.1. Let X be a K3 surface over a number field F . Let L be a polarization, $\epsilon > 0$ and $Y = Y(\epsilon, L)$ be the union of all F -rational curves $C \subset X$ (i.e., curves that are isomorphic to \mathbb{P}^1 over F) and have L -degree $\leq 2/\epsilon$. Then

$$N(X, L, B) = N(Y, L, B) + O(B^\epsilon), \quad \text{as } B \rightarrow \infty.$$

Theorem 4.6.2. [McK00] *Let $X \rightarrow \mathbb{P}^1 \times \mathbb{P}^1$ be a double cover ramified over a curve of bidegree $(4, 4)$. Then there exists an open cone $\Lambda \subset \Lambda_{\text{ample}}(X)$ such that for every $L \in \Lambda$ there exists a $\delta > 0$ such that*

$$N(X, L, B) = N(Y, L, B) + O(B^{2/d-\delta}), \quad \text{as } B \rightarrow \infty,$$

where d is the minimal L -degree of a rational curve on X and Y is the union of all F -rational curves of degree d .

This theory exhibits the first layer of an arithmetic stratification predicted in Conjecture 4.6.1.

4.7. Upper bounds. Fibration method

[BHBS06]

sect:circle

4.8. The circle method. Let $f \in \mathbb{Z}[x_0, \dots, x_n]$ be a homogeneous polynomial of degree d such that the hypersurface $X_f \subset \mathbb{P}^n$ is nonsingular. Let

$$N_f(B) := \#\{\mathbf{x} \in \mathbb{Z}^n \mid f(\mathbf{x}) = 0 \text{ and } \|\mathbf{x}\| \leq B\}$$

be the counting function. In this section we sketch a proof of the following

thm:birch

Theorem 4.8.1. [Bir62] *Assume that $n \geq 2^d(d+1)$. Then*

$$(4.2) \quad N_f(B) = \Theta \cdot B^{n+1-d}(1 + o(1)) \quad B \rightarrow \infty,$$

where

$$\Theta = \prod_p \tau_p \cdot \tau_\infty > 0,$$

provided $f(\mathbf{x}) = 0$ is solvable in \mathbb{Z}_p , for all p , and in \mathbb{R} .

The constants τ_p and τ_∞ admit an interpretation as *local densities*; these are explained in a more conceptual framework in Section 4.14.

eqn:circle-theta

Substantial efforts have been put into reducing the number of variables, especially for low degrees. (See HB 2006, 2007...) Another direction is the extension of the method to more general number fields [Ski97] and even function fields.

We now outline the main steps of the proof of the asymptotic formula 4.2. The first step is the introduction of a “delta”-function: for $x \in \mathbb{Z}$ we have

$$\int_0^1 e^{2\pi i \alpha x} d\alpha = \begin{cases} 0 & \text{if } x \neq 0 \\ 1 & \text{otherwise} \end{cases}$$

Now we can write

eqn:N (4.3)
$$N_f(B) = \int_0^1 S(\alpha) d\alpha$$

where

$$S(\alpha) := \sum_{\mathbf{x} \in \mathbb{Z}^{n+1}, \|\mathbf{x}\| \leq B} e^{2\pi i \alpha f(\mathbf{x})}.$$

The function $S(\alpha)$ is wildly oscillating (see Figure 4.8), with peaks at $\alpha = a/q$, with small q . Indeed, the probability that $f(\mathbf{x})$ is divisible by q is higher for small q , and each such term contributes 1 to $S(\alpha)$. The idea of the circle method is to analyze the asymptotic of the integral in equation 4.3, for $B \rightarrow \infty$, by extracting the contributions of α close to rational numbers a/q with small q , and finding appropriate bounds for integrals over the remaining intervals.

[?], [HB06]

More precisely, one introduces the *major arcs*

$$\mathfrak{M} := \bigcup_{(a,q)=1, q \leq B^\Delta} \mathfrak{M}_{a,q},$$

where $\Delta > 0$ is a parameter to be specified, and

$$\mathfrak{M}_{a,q} := \left\{ \alpha \mid \left| \alpha - \frac{a}{q} \right| \leq B^{-d+\delta} \right\}.$$


The *minor arcs* are the complement:

$$\mathfrak{m} := [0, 1] \setminus \mathfrak{M}.$$

The goal is to prove the bound

eqn:minor (4.4)
$$\int_{\mathfrak{m}} S(\alpha) d\alpha = O(B^{n-d-\epsilon}), \text{ for some } \epsilon > 0,$$

fig:s

FIGURE 1. Oscillations of $S(\alpha)$

and the asymptotic

$$(4.5) \quad \int_{\mathfrak{M}} S(\alpha) d\alpha \sim \prod_p \tau_p \cdot \tau_\infty \cdot B^{n+1-d} \quad \text{for } B \rightarrow \infty.$$

Remark 4.8.2. Modern refinements employ “smoothed out” intervals, i.e., the delta function of an interval in the major arcs is replaced by a smooth bell curve with support in this interval. In Fourier analysis, “rough edges” translate into bad bounds on the dual side, and should be avoided. An implementation of this idea, leading to savings in the number of variables can be found in [HB83].

There are various approaches to proving upper bounds in equation 4.4, most are a variant or refinement of Weyl’s bounds (1916) [Wey16]. Weyl considered the following exponential sums

$$s(\alpha) := \sum_{0 \leq x \leq B} e^{2\pi i \alpha x^d},$$

The main observation is that $|\mathfrak{s}(\alpha)|$ is “small”, when $|\alpha - a/q|$ “large”. This is easy to see when $d = 1$; summing the geometric series we get

$$|\mathfrak{s}(\alpha)| = \left| \frac{1 - e^{2\pi i \alpha (B+1)}}{1 - e^{2\pi i \alpha}} \right| \leq \frac{1}{2\alpha} \dots$$

We turn to major arcs. Let

$$\alpha = \frac{a}{q} + \beta$$

with β *very small*, and getting smaller as a function of B . Here we will assume that $|\beta| \leq B^{-d+\delta'}$, for some small $\delta' > 0$. We put $\mathbf{x} = q\mathbf{y} + \mathbf{z}$, with \mathbf{z} the corresponding residue class modulo q , and obtain

$$\begin{aligned} \mathfrak{S}(\alpha) &= \sum_{\mathbf{x} \in \mathbb{Z}^{n+1}, \|\mathbf{x}\| \leq B} e^{2\pi i \frac{a}{q} f(\mathbf{x})} e^{2\pi i \beta f(\mathbf{x})} \\ &= \sum_{\|\mathbf{x}\| \leq B} e^{2\pi i \frac{a}{q} f(q\mathbf{y} + \mathbf{z})} e^{2\pi i \beta f(\mathbf{x})} \\ &= \sum_{\mathbf{z}} e^{2\pi i \frac{a}{q} f(\mathbf{z})} \left(\sum_{\|\mathbf{y}\| \leq B/q} e^{2\pi i \beta f(\mathbf{x})} \right) \\ &= \sum_{\mathbf{z}} e^{2\pi i \frac{a}{q} f(\mathbf{z})} \int_{\|\mathbf{y}\| \leq B/q} e^{2\pi i \beta f(\mathbf{x})} d\mathbf{y} \\ &= \sum_{\mathbf{z}} \frac{e^{2\pi i \frac{a}{q} f(\mathbf{z})}}{q^{n+1}} \int_{\|\mathbf{x}\| \leq B} e^{2\pi i \beta f(\mathbf{x})} d\mathbf{x}, \end{aligned}$$

where $d\mathbf{y} = q^{n+1}d\mathbf{x}$. The passage $\sum \mapsto \int$ is justified for our choice of small β - the difference will be adsorbed in the error term in (4.2). We have obtained

$$\int_0^1 \mathfrak{S}(\alpha) d\alpha \sim \sum_{a,q} \sum_{\mathbf{z}} \frac{e^{2\pi i \frac{a}{q} f(\mathbf{z})}}{q^{n+1}} \cdot \int_{|\beta| \leq B^{-d+\delta}} \int_{\|\mathbf{x}\| \leq B} e^{2\pi i \beta f(\mathbf{x})} d\mathbf{x} d\beta.$$

We first deal with the integral on the right, called the *singular integral*. Put $\beta' = \beta B^d$ and $\mathbf{x}' = \mathbf{x}/B$. The change of variables leads to

$$\int_{|\beta| \leq \frac{1}{B^{d-\delta}}} d\beta \int_{\|\mathbf{x}\| \leq B} e^{2\pi i \beta B^d f(\frac{\mathbf{x}}{B})} B^{n+1} d(\frac{\mathbf{x}}{B}) = B^{n+1-d} \int_{|\beta'| \leq B^\delta} \int_{\|\mathbf{x}'\| \leq 1} e^{2\pi i \beta' f(\mathbf{x}')} d(\mathbf{x}').$$

We see the appearance of the main term B^{n-d} and the density

$$\tau_\infty := \int_0^1 d\beta' \int_{\|\mathbf{x}'\| \leq 1} e^{2\pi i \beta' f(\mathbf{x}')} d\mathbf{x}'.$$

Now we analyze the *singular integral*

$$\sigma_Q := \sum_{a,q} \sum_{\mathbf{z}} \frac{e^{2\pi i \frac{a}{q} f(\mathbf{z})}}{q^n},$$

where the outer sum runs over positive coprime integers a, q , $a < q$ and $q < Q$, and the inner sum over residue classes $\mathbf{z} \in (\mathbb{Z}/q)^{n+1}$. This sum has the following properties

- (1) multiplicativity in q , in particular we have

$$\sigma := \prod_p \left(\sum_{i=0}^{\infty} A(p^i) \right).$$

- (2) with $\sigma_Q \rightarrow \sigma$, for $Q \rightarrow \infty$, (with small error term),

$$\sum_{i=0}^k \frac{A(p^i)}{p^{i(n+1)}} = \frac{\varrho(f, p^k)}{p^{kn}},$$

where

$$\varrho(f, p^k) := \#\{ \mathbf{z} \pmod{p^k} \mid f(\mathbf{z}) = 0 \pmod{p^k} \}.$$

Here, a discrete version of equation (4.3) comes into play:

$$\#\{\text{solutions} \pmod{p^k}\} = \frac{1}{p^k} \sum_{a=0}^{p^k-1} \sum_{\mathbf{z}} e^{2\pi i \frac{af(\mathbf{z})}{p^k}}$$

However, our sums run over a with $(a, p) = 1$. A rearranging of terms leads to

$$\begin{aligned} \frac{\varrho(f, p^k)}{p^{kn}} &= \sum_{i=0}^k \sum_{(a, p^k)=p^i} \sum_{\mathbf{z}} \frac{1}{p^{k(n+1)}} e^{2\pi i \frac{a}{p^k} f(\mathbf{z})} \\ &= \sum_{i=0}^k \sum_{(\frac{a}{p^i}, p^{k-i})=1} \sum_{\mathbf{z}} \frac{1}{p^{k(n+1)}} e^{2\pi i \frac{a/p^i}{p^{k-i}} f(\mathbf{z})} \cdot p^{(n+1)i} \\ &= \sum_{i=0}^k \frac{1}{p^{(n+1)(k-i)}} \sum_{(a, p^{k-i})=1} \sum_{\mathbf{z}} e^{2\pi i \frac{a}{p^{k-i}} f(\mathbf{z})} \\ &= \sum_{i=0}^k \frac{1}{p^{(n+1)(k-i)}} \cdot A(p^{k-i}). \end{aligned}$$

In conclusion,

eqn:euler (4.6)
$$\sigma = \prod_p \tau_p, \quad \text{where } \tau_p = \lim_{k \rightarrow \infty} \frac{\varrho(f, p^k)}{p^{nk}}.$$

As soon as there is at least one (nonsingular) solution $f(\mathbf{z}) = 0 \pmod p$, $\tau_p \neq 0$, and in fact, for almost all p ,

$$\frac{\varrho(f, p^k)}{p^{nk}} = \frac{\varrho(f, p)}{p^n},$$

by Hensel's lemma. Moreover, if $\tau_p \neq 0$ for all p , the the Euler product in equation (4.6) converges.

Let us illustrate this in the example of Fermat type equations

$$f(\mathbf{x}) = a_0x^d + \dots + a_nx_n^d = 0.$$

Using properties of Jacobi sums one can show that

$$\varrho(f, p) = p^n + E, \quad \text{with } E = O((p-1)p^{(n-1)/2}),$$

so that

$$\left| \frac{\varrho(f, p)}{p^{n-1}} - 1 \right| \leq \frac{C}{p^{(n+1)/2}}.$$

The corresponding Euler product

$$\prod_p \frac{\varrho(f, p)}{p^n} \ll \prod_p \left(1 + \frac{C}{p^{(n+1)/2}}\right)$$

is convergent.

Some historical background: the cicle method was firmly established in the series of papers of Hardy and Littlewood *Partitio numerum*. They comment: "A method of great power and wide scope, applicable to almost any problem concernign the decomposition of integers into parts of a particular kind, and to many against which it is difficult to suggest any other obvious method of attack."

chap:funct

4.9. Function fields: heuristics. Let p be a prime and put $q = p^n$. Let Λ be a convex n -dimensional cone in \mathbb{R}^n with vertex at 0. Let

$$f_1, f_2 : \mathbb{R}^n \rightarrow \mathbb{R}$$

be two linear functions such that

- $f_i(\mathbb{Z}^n) \subset \mathbb{Z}$;
- $f_2(x) > 0$ for all $x \in \Lambda \setminus \{0\}$;
- there exists an $x \in \Lambda \setminus \{0\}$ such that $f(x) > 0$.

Put

$$M = \cup_{\lambda} M_{\lambda}, \quad \lambda \in \mathbb{Z}^n \cap \Lambda$$

and

$$|M_{\lambda}| := q^{\max(0, f_1(\lambda))}, \quad \varphi(m) := q^{f_2(\lambda)}, \quad \text{for } m \in M_{\lambda}.$$

Then the series

$$F(s) = \sum_{\lambda \in \Lambda \cap \mathbb{Z}^n} \frac{|M_{\lambda}|}{q^{s f_2(\lambda)}}$$

converges for

$$\Re(s) > a := \max_{x \in \Lambda} (f_1(x)/f_2(x)) > 0.$$

What happens around $s = a$? Choose an $\epsilon > 0$ and decompose the cone

$$\Lambda := \Lambda_{\epsilon}^{+} \cup \Lambda_{\epsilon}^{-},$$

where

$$\begin{aligned} \Lambda_{\epsilon}^{+} &:= \{x \in \Lambda \mid f_1(x)/f_2(x) \geq a - \epsilon\} \\ \Lambda_{\epsilon}^{-} &:= \{x \in \Lambda \mid f_1(x)/f_2(x) < a - \epsilon\} \end{aligned}$$

Therefore,

$$F(s) = F_{\epsilon}^{+} + F_{\epsilon}^{-},$$

where F_{ϵ}^{-} converges absolutely for $\Re(s) > a - \epsilon$.

Now we make some assumptions concerning Λ : suppose that for all $\epsilon \in \mathbb{Q}_{>0}$, the cone Λ_{ϵ}^{+} is a rational finitely generated polyhedral cone. Then

$$\Lambda_{\epsilon}^a := \{x \mid f_1(x)/f_2(x) = a\}$$

is a face of Λ_{ϵ}^{+} , and thus also finitely generated polyhedral.

Lemma 4.9.1. *There exists a function $G_{\epsilon}(s)$, holomorphic for $\Re(s) > a - \epsilon$, such that*

$$F_{\epsilon}(s) = \frac{G_{\epsilon}(s)}{(s - a)^b},$$

where b is the dimension of the face Λ_{ϵ}^a .

Proof. For $y \in \mathbb{Q}_{>0}$ we put

$$P(y) := \{x \mid x \in \Lambda, f_2(x) = y\}$$

Consider the expansion

$$F(s) = \sum_{y \in \mathbb{N}} \sum_{\lambda \in P(y) \cap \mathbb{Z}^n} q^{f_1(\lambda) - s f_2(\lambda)}$$

Replacing by the integral, we obtain (with $w = yz$)

$$\begin{aligned}
 &= \int_0^\infty dy \left(\int_{P(y)} q^{f_1(w) - sf_2(w)} dw \right) \\
 &= \int_0^\infty dy \left(\int_{P(1)} y^{n-1} q^{f_1(w) - sf_2(y)} dz \right) \\
 (4.7) \quad &= \int_{P(1)} dz \int_0^\infty y^{n-1} q^{(f_1(w) - s)y} dy \\
 &= \int_{P(1)} dz \frac{1}{(s - f_1(z))^n} \int_0^\infty u^{n-1} q^{-u} du \\
 &= \frac{\Gamma(n)}{(\log(q))^n} \int_{P(1)} \frac{1}{(s - f_1(z))^n} dz.
 \end{aligned}$$

It is already clear that we get a singularity at $s = \max(f_1(z))$ on $P(1)$, which is a . In general, let f be a linear function and

$$\Phi(s) := \int_{\Delta} (s - f(x))^{-n} d\Omega$$

where Δ is a polytope of dimension $n - 1$. Then Φ is a rational function in s , with an asymptotic at $s = a$ given by

$$\text{vol}_{f,a} \frac{(b-1)!}{(n-1)!} (s-a)^{-b},$$

where $\Delta_{f,a}$ is the polytope $\Delta \cap \{f(x) = a\}$, $\text{vol}_{f,a}$ is its volume and $b = 1 + \dim \Delta_{f,a}$. \square

Let C be a curve of genus g over the finite field \mathbb{F}_q and F its function field. Let X be a variety over \mathbb{F}_q of dimension n . Then $V := X \times S$ is a variety over F . Every F -rational point x of V gives rise to a section \tilde{x} of the map $V \rightarrow S$. We have a pairing

$$A^1(V) \times A^n(V) \rightarrow \mathbb{Z}$$

between the groups of (numerical) equivalence classes of codimension 1-cycles and codimension n -cycles. We have

$$A^n(V) = A^n(X) \otimes A^1(S) \oplus A^{n-1}(X) \otimes A^0(S)$$

and

$$\begin{aligned}
 A^1 &= A^1(X) \oplus \mathbb{Z} \\
 L &= (L_X, \ell) \\
 -K_V &= (-K_X, 2 - 2g)
 \end{aligned}$$

Let L be a very ample line bundle on V . Then

$$q^{(L, \tilde{x})}$$

is the height of the point x with respect to L . The height zeta function takes the form

$$\begin{aligned} Z(s) &= \sum_{x \in V(F)} q^{-(L, \tilde{x})s} \\ &= \sum_{y \in A^n(X)} \tilde{\mathbf{N}}(q) a^{-[(L_X, y) + \ell]s}, \end{aligned}$$

where

$$\tilde{\mathbf{N}}(q) := \#\{x \in V(F) \mid \text{cl}(x) = y\}.$$

We proceed to give some *heuristic*(!) bound on $\tilde{\mathbf{N}}(q)$. The cycles in a given class y are parametrized by an algebraic variety M_y and

$$\dim M_{y(\tilde{x})} \geq \chi(\mathcal{N}_{V|\tilde{x}})$$

(the Euler characteristic). More precisely, the local ring on the moduli space is the quotient of a powerseries ring with $h^0(\mathcal{N}_{V|\tilde{x}})$ generators by $h^1(\mathcal{N}_{V|\tilde{x}})$ relations. Our main heuristic assumption is that

$$\tilde{\mathbf{N}}(q) \sim q^{\dim M_y} \sim q^{\chi(\mathcal{N}_{V|\tilde{x}})}.$$

This assumption fails, for example, for points contained in “exceptional” (accumulating) subvarieties.

By the short exact sequence

$$0 \rightarrow \mathcal{T}_{\tilde{x}} \rightarrow \mathcal{T}_{V|\tilde{x}} \rightarrow \mathcal{N}_{V|\tilde{x}} \rightarrow 0$$

we have

$$\begin{aligned} \chi(\mathcal{T}_{V|\tilde{x}}) &= (-K_V, \tilde{x}) + (n+1)\chi(\mathcal{O}_{\tilde{x}}) \\ \chi(\mathcal{N}_{V|\tilde{x}}) &= (-K_X, \text{cl}(x)) + n\chi(\mathcal{O}_{\tilde{x}}) \end{aligned}$$

From now on we consider a *modified* height zeta function

$$Z_{\text{mod}}(s) := \sum q^{\chi(\mathcal{N}_{V|\tilde{x}}) - (L, \tilde{x})s}.$$

We observe that its analytic properties are determined by the ratio between two linear functions

$$(-K_X, \cdot) \text{ and } (L, \cdot).$$

The relevant cone Λ is the cone spanned by classes of (maximally moving) effective curves. The *finite* generation of this cone for Fano varieties is one of the main results of Mori's theory. We conclude that

$$\mathbf{N}_U(L, B) \sim \mathbf{B}^a(\log(\mathbf{B}))^{b-1},$$

where

$$a = a(L) = \max_{z \in \Lambda} ((-K_X, z)/(L, z))$$

and $b = b(L)$ is the dimension of the face of the cone where this maximum is achieved.

sect:metrization

4.10. Metrizations of line bundles. In this section we discuss a refined theory of height functions, based on the notion of an adelically metrized line bundle.

Let F be a number field and $\text{disc}(F)$ the discriminant of F (over \mathbb{Q}). The set of places of F will be denoted by $\text{Val}(F)$. We shall write $v|\infty$ if v is archimedean and $v \nmid \infty$ if v is nonarchimedean. For any place v of F we denote by F_v the completion of F at v and by \mathfrak{o}_v the ring of v -adic integers (for $v \nmid \infty$). Let q_v be the cardinality of the residue field \mathbb{F}_v of F_v for nonarchimedean valuations. The local absolute value $|\cdot|_v$ on F_v is the multiplier of the Haar measure, i.e., $d(ax_v) = |a|_v dx_v$ for some Haar measure dx_v on F_v . We denote by $\mathbb{A} = \mathbb{A}_F = \prod'_v F_v$ the adèle ring of F . We have the *product formula*

$$\prod_{v \in \text{Val}(F)} |a|_v = 1, \quad \text{for all } a \in F^*.$$

defn:metri

Definition 4.10.1. Let X be an algebraic variety over F and L a line bundle on X . A v -adic metric on L is a family $(\|\cdot\|_x)_{x \in X(F_v)}$ of v -adic Banach norms on the fibers L_x such that for all Zariski open subsets $U \subset X$ and every section $\mathbf{f} \in H^0(U, L)$ the map

$$U(F_v) \rightarrow \mathbb{R}, \quad x \mapsto \|\mathbf{f}\|_x,$$

is continuous in the v -adic topology on $U(F_v)$.

exam:metr

Example 4.10.2. Assume that L is generated by global sections. Choose a basis $(\mathbf{f}_j)_{j \in [0, \dots, n]}$ of $H^0(X, L)$ (over F). If \mathbf{f} is a section such that $\mathbf{f}(x) \neq 0$ then define

$$\|\mathbf{f}\|_x := \max_{0 \leq j \leq n} \left(\left| \frac{\mathbf{f}_j}{\mathbf{f}}(x) \right|_v \right)^{-1},$$

otherwise $\|0\|_x := 0$. This defines a v -adic metric on L . Of course, this metric depends on the choice of $(\mathbf{f}_j)_{j \in [0, \dots, n]}$.

defn:ad-metric

Definition 4.10.3. Assume that L is generated by global sections. An adelic metric on L is a collection of v -adic metrics, for every $v \in \text{Val}(F)$, such that for all but finitely many $v \in \text{Val}(F)$ the v -adic metric on L is defined by means of some *fixed* basis $(f_j)_{j \in [0, \dots, n]}$ of $H^0(X, L)$.

We shall write $\|\cdot\|_{\mathbb{A}} := (\|\cdot\|_v)$ for an adelic metric on L and call a pair $\mathcal{L} = (L, \|\cdot\|_{\mathbb{A}})$ an adelically metrized line bundle. Metrizations extend naturally to tensor products and duals of metrized line bundles, which allows to define adelic metrizations on arbitrary line bundles L (on projective X): represent L as $L = L_1 \otimes L_2^{-1}$ with very ample L_1 and L_2 . Assume that L_1, L_2 are adelically metrized. An adelic metrization of L is any metrization which for all but finitely many v is induced from the metrizations on L_1, L_2 .

defn:height

Definition 4.10.4. Let $\mathcal{L} = (L, \|\cdot\|_{\mathbb{A}})$ be an adelically metrized line bundle on X and f an F -rational section of L . Let $U \subset X$ be the maximal Zariski open subset of X where f is defined and does not vanish. For all $x = (x_v)_v \in U(\mathbb{A})$ we define the local

$$H_{\mathcal{L}, f, v}(x_v) := \|f\|_{x_v}^{-1}$$

and the global *height function*

$$H_{\mathcal{L}}(x) := \prod_{v \in \text{Val}(F)} H_{\mathcal{L}, f, v}(x_v).$$

By the product formula, the restriction of the global height to $U(F)$ does not depend on the choice of f .

exam:p1-height

Example 4.10.5. For $X = \mathbb{P}^1 = (x_0 : x_1)$ one has $\text{Pic}(X) = \mathbb{Z}$, spanned by the class $L = [(1 : 0)]$. For all $f = x_0/x_1 \in \mathbb{G}_a(\mathbb{A})$ we define

$$H_{\mathcal{L}, f, v}(x_v) = \max(1, |f|_v).$$

The restriction of $H_{\mathcal{L}} = \prod_v H_{\mathcal{L}, f, v}$ to $\mathbb{G}_a(F) \subset \mathbb{P}^1$ is the usual height on \mathbb{P}^1 (with respect to the usual metrization of $\mathcal{L} = \mathcal{O}(1)$).

exam:unipotent-height

Example 4.10.6. Let X be an equivariant compactification of a unipotent group G and L a very ample line bundle on X . The space $H^0(X, L)$, a representation space for G , has a *unique* G -invariant section f , modulo scalars. Indeed, if we had two nonproportional sections, their quotient would be a character of G , which are trivial.

Fix such a section. We have $f(g_v) \neq 0$, for all $g_v \in G(F_v)$. Put

$$H_{\mathcal{L}, f, v}(g_v) = \|f(g_v)\|^{-1} \quad \text{and} \quad H_{\mathcal{L}, f} = \prod_v H_{\mathcal{L}, f, v}.$$

By the product formula, the global height is independent of the choice of f .

sect:hei-z

4.11. Height zeta functions. Let X be an algebraic variety over a global field F , $\mathcal{L} = (L, \|\cdot\|_{\mathbb{A}})$ an adelicly metrized ample invertible sheaf on X , $H_{\mathcal{L}}$ a height function associated to \mathcal{L} , U a subvariety of X , $a_U(\mathcal{L})$ the convergence abscissa of the height zeta function

$$Z_U(\mathcal{L}, s) := \sum_{x \in U(F)} H_{\mathcal{L}}(x)^{-s}.$$

prop:al

Proposition 4.11.1.

- (1) *The value of $a_U(\mathcal{L})$ depends only on the class of L in $\text{NS}(X)$.*
- (2) *Either $0 \leq a_U(\mathcal{L}) < \infty$, or $a_U(\mathcal{L}) = -\infty$, the latter possibility corresponding to the case of finite $U(F)$. If $a_U(\mathcal{L}) > 0$ for one ample L then this is so for every ample L .*
- (3) *$a_U(\mathcal{L}^m) = \frac{1}{m}a_U(\mathcal{L})$. In general, $a_U(\mathcal{L})$ extends uniquely to a continuous function on $\Lambda_{\text{nef}}(X)^{\circ}$, which is inverse linear on each half-line unless it identically vanishes.*

Proof. All statements follow directly from the listed properties of heights (see [?]). In particular,

$$a_U(\mathcal{L}) \leq a(\mathbb{P}^n(F), \mathcal{O}(m)) = \frac{n+1}{m}$$

for some n, m . If $Z_U(\mathcal{L}, s)$ converges at some negative s , then it must be a finite sum. Since for two ample heights H, H' we have

$$cH^m < H' < c'H^n, \quad c, c', m, n > 0,$$

the value of a can only be simultaneously positive or zero. Finally, if L and L' are close in the (real) topology of $\text{NS}(V)_{\mathbb{R}}$, then $L - L'$ is a linear combination of ample classes with small coefficients, and so $a_U(\mathcal{L})$ is close to $a_U(\mathcal{L}')$. \square

nota:al

Notation 4.11.2. By Property (1) of Proposition 4.11.1, we may write $a_U(\mathcal{L}) = a_U(L)$.

exam:ab-v

Example 4.11.3. For an abelian variety X and ample invertible sheaf L we have

$$H_{\mathcal{L}} = \exp(q(x) + \ell(x) + O(1)),$$

where q is a positive definite quadratic form on the space $X(F) \otimes \mathbb{Q}$ and ℓ is a linear form. It follows that $a_X(L) = 0$, although $X(F)$ may well be Zariski dense in V . Also

$$N_X(\mathcal{L}, B) \sim \log(\mathbf{B})^{r/2}$$

where $r = \text{rk } X(F)$. Hence for $a = 0$, the power of $\log(\mathbf{B})$ in principle cannot be calculated geometrically: it depends on the arithmetic of X and F . The hope is that for $a > 0$ the situation is more stable.

defn:lin-gr

Definition 4.11.4. The *arithmetic hypersurface of linear growth* Σ_U is

$$\Sigma_U^{\text{arith}} := \{L \in \text{NS}(X)_{\mathbb{R}} \mid a_U(L) = 1\}.$$

Obviously, it defines a completely. More precisely,

prop:alpha

Proposition 4.11.5.

- If $a_U(L) > 0$ for some L , then Σ_U is nonempty and intersects each half-line in $\Lambda_{\text{eff}}(X)^\circ$ in exactly one point.
- $\Sigma_U^{\leq} := \{L \mid a_U(L) < 1\}$ is convex.

Proof. The first statement is clear. The second follows from the Hölder inequality: if

$$0 < \sigma, \sigma' \leq 1 \quad \text{and} \quad \sigma + \sigma' = 1$$

then

$$H_{\mathcal{L}}^{-\sigma}(x) H_{\mathcal{L}}^{\sigma'}(x) \leq \sigma H_{\mathcal{L}}(x)^{-1} + \sigma' H_{\mathcal{L}}(x)^{-1}$$

so that from $L, L' \in \Sigma_U^{\leq}$ it follows that $\sigma L + \sigma' L' \in \Sigma_U^{\leq}$. \square

When $\text{rk NS}(X) = 1$, Σ_U is either empty, or consists of one point. Schanuel's theorem ?? implies that for $\mathbb{P}^n(F)$, this point is the anticanonical class.

defn:accu-sub

Definition 4.11.6. A subvariety $X \subset U \subset V$ is called *point accumulating*, or simply *accumulating* (in U with respect to L), if

$$a_U(L) = a_X(L) > a_{U \setminus X}(L).$$

It is called *weakly accumulating*, if

$$a_U(L) = a_X(L) = a_{U \setminus X}(L).$$

exem:accum

Example 4.11.7. If we blow up an F -point of an abelian variety X , the exceptional divisor will be an accumulating subvariety in the resulting variety, although to prove this we must analyze the height with respect to the exceptional divisor, which is not quite obvious.

We will see in Theorem ?? that for the product $\mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_k} = X$, with $n_j > 0$, every fiber of a partial projection will be weakly accumulating with respect to the anticanonical class.

The role of accumulating subvarieties in our subsequent analysis will be different for various classes of varieties, but we shall generally try to pinpoint them in a geometric way. For example, on Fano varieties we shall have to remove the $-K_X$ -accumulating subvarieties if we want to ensure the linear growth conjecture.

The role of weakly accumulating subvarieties is that they open possibilities to obtain exact lower bounds for the power growth rate of $X(F)$ by analyzing subvarieties of smaller dimension.

sect:manin

4.12. Manin’s conjecture. The following picture emerged from the analysis of examples such as \mathbb{P}^n , flag varieties, complete intersections of small degree [?], [BM90].

Let X be a smooth projective variety with ample anticanonical class over a number field F_0 . The conjectures below describe the asymptotic of rational points of bounded height in a *stable* situation, i.e., after a sufficiently large finite extension F/F_0 and passing to a sufficiently small Zariski dense subset $X^\circ \subset X$.

conj:linear

Conjecture 4.12.1 (Linear growth conjecture). One has

eqn:manin

$$(4.8) \quad \mathbf{B}^1 \ll \mathbf{N}(X^\circ(F), -K_X) \ll \mathbf{B}^{1+\epsilon}.$$

conj:power

Conjecture 4.12.2 (The power of log).

eqn:log

$$(4.9) \quad \mathbf{N}(X^\circ(F), -K_X) \sim \mathbf{B}^1 \log(\mathbf{B})^{r-1},$$

where $r = \text{rk Pic}(X_F)$.

Conjecture 4.12.3 (General polarizations / linear growth). Every smooth projective X with $-K_X \in \Lambda_{\text{big}}(X)$ has a dense Zariski open subset X° such that

$$\Sigma_{X^\circ}^{\text{arith}} = \Sigma_X^{\text{geom}},$$

(see Definitions 4.11.4, ??).

The next level of precision requires that $\Lambda_{\text{eff}}(X)$ is a finitely generated polyhedral cone. By Theorem 1.1.5, this holds when X is Fano.

Conjecture 4.12.4 (General polarizations / power of log). Let L be an ample line bundle and $a(L), b(L)$ the constants defined in ... Then

eqn:gen-pol

$$(4.10) \quad \mathbf{N}(X^\circ(F), L) \sim B^{a(L)} \log(B)^{b(L)-1}.$$

sect:counter

4.13. Counterexamples. At present, no counterexamples to Conjecture 4.12.1 are known. However, Conjecture 4.12.2 fails in dimension 3. The geometric reason for this failure comes from Mori fiber spaces, more specifically from “unexpected” jumps in the rank of the Picard group in fibrations.

Let $X \subset \mathbb{P}^n$ be a smooth hypersurface. We know, by Lefschetz, that $\text{Pic}(X) = \text{Pic}(\mathbb{P}^n) = \mathbb{Z}$, for $n \geq 4$. However, this may fail when X has dimension 2. Moreover, the variation of the rank of the Picard group in a family of surfaces X_t over a number field F may be nontrivial, even when *geometrically*, i.e., over the algebraic closure \bar{F} of F , the rank is constant.

Concretely, consider a hypersurface $X \subset \mathbb{P}_{\mathbf{x}}^3 \times \mathbb{P}_{\mathbf{y}}^3$ given by a form of bidegree (1,3):

$$\sum_{j=0}^3 x_j y_j^3 = 0.$$

By Lefschetz, the Picard group $\text{Pic}(X) = \mathbb{Z}^2$, with the basis of hyperplane sections of $\mathbb{P}_{\mathbf{x}}^3$, resp. $\mathbb{P}_{\mathbf{y}}^3$, and the anticanonical class is computed as in Example ??

$$-K_X = (3, 1).$$

Projection onto $\mathbb{P}_{\mathbf{y}}^3$ exhibits X as a \mathbb{P}^2 -fibration over \mathbb{P}^3 . The second Mori fiber space structure on X is given by projection to $\mathbb{P}_{\mathbf{x}}^3$, with fibers diagonal cubic surfaces. The restriction of $-K_X$ to each (smooth) fiber $X_{\mathbf{x}}$ is the anticanonical class of the fiber.

By ..., $\text{rk Pic}(X_{\mathbf{x}})$ varies between 1 and 4. For example, if F contains $\sqrt{-3}$, then the $\text{rk Pic}(X_{\mathbf{x}}) = 4$ whenever all x_j are cubes in F . The lower bounds in Section ?? show that

$$\mathbf{N}_{X_{\mathbf{x}}^{\circ}}(-K_{X_{\mathbf{x}}}, \mathbf{B}) \sim \mathbf{B} \log(\mathbf{B})^3$$

for all such fibers, all dense Zariski open subsets $X_{\mathbf{x}}^{\circ}$ and all F . On the other hand, Conjecture ?? implies that

$$\mathbf{N}_{X^{\circ}}(-K_X, \mathbf{B}) \sim \mathbf{B} \log(\mathbf{B}),$$

for some Zariski open $X^{\circ} \subset X$, over a sufficiently large number field F . However, every Zariski open subset $X^{\circ} \subset X$ intersects infinitely many fibers $X_{\mathbf{x}}$ with $\text{rk Pic}(X_{\mathbf{x}}) = 4$ in a dense Zariski open subset. This is a contradiction.

secct:refine

4.14. Peyre’s refinement. The refinement concerns the conjectured asymptotic formula (4.9). Fix a metrization of $-K_X$. The expectation is that

eqn:peyre

$$(4.11) \quad \mathbf{N}(U(F), K_X) = c(-K_X) \cdot \mathbf{B}^1 \log(\mathbf{B})^{r-1} (1 + o(1)), \quad \text{as } \mathbf{B} \rightarrow \infty,$$

with $r = \text{rk Pic}(X)$. Peyre’s achievement was to give a conceptual interpretation of the constant $c(-K_X)$. Here we explain the key steps of his construction.

Let F be a number field and F_v its v -adic completion. Let X be a smooth algebraic variety over F of dimension d equipped with an adelicly metrized line bundle $\mathcal{K} = \mathcal{K}_X = (K_X, \|\cdot\|_{\mathbb{A}})$. Fix a point $x \in X(F_v)$ and let x_1, \dots, x_d be local analytic coordinates in an analytic neighborhood U_x of x giving a homeomorphism

$$\phi : U \xrightarrow{\sim} F_v^d.$$

Let $dy_1 \wedge \dots \wedge dy_d$ be the standard differential form on F_v^d and $\mathbf{f} := \phi^*(dy_1 \wedge \dots \wedge dy_d)$ its pullback to U . Note that \mathbf{f} is a local section of the canonical sheaf K_X and that a v -adic metric $\|\cdot\|_v$ on K_X gives rise to a norm $\|\mathbf{f}(u)\|_v \in \mathbb{R}_{>0}$, for each $u \in U_x$. Let $d\mu_v = dy_1 \cdots dy_d$ be the standard Haar measure, normalized by

$$\int_{\mathfrak{o}_v^d} d\mu_v = \frac{1}{\mathfrak{d}_v^{d/2}},$$

where \mathfrak{d}_v is the local different (which equals 1 for almost all v).

Define the local v -adic measure $\tilde{\omega}_{\mathcal{K},v}$ on U_x via

$$\int_W \tilde{\omega}_{\mathcal{K},v} = \int_{\phi(W)} \|\mathbf{f}(\phi^{-1}(y))\|_v d\mu_v,$$

for every open $W \subset U_x$. This local measure glues to a measure $\tilde{\omega}_{\mathcal{K},v}$ on $X(F_v)$. Indeed, changing the coordinates, ...

Let \mathcal{X} be a model of X over the integers \mathfrak{o}_F and let v be a place of good reduction. Let $\mathbb{F}_v = \mathfrak{o}_v/\mathfrak{m}_v$ be the corresponding finite field and put $q_v = \#\mathbb{F}_v$. Since X is projective, we have

$$X(F_v) = \mathcal{X}(\mathfrak{o}_v) \rightarrow \mathcal{X}(\mathbb{F}_v).$$

We have

$$\begin{aligned} \int_{X(F_v)} \tilde{\omega}_{\mathcal{K},v} &= \sum_{\bar{x}_v \in X(\mathbb{F}_v)} \int \tilde{\omega}_{\mathcal{K},v} \\ &= \frac{X(\mathbb{F}_v)}{q_v^d} \\ &= 1 + \frac{\mathrm{Tr}_v(\mathrm{H}_{et}^{2d-1}(X_{\mathbb{F}_v}))}{\sqrt{q_v}} + \frac{\mathrm{Tr}_v(\mathrm{H}_{et}^{2d-2}(X_{\mathbb{F}_v}))}{q_v} + \cdots + \frac{1}{q_v^d}, \end{aligned}$$

where Tr_v is the trace of the v -Frobenius on the ℓ -adic cohomology of X . Trying to integrate the product measure over $X(\mathbb{A})$ is problematic, since the Euler product

$$\prod_v \frac{X(\mathbb{F}_v)}{q_v^d}$$

diverges. In all examples of interest to us, the cohomology group $\mathrm{H}_{et}^{2d-1}(X_{\mathbb{F}_v}, \mathbb{Q}_\ell)$ vanishes. For instance, this holds if the anticanonical class is ample, by ???. Still the product diverges, since the $1/q_v$ term does not vanish, for projective X . There is a standard regularization procedure: Choose a finite set $S \subset \mathrm{Val}(F)$, including all $v \mid \infty$ and all places of bad reduction. Put

$$\lambda_v = \begin{cases} L_v(1, \mathrm{Pic}(X_{\mathbb{Q}})) & v \notin S \\ 1 & v \in S \end{cases},$$

where $L_v(s, \mathrm{Pic}(X_{\mathbb{Q}}))$ is the local factor of the Artin L -function associated to the Galois representation on the geometric Picard group. Define the regularized *Tamagawa measure*

$$\omega_{\mathcal{K},v} := \lambda_v^{-1} \tilde{\omega}_{\mathcal{K},v}.$$

Write $\omega_{\mathcal{K},S} := \prod_v \omega_{\mathcal{K},v}$ and define

eqn:tamagawa

$$(4.12) \quad \tau(-\mathcal{K}_X) := L_S^*(1, \mathrm{Pic}(X_{\mathbb{Q}})) \cdot \int_{X(F)} \omega_{\mathcal{K},S},$$

where

$$L_S^*(1, \mathrm{Pic}(X_{\mathbb{Q}})) := \lim_{s \rightarrow 1} (s-1)^r L_S^*(s, \mathrm{Pic}(X_{\mathbb{Q}}))$$

and r is the rank of $\mathrm{Pic}(X_F)$.

exam:group

Example 4.14.1. Let G be a linear algebraic group over F . It carries an F -rational d -form ω , where $d = \dim(G)$. This form is unique, modulo multiplication by nonzero constants. Fixing ω , we obtain an isomorphism $K_X \simeq \mathcal{O}_G$, the structure sheaf, which carries a natural adelic metrization $(\|\cdots\|_{\mathbb{A}})$.

$$\boxed{\text{eqn:c}} \quad (4.13) \quad c(-\mathcal{K}_X) = \alpha(X)\beta(X)\tau(-\mathcal{K}_X),$$

where

- $\alpha(X) := \alpha(\Lambda_{\text{eff}}(X), -K_X)$,
- $\beta(X) := \text{Br}(X)/\text{Br}(F)$,

and $\tau(-\mathcal{K}_X)$ is the constant defined in equation 4.12.

- $f = f(x_0, \dots, x_n) \in \mathbb{Z}[x_0, \dots, x_n]$, homogeneous, degree d
- $X = X_f \subset \mathbb{P}^n$ corresponding hypersurface
- $x \in X(\mathbb{Q}) \Leftrightarrow x = (x_0 : \dots : x_n) \in (\mathbb{Q}^{n+1} \setminus 0) / \pm$ with $f(x) = 0$
- choose a representative $x \in (\mathbb{Z}_{\text{prim}}^{n+1} \setminus 0) / \pm$ and put

$$H(x) := \max_j (|x_j|).$$

- define the counting function

$$N(X, B) := \#\{x \mid f(x) = 0, x \text{ primitive}, H(x) \leq B\}$$

ect:general-polarize

4.15. General polarizations. I follow closely the exposition in [BT98a]. Let E/F be some finite Galois extension such that all of the following constructions are defined over E . Let (X, \mathcal{L}) be a smooth *quasi-projective* d -dimensional variety together with a metrized very ample line bundle \mathcal{L} which embeds X in some projective space \mathbb{P}^n . We denote by $\overline{X}^{\mathcal{L}}$ the normalization of the projective closure of $X \subset \mathbb{P}^n$. In general, $\overline{X}^{\mathcal{L}}$ is singular. We will introduce several notions relying on a resolution of singularities

$$\rho : X \rightarrow \overline{X}^{\mathcal{L}}.$$

Naturally, the defined objects will be independent of the choice of the resolution.

For $\Lambda \subset \text{NS}(X)_{\mathbb{R}}$ we define

$$a(\Lambda, \mathcal{L}) := a(\Lambda, \rho^* \mathcal{L}).$$

We will always assume that $a(\Lambda_{\text{eff}}(X), \mathcal{L}) > 0$.

defn:primitive

Definition 4.15.1. A pair (X, \mathcal{L}) is called *primitive* if $a(\Lambda_{\text{eff}}(X), \mathcal{L}) \in \mathbb{Q}_{>0}$ and if there exists a resolution of singularities

$$\rho : X \rightarrow \overline{X}^{\mathcal{L}}$$

such that for some $k \in \mathbb{N}$

$$((\rho^* \mathcal{L})^{\otimes a(\Lambda_{\text{eff}}(X), \mathcal{L})} \otimes K_X)^{\otimes k} = \mathcal{O}(D),$$

where D is a *rigid* effective divisor ($h^0(X, \mathcal{O}(\nu D)) = 1$ for all $\nu \gg 0$).

Example 4.15.2. of a primitive pair: $(X, -\mathcal{K}_X)$, where X is a smooth projective Fano variety and $-\mathcal{K}_X$ is a metrized anticanonical line bundle.

Let $k \in \mathbb{N}$ be such that $a(\Lambda, \mathcal{L})k \in \mathbb{N}$ and consider

$$R(\Lambda, \mathcal{L}) := \bigoplus_{\nu \geq 0} \mathbf{H}^0(X, (((\rho^* \mathcal{L})^{a(\Lambda, \mathcal{L})} \otimes K_X)^{\otimes k})^{\otimes \nu}).$$

As explained in Section ??, in both cases ($\Lambda = \Lambda_{\text{ample}}$ or $\Lambda = \Lambda_{\text{eff}}$) it is expected that $R(\Lambda, \mathcal{L})$ is finitely generated and that we have a fibration

$$\pi = \pi_{\mathcal{L}} : X \rightarrow Y^{\mathcal{L}},$$

where $Y^{\mathcal{L}} = \text{Proj}(R(\mathcal{L}, \Lambda))$. For $\Lambda = \Lambda_{\text{eff}}(X)$ the generic fiber of π is (expected to be) a primitive variety in the sense of Definition 4.15.1. More precisely, there should be a diagram:

$$\begin{array}{ccc} \rho : X & \rightarrow & \overline{X}^{\mathcal{L}} \supset X \\ & & \downarrow \\ & & Y^{\mathcal{L}} \end{array}$$

such that:

- $\dim(Y^{\mathcal{L}}) < \dim(X)$;
- there exists a Zariski open $U \subset Y^{\mathcal{L}}$ such that for all $y \in U(\mathbb{C})$ the pair (X_y, \mathcal{L}_y) is primitive (here $X_y = \pi^{-1}(y) \cap X$ and \mathcal{L}_y is the restriction of \mathcal{L} to X_y);
- for all $y \in U(\mathbb{C})$ we have $a(\Lambda_{\text{eff}}(X), \mathcal{L}) = a(\Lambda_{\text{eff}}(X_y), \mathcal{L}_y)$;
- For all $k \in \mathbb{N}$ such that $a(\Lambda_{\text{eff}}(X), \mathcal{L})k \in \mathbb{N}$ the vector bundle

$$\mathcal{L}_k := R^0 \pi_* (((\rho^* \mathcal{L})^{\otimes a(\Lambda_{\text{eff}}(X), \mathcal{L})} \otimes K_X)^{\otimes k})$$

is in fact an ample invertible sheaf on $Y^{\mathcal{L}}$.

Such a fibration will be called an \mathcal{L} -primitive fibration. A variety may admit several primitive fibrations.

Example 4.15.3. Let $X \subset \mathbb{P}_1^n \times \mathbb{P}_2^n$ ($n \geq 2$) be a hypersurface given by a bi-homogeneous form of bi-degree (d_1, d_2) . Both projections $X \rightarrow \mathbb{P}_1^n$ and $X \rightarrow \mathbb{P}_2^n$ are \mathcal{L} -primitive, for appropriate \mathcal{L} . In particular, for $n = 3$ and $(d_1, d_2) = (1, 3)$ there are *two* distinct $-\mathcal{K}_X$ -primitive fibrations: one onto a point and another onto \mathbb{P}_1^3 .

sect:tama

4.16. Tamagawa numbers. For smooth projective Fano varieties X with an adelicly metrized anticanonical line bundle Peyre defined in [Pey95] a Tamagawa number, generalizing the classical construction for linear algebraic groups. We need to further generalize this to primitive pairs.

Abbreviate $a(\mathcal{L}) = a(\Lambda_{\text{eff}}(X), \mathcal{L})$ and let (X, \mathcal{L}) be a primitive pair such that

$$\mathcal{O}(D) := ((\rho^* \mathcal{L})^{\otimes a(\mathcal{L})} \otimes K_X)^{\otimes k},$$

where k is such that $a(\mathcal{L})k \in \mathbb{N}$ and D is a rigid effective divisor as in Definition 4.15.1. Choose an F -rational section $g \in H^0(X, \mathcal{O}(D))$; it is unique up to multiplication by F^* . Choose local analytic coordinates $x_{1,v}, \dots, x_{d,v}$ in a neighborhood U_x of $x \in X(F_v)$. In U_x the section g has a representation

$$g = f^{ka(\mathcal{L})} (dx_{1,v} \wedge \dots \wedge dx_{d,v})^k,$$

where f is a local section of L . This defines a local v -adic measure in U_x by

$$\omega_{\mathcal{L},g,v} := \|f\|_{x_v}^{a(\mathcal{L})} dx_{1,v} \cdots dx_{d,v},$$

where $dx_{1,v} \cdots dx_{d,v}$ is the Haar measure on F_v^d normalized by $\text{vol}(\mathfrak{o}_v^d) = 1$. A standard argument shows that $\omega_{\mathcal{L},g,v}$ glues to a v -adic measure on $X(F_v)$. The restriction of this measure to $X(F_v)$ does not depend on the choice of the resolution $\rho : X \rightarrow \overline{X}^{\mathcal{L}}$. Thus we have a measure on $X(F_v)$.

Denote by $(D_j)_{j \in \mathcal{J}}$ the irreducible components of the support of D and by

$$\text{Pic}(X, \mathcal{L}) := \text{Pic}(X \setminus \cup_{j \in \mathcal{J}} D_j).$$

The Galois group Γ acts on $\text{Pic}(X, \mathcal{L})$. Let S be a finite set of valuations of bad reduction for the data $(\rho, D_j, \text{etc.})$, including the archimedean valuations. Put $\lambda_v = 1$ for $v \in S$, $\lambda_v = L_v(1, \text{Pic}(X, \mathcal{L}))$ (for $v \notin S$) and

$$\omega_{\mathcal{L}} := L_S^*(1, \text{Pic}(X, \mathcal{L})) |\text{disc}(F)|^{-d/2} \prod_v \lambda_v^{-1} \omega_{\mathcal{L},g,v}.$$

(Here L_v is the local factor of the Artin L-function associated to the Γ -module $\text{Pic}(X, \mathcal{L})$ and $L_S^*(1, \text{Pic}(X, \mathcal{L}))$ is the residue at $s = 1$ of the partial Artin L-function.) By the product formula, the measure does

not depend on the choice of the F -rational section g . Define

$$\tau_{\mathcal{L}}(X) := \int_{\overline{X(F)}} \omega_{\mathcal{L}},$$

where $\overline{X(F)} \subset X(\mathbb{A})$ is the closure of $X(F)$ in the direct product topology. The convergence of the Euler product follows from

$$h^1(X, \mathcal{O}_X) = h^2(X, \mathcal{O}_X) = 0.$$

We have a map

$$\tilde{\rho} : \text{Pic}(X)_{\mathbb{R}} \rightarrow \text{Pic}(X, \mathcal{L})_{\mathbb{R}}$$

and we denote by

$$\Lambda_{\text{eff}}(X, \mathcal{L}) := \tilde{\rho}(\Lambda_{\text{eff}}(X)) \subset \text{Pic}(X, \mathcal{L})_{\mathbb{R}}.$$

Let (A, Λ) be a pair consisting of a lattice and a strictly convex (closed) cone in $A_{\mathbb{R}}$: $\Lambda \cap -\Lambda = 0$. Let $(\check{A}, \check{\Lambda})$ the pair consisting of the dual lattice and the dual cone defined by

$$\check{\Lambda} := \{\check{\lambda} \in \check{A}_{\mathbb{R}} \mid \langle \lambda', \check{\lambda} \rangle \geq 0, \forall \lambda' \in \Lambda\}.$$

The lattice \check{A} determines the normalization of the Lebesgue measure $d\check{a}$ on $\check{A}_{\mathbb{R}}$ (covolume = 1). For $a \in A_{\mathbb{C}}$ define

$$\boxed{\text{eqn:gam}} \quad (4.14) \quad \mathcal{X}_{\Lambda}(a) := \int_{\check{\Lambda}} e^{-\langle a, \check{a} \rangle} d\check{a}.$$

The integral converges absolutely and uniformly for $\Re(a)$ in compacts contained in the interior Λ° of Λ .

$\boxed{\text{defn:chi}}$ **Definition 4.16.1.** Assume that X is smooth, $\text{NS}(X) = \text{Pic}(X)$ and that $-K_X$ is in the interior of $\Lambda_{\text{eff}}(X)$. We define

$$\alpha(X) := \mathcal{X}_{\Lambda_{\text{eff}}(X)}(-K_X).$$

Remark 4.16.2. This constant measures the volume of the polytope obtained by intersecting the affine hyperplane $(-K_X, \cdot) = 1$ with the *dual* to the cone of effective divisors $\Lambda_{\text{eff}}(X)$ in the dual to the Néron-Severi group. The explicit determination of $\alpha(X)$ can be a serious problem. For Del Pezzo surfaces, these volumes are given in Section 1.9. For example, let X be the moduli space $\bar{M}_{0,6}$ (see Example ??). The dual to the cone $\Lambda_{\text{eff}}(X)$ has 3905 generators (in a 16-dimensional vector space), forming 25 orbits under the action of the symmetric group \mathbb{S}_6 .

$\boxed{\text{defn:tau}}$ **Definition 4.16.3.** Let (X, \mathcal{L}) be a primitive pair as above. Define

$$c(X, \mathcal{L}) := \mathcal{X}_{\Lambda_{\text{eff}}(X, \mathcal{L})}(\tilde{\rho}(-K_X)) \cdot |\text{H}^1(\Gamma, \text{Pic}(X, \mathcal{L}))| \cdot \tau_{\mathcal{L}}(X).$$

Example 4.16.4. Let us return to the Example ???. For the image of $\bar{\mathcal{M}}_{0,6}$ under f_3 (the Segre cubic) one knows an upper and lower bound:

$$c' \mathbf{B}^2 \log(\mathbf{B})^5 \leq \mathbf{N}_U(L_3, \mathbf{B}) \leq c'' \mathbf{B}^2 \log(\mathbf{B})^5,$$

for an appropriate Zariski open U and some constants $c', c'' \geq 0$ [VW95]. An *asymptotic* formula for $\mathbf{N}_U(L_3, \mathbf{B})$ of this shape would be compatible with the description in Theorem ???. Moreover, we are now in the position to specify a constant c which should appear in this asymptotic (answering a question in [VW95]). Indeed, the Segre cubic threefold is singular (it has 10 isolated double points). The blow-up $X = \tilde{S}_3$ of 5 points on \mathbb{P}^3 is a desingularization of S_3 . Its Picard group $\text{Pic}(X)$ is freely generated by the classes L, E_j (for $j = 1, \dots, 5$). The effective cone is generated by the classes $E_j, L - (E_i + E_j + E_k)$ and the anticanonical class is given by

$$-K_X = 4L - 2(E_1 + \dots + E_5).$$

The line bundle on S_3 giving the Segre embedding pulls back to $L = -1/2 \cdot K_X$. Clearly, $a(L) = 2$ and $b(L) = \text{rk Pic}(X) = 6$ (see also [BT98b], Section 5.2). The predicted leading constant $c = \alpha \cdot \tau$, where

$$\alpha = \frac{2^5}{5!} \mathcal{X}_\Lambda(-K_X)$$

and

$$\tau = \tau_\infty \cdot \prod_p (1 - 1/p)^6 (1 + 6/p + 6/p^2 + 1/p^3),$$

(τ_∞ is the ‘singular integral’ - the archimedean density of X).

If (X, \mathcal{L}) is not primitive then, by Section ??, some Zariski open subset $U \subset X$ admits a primitive fibration: there is a diagram

$$\begin{array}{ccc} X & \rightarrow & \bar{X}^{\mathcal{L}} \\ \downarrow & & \\ Y^{\mathcal{L}} & & \end{array}$$

such that for all $y \in Y^{\mathcal{L}}(F)$ the pair (U_y, \mathcal{L}_y) is primitive. Then

$$c(U, \mathcal{L}) := \sum_{y \in Y^0} c(U_y, \mathcal{L}_y),$$

where the right side is a (possibly infinite converging (!)) sum over the subset $Y^0 \subset Y^{\mathcal{L}}(F)$ of all those fibers U_y where

$$a(\mathcal{L}) = a(\mathcal{L}_y) \quad \text{and} \quad \text{rk Pic}(X, \mathcal{L})^{\mathbb{G}_a} = \text{rk Pic}(X_y, \mathcal{L}_y)^{\mathbb{G}_a}.$$

In Section 4.1 we will see that even if we start with pairs (X, \mathcal{L}) where X is a smooth projective variety and \mathcal{L} is a very ample adelically metrized line bundle on X we still need to consider singular varieties.

sect:tam-height

4.17. Tamagawa number as a height. For $\mathbf{a} \in \mathbb{P}(\mathbb{Q})$, let $S_{\mathbf{a}} \subset \mathbb{P}^3$ be the diagonal cubic surface fibration

eqn:aa (4.15)
$$a_0x_0^3 + a_1x_1^3 + a_2x_2^3 + a_3x_3^3 = 0,$$

considered in Section 4.13. Let $H : \mathbb{P}^3(\mathbb{Q}) \rightarrow \mathbb{R}_{>0}$ be the standard height as in Section 4.1.

exam:cub-tam

Theorem 4.17.1. [EJ] *For all $\epsilon > 0$ there exists a constant $c = c(\epsilon)$ such that*

$$\frac{1}{\tau(S_{\mathbf{a}})} \geq cH\left(\frac{1}{a_0} : \dots : \frac{1}{a_3}\right)^{1/3-\epsilon}$$

In particular, we have the following fundamental finiteness property: for $B > 0$ there are finitely many $\mathbf{a} \in \mathbb{P}^3(\mathbb{Q})$ such that $\tau(S_{\mathbf{a}}) > B$.

A similar, but unconditional, result holds for 3 dimensional cubics and quartics

Theorem 4.17.2.

sect:smallest

4.18. Smallest points. [Sie69]

For a discussion of bounds of diophantine equations in terms of the *height* of the equation, see [Mas02].

[NP89]

A sample result in this direction is [Pit71], [NP89]: Let

eqn:diag (4.16)
$$\sum_{i=0}^n a_i x_i^d = 0,$$

with d odd and let $\mathbf{a} = (a_0 \text{ ldots } a_n) \in \mathbb{Z}^{n+1}$ be a vector with nonzero coordinates. For $n \gg d$ (e.g., $n = 2^d + 1$) and any $\epsilon > 0$ there exists a constant c such that (4.16) has a solution \mathbf{x} with

$$\sum_{i=0}^n |a_i x_i^d| < c \prod |a_i|^{d+\epsilon}.$$

For $d \geq 12$, one can work with $n \sim 4d^2 \log(d)$. There have been a several improvements of this result for specific values of d , e.g. [Cas55], [Die03] for quadrics and [Bak89], [Brü94] for $d = 3$.

thm:smallest-height

Theorem 4.18.1. [EJ] *Let $S_a \subset \mathbb{P}^3$ be the cubic surface given by*

$$ax_0^3 + 4x_1^3 + 2x_2^3 + x_3^3 = 0.$$

Assume the Generalized Riemann Hypothesis. Then there does not exist a constant $c > 0$ such that

$$\min \{H(\mathbf{x}) \mid \mathbf{x} \in S_a(\mathbb{Q})\} \leq \frac{c}{\tau(S_a)},$$

for all $a \in \mathbb{Z}$.

5. COUNTING POINTS VIA UNIVERSAL TORSORS

sect:count-torsors

5.1. The formalism. [Pey04], [Sal98]

sect:toric-torsors

5.2. Toric Del Pezzo surfaces.

Example 5.2.1. Let $X = \text{Bl}_Y(P^2)$ be the blowup of the projective plane in the subscheme

$$Y := (1 : 0 : 0) \cup (0 : 1 : 0) \cup (0 : 0 : 1),$$

a toric Del Pezzo surface of degree 6. We can realize it as a subvariety $X \subset \mathbb{P}_{\mathbf{x}}^1 \times \mathbb{P}_{\mathbf{y}}^1 \times \mathbb{P}_{\mathbf{z}}^1$ given by $x_0y_0z_0 = x_1y_1z_1$. The anticanonical height is given by

$$\max(|x_0|, |x_1|) \times \max(|y_0|, |y_1|) \times \max(|z_0|, |z_1|).$$

There are six exceptional curves: the preimages of the 3 points and the preimages of lines joining two of these points.

$$xyz = t^3 \dots x^2yz = t^4$$

5.3. Torsors over Del Pezzo surfaces.

Example 5.3.1. Quintic DP over \mathbb{Q} and Grassmannian. [dlB02]

sect:quartic-dp

Example 5.3.2. A quartic Del Pezzo surface S with two singularities of type A_1 can be realized as a blow-up of the following points

$$\begin{aligned} p_1 &= (0 : 0 : 1) \\ p_2 &= (1 : 0 : 0) \\ p_3 &= (0 : 1 : 0) \\ p_4 &= (1 : 0 : 1) \\ p_5 &= (0 : 1 : 1) \end{aligned}$$

in $\mathbb{P}^2 = (x_0 : x_1 : x_2)$. The anticanonical line bundle embeds S into \mathbb{P}^4 :

$$(x_0^2x_1 : x_0x_1^2 : x_0x_1x_2 : x_0x_2(x_0 + x_1 - x_2) : x_1x_2(x_0 + x_1 - x_2)).$$

The Picard group is spanned by

$$\text{Pic}(S) = \langle L, E_1, \dots, E_5 \rangle$$

and $\Lambda_{\text{eff}}(S)$ by

$$\begin{aligned} & E_1, \dots, E_5 \\ L - E_2 - E_3, L - E_3 - E_4, L - E_4 - E_5, L - E_2 - E_5 \\ & L - E_1 - E_3 - E_5, L - E_1 - E_2 - E_4. \end{aligned}$$

The universal torsor is given by the following equations

$$\begin{aligned} (23)(3) - (1)(124)(4) + (25)(5) &= 0 \\ (23)(2) - (1)(135)(5) + (34)(4) &= 0 \\ (124)(1)(2) - (34)(3) + (45)(5) &= 0 \\ (25)(2) - (135)(1)(3) + (45)(4) &= 0 \\ (23)(45) + (34)(25) - (1)^2(124)(135) &= 0. \end{aligned}$$

(with variables labeled by the corresponding exceptional curves). Introducing additional variables

$$(24)' := (1)(124), \quad (35)' := (1)(135)$$

we see that the above equations define a \mathbb{P}^1 -bundle over a codimension one subvariety of the Grassmannian $\text{Gr}(2, 5)$.

We need to estimate the number of 11-tuples of nonzero integers, satisfying the equations above and subject to the inequalities:

$$\begin{aligned} |(135)(124)(23)(1)(2)(3)| &\leq B \\ |(135)(124)(34)(1)(3)(4)| &\leq B \\ &\dots \end{aligned}$$

By symmetry, we can assume that $|(2)| \geq |(4)|$ and write $(2) = (2)'(4) + r_2$. Now we weaken the first inequality to

$$|(135)(124)(23)(1)(4)(2)'(3)| \leq B.$$

There are $O(B \log(B)^6)$ 7-tuples of integers satisfying this inequality.

Step 1. Use equation $(23)(3) - (1)(124)(4) + (25)(5)$ to reconstruct $(25), (5)$ with ambiguity $O(\log(B))$.

Step 2. Use $(25)(2) - (135)(1)(3) + (45)(4) = 0$ to reconstruct the residue r_2 modulo (4) . Notice that (25) and (4) are “almost” coprime since the corresponding exceptional curves are disjoint.

Step 3. Reconstruct (2) and (45) .

Step 4. Use $(23)(2) - (1)(135)(5) + (34)(4)$ to reconstruct (34).

In conclusion, if $U \subset S$ is the complement to exceptional curves then

$$\mathbf{N}_U(-K_S, \mathbf{B}) = O(\mathbf{B} \log(\mathbf{B})^7).$$

As explained in Remark ??, we expect that

$$\mathbf{N}_U(-K_S, \mathbf{B}) = \mathbf{B} \log(\mathbf{B})^5(1 + o(1))$$

as $B \rightarrow \infty$, where c is the constant defined in Chapter ??.

`exem:dp4-tors`

Example 5.3.3. The universal torsor of a smooth quartic Del Pezzo surface, given as a blow up of the five points

$$\begin{aligned} p_1 &= (1 : 0 : 0) \\ p_2 &= (0 : 1 : 0) \\ p_3 &= (0 : 0 : 1) \\ p_4 &= (1 : 1 : 1) \\ p_5 &= (1 : a_2 : a_3), \end{aligned}$$

assumed to be in general position, is given by the vanishing of

$(14)(23)$	+	$(12)(34)$	-	$(13)(24)$		$(00)(05) - (12)(34) + (13)(24) - (14)(23)$
$(00)(05)$	+	$a_3(a_2 - 1)(12)(34)$	-	$a_2(a_3 - 1)(13)(24)$		
$(23)(03)$	+	$(24)(04)$	-	$(12)(01)$		$(12)(01) - (23)(03) + (24)(04) - (25)(05)$
$a_2(23)(03)$	+	$(25)(05)$	-	$(12)(01)$		
$(12)(35)$	-	$(13)(25)$	+	$(15)(23)$		$(00)(04) - (12)(35) + (13)(25) - (15)(23)$
$(a_2 - 1)(12)(35)$	+	$(00)(04)$	-	$(a_3 - 1)(13)(25)$		
$(12)(45)$	+	$(14)(25)$	-	$(15)(24)$		$(00)(03) - (12)(45) + (14)(25) - (15)(24)$
$(00)(03)$	+	$a_3(14)(25)$	-	$(15)(24)$		
$(13)(45)$	+	$(14)(35)$	-	$(15)(34)$		$(00)(02) - (13)(45) + (14)(35) - (15)(34)$
$(00)(02)$	+	$a_2(14)(35)$	-	$(15)(34)$		
$(23)(45)$	+	$(24)(35)$	-	$(25)(34)$		$(00)(01) - (23)(45) + (24)(35) - (25)(34)$
$(00)(01)$	+	$a_2(24)(35)$	-	$a_3(25)(34)$		
$(04)(34)$	+	$(02)(23)$	-	$(01)(13)$		$(13)(01) - (23)(02) + (34)(04) + (35)(05)$
$(05)(35)$	+	$a_3(02)(23)$	-	$(01)(13)$		
$(a_2 - 1)(03)(34)$	+	$(05)(45)$	-	$(a_3 - 1)(02)(24)$		$(14)(01) - (24)(02) + (34)(03) - (45)(05)$
$(03)(34)$	+	$(01)(14)$	-	$(02)(24)$		
$(04)(14)$	+	$(03)(13)$	-	$(02)(12)$		$(12)(02) - (13)(03) + (14)(04) - (15)(05)$
$(05)(15)$	+	$a_2(03)(13)$	-	$a_3(02)(12)$		
$a_3(02)(25)$	-	$a_2(03)(35)$	-	$(01)(15)$		$(15)(01) - (25)(02) + (35)(03) - (45)(04)$
$(a_3 - 1)(02)(25)$	-	$(a_2 - 1)(03)(35)$	-	$(04)(45)$		

Connection to the D_5 -Grassmannian

exam:cay-tors

Example 5.3.4. The Cayley cubic X_3 is the unique cubic hypersurface in \mathbb{P}^3 with 4 double points (A_1 -singularities), the maximal number of double points on a cubic surface. It can be given by the equation

$$y_0y_1y_2 + y_0y_1y_3 + y_0y_2y_3 + y_1y_2y_3 = 0.$$

The double points correspond to

$$(1 : 0 : 0 : 0), (0 : 1 : 0 : 0), (0 : 0 : 1 : 0), (0 : 0 : 0 : 1).$$

It can be realized as the blow-up of $\mathbb{P}^2 = (x_1 : x_2 : x_3)$ in the points

$$q_1 = (1 : 0 : 0), q_2 = (0 : 1 : 0), q_3 = (0 : 0 : 1), q_4 = (1 : -1 : 0), q_5 = (1 : 0 : -1), q_6 = (0 : 1 : -1)$$

The points lie on a rigid configuration of 7 lines

$$\begin{array}{ll} x_1 = 0 & (12)(13)(14)(1) \\ x_2 = 0 & (12)(23)(24)(2) \\ x_3 = 0 & (13)(23)(34)(3) \\ x_4 = x_1 + x_2 + x_3 & (14)(24)(34)(4) \\ x_1 + x_3 = 0 & (13)(24)(13, 24) \\ x_2 + x_3 = 0 & (23)(14)(14, 23) \\ x_1 + x_2 = 0 & (12)(34)(12, 34). \end{array}$$

The proper transform of the line x_j is the (-2) -curve corresponding to (j) . The curves corresponding to (ij) , (ij, kl) are (-1) -curves. The accumulating subvarieties are exceptional curves. The (anticanonical) embedding $X_3 \hookrightarrow \mathbb{P}^3$ is given by the linear system:

$$\begin{array}{ll} s_1 & = x_1x_2x_3 \\ s_2 & = x_2x_3x_4 \\ s_3 & = x_1x_3x_4 \\ s_4 & = x_1x_2x_4 \end{array}$$

The counting problem is: estimate

$$N(\mathbf{B}) = \#\{(x_1, x_2, x_3) \in \mathbb{Z}_{\text{prim}}^3 / \pm, \mid \max i(|s_i|) / \gcd(s_i) \leq \mathbf{B}\},$$

where the triple x_j is subject to the conditions

$$x_i \neq 0, (i = 1, \dots, 4) \quad x_j + x_i \neq 0 (1 \leq i < j \leq 3).$$

We expect $\sim \mathbf{B} \log(\mathbf{B})^6$ solutions. After dividing by the coordinates by their gcd, we obtain

$$\begin{array}{ll} s'_1 & = (1)(2)(3)(12)(13)(23) \\ s'_2 & = (2)(3)(4)(23)(24)(34) \\ s'_3 & = (1)(3)(4)(13)(14)(34) \\ s'_4 & = (1)(2)(4)(12)(14)(24) \end{array}$$

(These are special sections in the anticanonical series, other decomposable sections are: $(1)(2)(12)^2(12, 34)$ and $(12, 34)(13, 24)(14, 23)$, for

example.) The conic bundles on X_3 produce the following equations for the universal torsor:

$$\begin{array}{llll}
 \text{I} & (1)(13)(14) & + & (2)(23)(24) = (34)(12, 34) \\
 \text{II} & (1)(12)(14) & + & (3)(23)(34) = (24)(13, 24) \\
 \text{III} & (2)(12)(24) & + & (3)(13)(34) = (14)(14, 23) \\
 \text{IV} & -(3)(13)(23) & + & (4)(14)(24) = (12)(12, 34) \\
 \text{V} & -(2)(12)(23) & + & (4)(14)(34) = (13)(13, 24) \\
 \text{VI} & -(1)(12)(1) & + & (4)(24)(34) = (23)(14, 23) \\
 \text{VII} & (2)(4)(24)^2 & + & (1)(3)(13)^2 = (12, 34)(14, 23) \\
 \text{VIII} & -(1)(2)(12)^2 & + & (3)(4)(34)^2 = (13, 24)(14, 23) \\
 \text{IX} & (1)(4)(14)^2 & - & (2)(3)(23)^2 = (12, 34)(13, 24)
 \end{array}$$

The counting problem is to estimate the number of 13-tuples of *nonzero* integers, satisfying the equations above and subject to the inequality $\max_i \{|s'_i|\} \leq B$. Heath-Brown proved in [HB03] that there exists constants $0 < c < c'$ such that

$$cB \log(B)^6 \leq N(B) \leq c'B \log(B)^6.$$

sect:tors-segre

5.4. Torsors over the Segre cubic. In this section we work over \mathbb{Q} . The threefold $X = \bar{M}_{0,6}$ can be realized as the blow-up of \mathbb{P}^3 in the points

$$\begin{array}{cccc}
 & x_0 & x_1 & x_2 & x_3 \\
 q_1 & 1 & 0 & 0 & 0 \\
 q_2 & 0 & 1 & 0 & 0 \\
 q_3 & 0 & 0 & 1 & 0 \\
 q_4 & 0 & 0 & 0 & 1 \\
 q_5 & 1 & 1 & 1 & 1
 \end{array}$$

and in the proper transforms of lines joining two of these points. The Segre cubic is given as the image of X in \mathbb{P}^4 under the linear system $2L - (E_1 + \dots + E_5)$ (quadrics passing through the 5 points):

$$\begin{array}{ll}
 s_1 & = (x_2 - x_3)x_1 \\
 s_2 & = x_3(x_0 - x_1) \\
 s_3 & = x_0(x_1 - x_3) \\
 s_4 & = (x_0 - x_1)x_2 \\
 s_5 & = (x_1 - x_2)(x_0 - x_3)
 \end{array}$$

It can be realized in $\mathbb{P}^5 = (y_0 : \dots : y_5)$ as

$$\mathcal{S}_3 := \left\{ \sum_{i=0}^5 y_i^3 = \sum_{i=0}^5 y_i = 0 \right\}$$

(exhibiting the \mathfrak{S}_6 -symmetry.) It contains 15 planes, given by the \mathfrak{S}_6 -orbit of

$$y_0 + y_3 = y_1 + y_4 = y_2 + y_5 = 0,$$

and 10 singular double points, given by the \mathfrak{S}_6 -orbit of

$$(1 : 1 : 1 : -1 : -1 : -1).$$

This is the maximal number of nodes on a cubic threefold and \mathcal{S}_3 is the unique cubic with this property. The hyperplane sections $\mathcal{S}_3 \cap \{y_i = 0\}$ are *Clebsch* diagonal cubic surfaces (unique cubic surfaces with \mathfrak{S}_5 as symmetry group). The hyperplane sections $\mathcal{S}_3 \cap \{y_i = 0\}$ are *Clebsch* cubics, a unique cubic surface with \mathfrak{S}_5 -symmetry. The hyperplane sections $\mathcal{S}_3 \cap \{y_i - y_j = 0\}$ are *Cayley* cubic surfaces (see Section ??). The geometry and symmetry of these and similar varieties are described in detail in [Hun96].

The counting problem on \mathcal{S}_3 is: find the number $N(\mathbf{B})$ of all 4-tupels of $(x_0, x_1, x_2, x_3) \in \mathbb{Z}^4/\pm$ such that

- $\gcd(x_0, x_1, x_2, x_3) = 1$;
- $\max_{j=1, \dots, 5}(|s_j|) / \gcd(s_1, \dots, s_5) \leq \mathbf{B}$;
- $x_i \neq 0$ and $x_i - x_j \neq 0$ for all $i, j \neq i$.

The last condition is excluding rational points contained in accumulating subvarieties (there are \mathbf{B}^3 rational points on planes $\mathbb{P}^2 \subset \mathbb{P}^4$, with respect to the $\mathcal{O}(1)$ -height). The second condition is the bound on the *height*.

First we need to determine

$$a(L) = \inf\{a \mid aL + K_X \in \Lambda_{\text{eff}}(X)\},$$

where L is the line bundle giving the map to \mathbb{P}^4 . We claim that $a(L) = 2$. This follows from the fact that

$$\sum_{i,j} (ij)$$

is on the boundary of $\Lambda_{\text{eff}}(X)$ (where (ij) is the class in $\text{Pic}(X)$ of the preimage in X of the line $l_{ij} \subset \mathbb{P}^4$ through q_i, q_j).

Therefore, we expect

$$N(\mathbf{B}) = O(\mathbf{B}^{2+\epsilon})$$

as $\mathbf{B} \rightarrow \infty$. In fact, it was shown in [BT98a] that $b(L) = 6$. Consequently, one expects

$$N(\mathbf{B}) \sim c\mathbf{B} \log(\mathbf{B})^5, \quad \text{as } \mathbf{B} \rightarrow \infty.$$

Remark 5.4.1. The difficult part is to keep track of $\gcd(s_1, \dots, s_5)$. Indeed, if we knew that this $\gcd = 1$ we could easily prove the bound $O(\mathbf{B}^{2+\epsilon})$ by observing that there are $O(\mathbf{B}^{1+\epsilon})$ pairs of (positive) integers $(x_2 - x_3, x_1)$ (resp. $(x_0 - x_1, x_2)$) satisfying $(x_2 - x_3)x_1 \leq \mathbf{B}$ (resp. $(x_0 - x_1)x_2 \leq \mathbf{B}$). Then we could reconstruct the quadruple

$$(x_2 - x_3, x_1, x_0 - x_1, x_2)$$

and consequently

$$(x_0, x_1, x_2, x_3)$$

up to $O(\mathbf{B}^{2+\epsilon})$.

Thus it is necessary to introduce \gcd between x_j , etc. Again, we use the symbols (i) , (ij) , (ijk) for variables on the torsor for X corresponding to the classes of the preimages of points, lines, planes resp. Once we fix a point $(x_0, x_1, x_2, x_3) \in \mathbb{Z}^4$ (such that $\gcd(x_0, x_1, x_2, x_3) = 1$), the values of these coordinates over the corresponding point on X can be expressed as greatest common divisors. For example, we can write

$$x_3 = (123)(12)(13)(23)(1)(2)(3),$$

a product of integers (neglecting the sign of x_3 ; in the torsor language, we are looking at the orbit of $\mathbb{T}_{\text{NS}}(\mathbb{Z})$). Here is a self-explanatory list:

(123)	x_3	(12)	x_2, x_3
(124)	x_2	(13)	x_1, x_3
(125)	$x_2 - x_3$	(14)	x_1, x_2
(134)	x_1	(15)	$x_1 - x_3, x_1 - x_2$
(135)	$x_1 - x_3$	(23)	$x_3, x_0 - x_3$
(145)	$x_1 - x_2$	(24)	x_2, x_0
(234)	x_0	(25)	$x_3 - x_2, x_0 - x_3$
(235)	$x_0 - x_3$	(34)	x_1, x_0
(245)	$x_0 - x_2$	(35)	$x_1 - x_3, x_0 - x_1$
(345)	$x_0 - x_1$	(45)	$x_1 - x_2, x_0 - x_1$.

After dividing s_j by the \gcd , we get

$$\begin{aligned} s'_1 &= (125)(134)(12)(15)(25)(13)(14)(34)(1) \\ s'_2 &= (123)(245)(12)(13)(23)(24)(25)(45)(2) \\ s'_3 &= (234)(135)(23)(24)(34)(13)(15)(35)(3) \\ s'_4 &= (345)(124)(34)(35)(45)(12)(14)(24)(4) \\ s'_5 &= (145)(235)(14)(15)(45)(23)(35)(25)(5) \end{aligned}$$

(observe the symmetry with respect to the permutation (12345)). We claim that $\gcd(s'_1, \dots, s'_5) = 1$. One can check this directly using the

definitions of $(i), (ij), (ijk)$'s as gcd's. For example, let us check that nontrivial divisors $d \neq 1$ of (1) cannot divide any other s'_j . Such a d must divide (123) or (12) or (13) (see s'_2). Assume it divides (12). Then it doesn't divide (13), (14) and (15) (the corresponding divisors are disjoint). Therefore, d divides (135) (by s'_3) and (235) (by s'_5). Contradiction (indeed, (135) and (235) correspond to disjoint divisors). Assume that d divides (123). Then it has to divide either (13) or (15) (from s'_3) and either (12) or (14) (from s'_4). Contradiction.

The integers $(i), (ij), (ijk)$ satisfy a system of relations (these are equations for the torsor induced from fibrations of $\bar{M}_{0,6}$ over \mathbb{P}^1):

I	x_0	x_1	$x_0 - x_1$
II	x_0	x_2	$x_0 - x_2$
III	x_0	x_3	$x_0 - x_3$
IV	x_1	x_2	$x_1 - x_2$
V	x_1	x_3	$x_1 - x_3$
VI	x_2	x_3	$x_2 - x_3$
VII	$x_0 - x_1$	$x_0 - x_2$	$x_1 - x_2$
VIII	$x_0 - x_1$	$x_0 - x_3$	$x_1 - x_3$
IX	$x_1 - x_2$	$x_1 - x_3$	$x_2 - x_3$
X	$x_2 - x_3$	$x_0 - x_3$	$x_0 - x_2$

which translates to

I	$(234)(23)(24)(2) - (134)(13)(14)(1)$	$=$	$(345)(45)(35)(5)$
II	$(234)(23)(34)(3) - (124)(12)(14)(1)$	$=$	$(245)(25)(45)(5)$
III	$(234)(24)(34)(4) - (123)(12)(13)(1)$	$=$	$(235)(25)(35)(5)$
IV	$(134)(13)(34)(3) - (124)(12)(24)(2)$	$=$	$(145)(15)(45)(5)$
V	$(134)(14)(34)(4) - (123)(12)(23)(2)$	$=$	$(135)(15)(35)(5)$
VI	$(124)(14)(24)(4) - (123)(13)(23)(3)$	$=$	$(125)(15)(25)(5)$
VII	$(345)(34)(35)(3) - (245)(24)(25)(2)$	$=$	$-(145)(14)(15)(1)$
VIII	$(345)(34)(45)(4) - (235)(23)(25)(2)$	$=$	$-(135)(13)(15)(1)$
IX	$(145)(14)(45)(4) - (135)(13)(35)(3)$	$=$	$-(125)(12)(25)(2)$
X	$(125)(12)(15)(1) + (235)(23)(35)(3)$	$=$	$-(245)(24)(45)(4)$

The counting problem now becomes: find all 25-tuples of *nonzero* integers satisfying the equations I–X and the inequality $\max(|s'_j|) \leq \mathbf{B}$.

rema:gr26

Remark 5.4.2. Note the analogy to the the case of $\bar{M}_{0,5}$ (the unique split Del Pezzo surface of degree 5): the variety defined by the above equations is the Grassmannian $\text{Gr}(2, 6)$ (in its Plücker embedding into \mathbb{P}^{24}).

Remark 5.4.3. Vaughan and Wooley show in [VW95] that there exist constants $c, c' > 0$ such that

$$c\mathbf{B} \log(\mathbf{B})^5 \leq \mathbf{N}(\mathbf{B}) \leq c'\mathbf{B} \log(\mathbf{B})^5.$$

They use a different (an intermediate) torsor over X - the determinantal variety given by

$$\det(x_{ij})_{3 \times 3} = 0.$$

sect:flag-tors

5.5. Flag varieties and torsors. Natural examples of torsors under tori arise from flag varieties. Let G be a semi-simple algebraic group, $P \subset G$ a parabolic subgroup. The flag variety $P \backslash G$ admits an action by any subtorus of the maximal torus in G on the right. Choosing a linearization for this action and passing to the quotient we obtain a plethora of examples of *nonhomogeneous* varieties X whose torsors carry additional symmetries. These may be helpful in the counting rational points on X . Indeed, we have already seen such examples (Del Pezzo surface of degree 5 ?? and the Segre cubic, Section 5.4).

In this section we expand on these examples.

exam:tors-g2

Example 5.5.1. A flag variety for the group G_2 is the quadric hypersurface

$$v_1u_1 + v_2u_2 + v_3u_3 + z^2 = 0,$$

where the torus $G_m^2 \subset G_2$ acts as

$$\begin{aligned} v_j &\mapsto \lambda_j v_j, & j = 1, 2 & \quad v_3 \mapsto (\lambda_1 \lambda_2)^{-1} v_3 \\ u_j &\mapsto \lambda_j^{-1} u_j, & j = 1, 2 & \quad u_3 \mapsto \lambda_1 \lambda_2 u_3. \end{aligned}$$

The quotient by G_m^2 is a subvariety in the weighted projective space $\mathbb{P}(1, 2, 2, 2, 3, 3) = (z : x_1 : x_2 : x_3 : y_1 : y_2)$ with the equations

$$x_0 + x_1 + x_2 + z^2 = 0 \quad \text{and} \quad x_1 x_2 x_3 = y_1 y_2.$$

ssettzeta

6. HEIGHT ZETA FUNCTIONS

6.1. Tools from analysis. In this section we collect technical results from complex and harmonic analysis which will be used in the treatment of height zeta functions.

For $U \subset \mathbb{R}^n$ let

$$\mathbb{T}_U := \{s \in \mathbb{C} \mid \Re(s) \in U\}$$

be the tube domain over U .

thm:convex

Theorem 6.1.1 (Convexity principle). *Let $U \subset \mathbb{R}^n$ be a connected open subset and \bar{U} the convex envelope of U , i.e., the smallest convex open set containing U . Let $Z(s)$ be a function holomorphic in \mathbb{T}_U . Then $Z(s)$ is holomorphic in $\mathbb{T}_{\bar{U}}$.*

thm:phragmen

Theorem 6.1.2 (Phragmen-Lindelöf principle). *Let ϕ be a holomorphic function for $\Re(s) \in [\sigma_1, \sigma_2]$. Assume that in this domain ϕ satisfies the following bounds*

- $|\phi(s)| = O(e^{\epsilon|t|})$, for all $\epsilon > 0$;
- $|\phi(\sigma_1 + it)| = O(|t|^{k_1})$ and $|\phi(\sigma_2 + it)| = O(|t|^{k_2})$.

Then, for all $\sigma \in [\sigma_1, \sigma_2]$ one has

$$|\phi(\sigma + it)| = O(|t|^k), \quad \text{where } \frac{k - k_1}{\sigma - \sigma_1} = \frac{k_2 - k_1}{\sigma_2 - \sigma_1}.$$

Using the functional equation and known bounds for $\Gamma(s)$ in vertical strips one derives the *convexity bound*

eqn:bound-zeta

$$(6.1) \quad \left| \zeta\left(\frac{1}{2} + it\right) \right| = O(|t|^{1/4+\epsilon}).$$

More generally, we have the following

prop:phragmen

Proposition 6.1.3. *Let χ be an unramified character of $\mathbb{G}_m(\mathbb{A}_F)/\mathbb{G}_m(F)$, i.e., χ_v is trivial on $\mathbb{G}(\mathfrak{o}_v)$, for all $v \nmid \infty$. For all $\epsilon > 0$ there exists a $\delta > 0$ such that*

eqn:l-estim

$$(6.2) \quad |L(s, \chi)| \ll (1 + |\mathfrak{S}(\chi)| + |\mathfrak{S}(s)|)^\epsilon, \quad \text{for } \Re(s) > 1 - \delta.$$

Here $\mathfrak{S}(\chi) \in \bigoplus_{v|\infty} \mathbb{G}_m(F_v)/\mathbb{G}_m(\mathfrak{o}_v) \simeq \mathbb{R}^{r_1+r_2-1}$, with r_1, r_2 the number of real, resp. pairs of complex embeddings of F .

thm:tauber

Theorem 6.1.4 (Tauberian theorem). *Let $\{\lambda_n\}$ be an increasing sequence of positive real numbers, with $\lim_{n \rightarrow \infty} \lambda_n = \infty$. Let $\{a_n\}$ be another sequence of positive real numbers and put*

$$Z(s) := \sum_{n \geq 1} \frac{a_n}{\lambda_n^s}.$$

Assume that this series converges absolutely and uniformly to a holomorphic function in the tube domain $\mathbb{T}_{>a} \subset \mathbb{C}$, for some $a > 0$, and that it admits a representation

$$Z(s) = \frac{h(s)}{(s-a)^b},$$

where h is holomorphic in $\mathbb{T}_{>a-\epsilon}$, for some $\epsilon > 0$, with $h(a) = c > 0$, and $b \in \mathbb{N}$. Then

$$N(B) := \sum_{\lambda_n \leq B} a_n \sim \frac{c}{a(b-1)!} B^a \log(B)^{b-1}, \quad \text{for } B \rightarrow \infty.$$

A frequently employed result in analytic number theory is

Theorem 6.1.5 (Poisson formula). *Let G be a locally compact abelian group with Haar measure dg and $H \subset G$ a closed subgroup. Let \hat{G} be the Pontryagin dual of G , i.e., the group of characters, i.e., continuous homomorphisms,*

$$\chi : G \rightarrow \mathbb{S}^1 \subset \mathbb{C}^*$$

into the unit circle. Let $f : G \rightarrow \mathbb{C}$ be a function, satisfying some mild assumptions (integrability, continuity) and let

$$\hat{f}(\chi) := \int_G f(g) \cdot \chi(g) dg$$

be its Fourier transform. Then there exist Haar measures dh on H and dh^\perp on H^\perp , the subgroup of characters trivial on H , such that

eqn:poisson

$$(6.3) \quad \int_H f dh = \int_{H^\perp} \hat{f} dh^\perp.$$

A standard application is to $H = \mathbb{Z} \subset \mathbb{R} = G$. In this case $H^\perp = H = \mathbb{Z}$, and the formula reads

$$\sum_{n \in \mathbb{Z}} f(n) = \sum_{n \in \mathbb{Z}} \hat{f}(n).$$

This is a powerful identity which is used, e.g., to prove the functional equation and meromorphic continuation of the Riemann zeta function. We will apply Equation (6.3) in the case when G is the group of adelic points of an algebraic torus or an additive group, and H is the subgroup of rational points. This will allow us to establish a meromorphic continuation of height zeta functions for equivariant compactifications of these groups.

Another application of the Poisson formula arises as follows: Let A be a lattice and Λ a convex cone in $A_\mathbb{R}$. Let $d\check{a}$ be the Lebesgue measure on the dual space $\check{A}_\mathbb{R}$ normalized by the dual lattice \check{A}). Let

$$\alpha(\Lambda, \mathbf{s}) := \int_{\check{\Lambda}} e^{-\langle \mathbf{s}, \check{a} \rangle} d\check{a}, \quad \Re(\mathbf{s}) \in \Lambda^\circ.$$

be the Laplace transform of the set-theoretic characteristic function of the dual cone $\check{\Lambda}$.

Let $\pi : (A, \Lambda) \rightarrow (\tilde{A}, \tilde{\Lambda})$ be a homomorphism, with finite cokernel A' and kernel $B \subset A$. Normalize $db : \text{vol}(B_\mathbb{R}/B) = 1$. Then

$$\alpha(\tilde{\Lambda}, \pi(\mathbf{s})) = \frac{1}{(2\pi)^{k-\tilde{k}}} \frac{1}{|A'|} \int_{B_\mathbb{R}} \alpha(\Lambda, \mathbf{s} + ib) db.$$

In particular,

$$\alpha(\Lambda, \mathbf{s}) = \frac{1}{(2\pi)^d} \int_{M_{\mathbb{R}}} \prod_{j=1}^n \frac{1}{(s_j + im_j)} dm.$$

Oh's results

some number theoretic estimates??

sect:setup-groups

6.2. Compactifications of groups and homogeneous spaces. As already mentioned in Section 3, an easy way to generate examples of algebraic varieties with many rational points is to use actions of algebraic groups. Here we discuss the geometric properties of groups and their compactifications.

Let G be a linear algebraic group over a field F , and

$$\varrho : G \rightarrow \mathrm{PGL}_{n+1}$$

an algebraic representation over F . Let $x \in \mathbb{P}^n(F)$ be a point. The orbit $\varrho(G) \cdot x \subset \mathbb{P}^n$ inherits rational points from $G(F)$. Let $H \subset G$ be the stabilizer of x . In general, we have an exact sequence

$$1 \rightarrow H(F) \rightarrow G(F) \rightarrow G/H(F) \rightarrow H^1(F, H) \rightarrow \dots$$

We will only consider examples when $(G/H)(F) = G(F)/H(F)$.

By construction, the Zariski closure X of $\varrho(G) \cdot x$ is geometrically isomorphic to an *equivariant* compactification of the homogeneous space G/H . We have a dictionary

$$(\varrho, x \in \mathbb{P}^n) \Leftrightarrow \begin{cases} \text{equivariant compactification } X \supset G/H, \\ G\text{-linearized very ample line bundle } L \text{ on } X. \end{cases}$$

Representations of semisimple groups do not deform, and can be characterized by combinatorial data: lattices, polytopes etc. Note, however, that the choice of the initial point $x \in \mathbb{P}^n$ can still give rise to moduli. On the other hand, the classification of representations of unipotent groups is a *wild* problem, already for $G = \mathbb{G}_a^2$. In this case, understanding the moduli of representations of a fixed dimension is equivalent to classifying pairs of commuting matrices, up to conjugacy (see [GP69]).

recent Gorodnik/Oh

Borovoi

sect:principles

6.3. Basic principles. Here we explain some common features in the study of height zeta functions of compactifications of groups and homogeneous spaces.

In all examples, we have $\text{Pic}(X) = \text{NS}(X)$, a torsion-free abelian group. Choose a basis L_1, \dots, L_r of $\text{Pic}(X)$ and metrizations $\mathcal{L}_j = (L_j, \|\cdot\|_v)$. We obtain a *height system*:

$$H_j : X(F) \rightarrow \mathbb{R}_{>0}, \quad \text{for } j = 1, \dots, r,$$

which can be extended to $\text{Pic}(X)_{\mathbb{C}}$, by linearity:

eqn:heights

$$(6.4) \quad \begin{aligned} H : X(F) \times \text{Pic}(X)_{\mathbb{C}} &\rightarrow \mathbb{R}_{>0} \\ (x, \mathbf{s}) &\mapsto \prod_{j=1}^r H_{\mathcal{L}_j}(x)^{s_j}, \end{aligned}$$

where $\mathbf{s} := \sum_{j=1}^r s_j L_j$. For each j , the 1-parameter zeta function

$$Z_X(\mathcal{L}_j, s) = \sum_{x \in X(F)} H_{\mathcal{L}_j}(x)^{-s}$$

converges absolutely to a holomorphic function, for $\Re(s) \gg 0$ (see Remark ??). It follows that

$$Z_X(\mathbf{s}) := \sum_{x \in X(F)} H(\mathbf{s}, x)^{-1}$$

converges absolutely to a holomorphic function for $\Re(\mathbf{s})$ contained some cone in $\text{Pic}(X)_{\mathbb{R}}$.

Step 1. One introduces a generalized height pairing

eqn:heights-adelic

$$(6.5) \quad H = \prod_v H_v : G(\mathbb{A}) \times \text{Pic}(X)_{\mathbb{C}} \rightarrow \mathbb{C},$$

such that the restriction of H to $\text{Pic}(X) \times G(F)$ coincides with the pairing in (6.4). Since X is projective, the height zeta function

eqn:zz

$$(6.6) \quad Z(g, \mathbf{s}) := \sum_{\gamma \in G(F)} H(\gamma g, \mathbf{s})^{-1}$$

converges to a function which is continuous in g and holomorphic in \mathbf{s} for $\Re(\mathbf{s})$ contained in some cone $\Lambda \subset \text{Pic}(X)_{\mathbb{R}}$. The standard height zeta function is obtained by setting $g = e$, the identity in $G(\mathbb{A})$. Our goal is to obtain a meromorphic continuation to the tube domain \mathbb{T} over an open neighborhood of $[-K_X] = \kappa \in \text{Pic}(X)_{\mathbb{R}}$ and to identify the poles of Z in this domain.

Step 2. It turns out that

$$Z(g, \mathbf{s}) \in L^2(G(F) \backslash G(\mathbb{A})),$$

for $\Re(\mathbf{s}) \gg 0$. This is immediate in the cocompact case, e.g., for G unipotent or semisimple anisotropic, and requires an argument in other cases. The L^2 -space decomposes into unitary irreducible representations for the natural action of $G(\mathbb{A})$. We get a formal identity

eqn:formalz

$$(6.7) \quad Z(g, \mathbf{s}) = \sum_{\varrho} Z_{\varrho}(g, \mathbf{s}),$$

where the summation is over all irreducible unitary representations $(\varrho, \mathcal{H}_{\varrho})$ of $G(\mathbb{A})$ occurring in the right regular representation of $G(\mathbb{A})$ in $L^2(G(F) \backslash G(\mathbb{A}))$.

Step 3. In many cases, the leading pole of $Z(g, \mathbf{s})$ arises from the trivial representation, i.e., from the integral

eqn:trivial

$$(6.8) \quad \int_{G(\mathbb{A}_F)} \mathbf{H}(g, \mathbf{s})^{-1} dg = \prod_v \int_{G(F_v)} \mathbf{H}_v(g_v, \mathbf{s})^{-1} dg_v,$$

where dg_v is a Haar measure on $G(F_v)$. To simplify the exposition we assume that

$$X \setminus G = D = \cup_{i \in \mathcal{I}} D_i,$$

where D is a divisor with normal crossings whose components D_i are geometrically irreducible.

We choose integral models for X and D_i and observe

$$G(F_v) \subset X(F_v) \xrightarrow{\sim} X(\mathcal{O}_v) \rightarrow X(\mathbb{F}_q) = \cup_{I \subset \mathcal{I}} D_I^{\circ}(\mathbb{F}_q),$$

where

$$D_I := \cap_{i \in I} D_i, \quad D_I^{\circ} := D_I \setminus \cup_{I' \supsetneq I} D_{I'}.$$

Write $\mathbf{s} = \sum_i s_i D_i$ and For almost all v , we have:

eqn:local-intt

$$(6.9) \quad \int_{G(F_v)} \mathbf{H}_v(g_v, \mathbf{s})^{-1} dg_v = \tau_v(G)^{-1} \left(\sum_{I \subset \mathcal{I}} \frac{\#D_I^{\circ}(\mathbb{F}_q)}{q^{\dim(X)}} \prod_{i \in I} \frac{q-1}{q^{s_i - \kappa_i + 1} - 1} \right),$$

where $\tau_v(G)$ is the local Tamagawa number of G and κ_i is the order of the pole of the (unique modulo constants) top-degree differential form on G along D_i (SEE ...). The height integrals are geometric versions of Igusa's integrals. They are closely related to "motivic" integrals of Batyrev, Kontsevich, Denef and Loeser (see [?], [?] and [?]).

This allows to regularize explicitly this adelic integral. For example, for unipotent G we have

`eqn:global-int`

$$(6.10) \quad \int_{G(\mathbb{A}_F)} \mathbf{H}(g, \mathbf{s})^{-1} dg = \prod_i \zeta_F(s_i - \kappa_i + 1) \cdot \Phi(\mathbf{s}),$$

with $\Phi(s)$ holomorphic and absolutely bounded for $\Re(s_i) > \kappa_i - \delta$, for all i .

Step 4. Next, one has to identify the leading poles of $Z_\varrho(g, \mathbf{s})$, and to obtain bounds which are sufficiently uniform in ϱ to yield a meromorphic continuation of the right side of (6.15). This is nontrivial already for abelian groups G (see Section ?? for the case when $G = \mathbb{G}_a^n$). Moreover, will need to show *pointwise* convergence of the series, as a function of $g \in G(\mathbb{A})$.

For G abelian, e.g., an algebraic torus, all unitary representation have dimension one, and equation (6.15) is nothing but the usual Fourier expansion of a “periodic” function. The adelic Fourier coefficient is an Euler product, and the local integrals can be evaluated explicitly.

For other groups, it is important to have some sort of parametrization of representations occurring on the right side of the spectral expansion. For example, for unipotent groups such a representation is provided by Kirillov’s orbit method (see Section 6.6). For semisimple groups one has to appeal to Langland’s theory of automorphic representations (see Section ??).

`sect:additive`

6.4. Additive groups. Let X be an equivariant compactification of an additive group $G = \mathbb{G}_a^n$. For example, any blowup $X = \text{Bl}_Y(\mathbb{P}^n)$, with $Y \subset \mathbb{P}^{n-1} \subset \mathbb{P}^n$, can be equipped with a structure of an equivariant compactification of \mathbb{G}_a^n . In particular, the Hilbert schemes of all algebraic subvarieties of \mathbb{P}^{n-1} appear in the moduli of equivariant compactifications X as above. Some features of the geometry of such compactifications have been explored in [HT99]. The analysis of height zeta functions has to capture this geometric complexity. In this section we present an approach to height zeta functions developed in [CLT99], [CLT00], and [CLT02].

Indeed, the height pairing (6.5) determines X uniquely ??? (SEE ABOVE).

The Poisson formula yields

$$\begin{aligned} Z(\mathbf{s}) &= \sum_{x \in G(F)} H(x, \mathbf{s})^{-1} \\ &= \int_{G(\mathbb{A}_F)} H(x, \mathbf{s})^{-1} dx + \sum_{\psi \neq \psi_0} \hat{H}(\psi, \mathbf{s}), \end{aligned}$$

where the sum runs over all nontrivial characters $\psi \in (G(\mathbb{A}_F)/G(F))^*$ and

$$\hat{H}(\psi, \mathbf{s}) = \int_{G(\mathbb{A}_F)} H(x, \mathbf{s})^{-1} \psi(x) dx$$

is the Fourier transform, with an appropriately normalized Haar measure dx .

exam:p1

Example 6.4.1. The simplest case is $G = \mathbb{G}_a = \mathbb{A}^1 \subset \mathbb{P}^1$, over $F = \mathbb{Q}$, with the standard height

$$H_p(x) = \max(1, |x|_p), \quad H_\infty(x) = \sqrt{1 + x^2}.$$

We have

eqn:ga-zeta

$$(6.11) \quad Z(s) = \sum_{x \in \mathbb{Q}} H(x)^{-s} = \int_{\mathbb{A}_\mathbb{Q}} H(x)^{-s} dx + \sum_{\psi} \hat{H}(\psi, s).$$

The local Haar measure dx_p is normalized by $\text{vol}(\mathbb{Z}_p) = 1$ so that

$$\text{vol}(|x|_p = p^j) = p^j \left(1 - \frac{1}{p}\right).$$

We have

$$\begin{aligned} \int_{\mathbb{Q}_p} H_p(x)^{-s} dx_p &= \int_{\mathbb{Z}_p} H_p(x)^{-s} dx_p + \sum_{j \geq 1} \int_{|x|_p = p^j} H_p(x)^{-s} dx_p \\ &= 1 + \sum_{j \geq 1} p^{-js} \text{vol}(|x|_p = p^j) = \frac{1 - p^{-s}}{1 - p^{-(s-1)}} \\ \int_{\mathbb{R}} (1 + x^2)^{-s/2} dx &= \frac{\Gamma((s-1)/2)}{\Gamma(s/2)}. \end{aligned}$$

Now we analyze the contributions from nontrivial characters. Each such character ψ decomposes as a product of local characters, defined as follows:

$$\begin{aligned} \psi_p &= \psi_{p, a_p} : x_p \mapsto e^{2\pi i a_p \cdot x_p}, \quad a_p \in \mathbb{Q}_p, \\ \psi_\infty &= \psi_{\infty, a_\infty} : x \mapsto e^{2\pi i a_\infty \cdot x}, \quad a \in \mathbb{R}. \end{aligned}$$

A character is *unramified* at p if it is trivial on \mathbb{Z}_p , i.e., $a_p \in \mathbb{Z}_p$. Then $\psi = \psi_a$, with $a \in \mathbb{A}_{\mathbb{Q}}$. A character $\psi = \psi_a$ is unramified for all p iff $a \in \mathbb{Z}$. Pontryagin duality identifies $\hat{\mathbb{Q}}_p = \mathbb{Q}_p$, $\hat{\mathbb{R}} = \mathbb{R}$, and $(\mathbb{A}_{\mathbb{Q}}/\mathbb{Q})^* = \mathbb{Q}$.

Since H_p is invariant under the translation action by \mathbb{Z}_p , the local Fourier transform $\hat{H}_p(\psi_{a_p})$ vanishes unless ψ_p is unramified at p . In particular, only unramified characters are present in the expansion (6.11), i.e., we may assume that $\psi = \psi_a$ with $a \in \mathbb{Z} \setminus 0$. For $p \nmid a$, we compute

$$\hat{H}_p(s, \psi_a) = 1 + \sum_{j \geq 1} p^{-sj} \int_{|x|_p=p^j} \psi_a(x_p) dx_p = 1 - p^{-s}.$$

Putting together we obtain

$$\begin{aligned} Z(s) &= \frac{\zeta(s-1)}{\zeta(s)} \cdot \frac{\Gamma((s-1)/2)}{\Gamma(s/2)} \\ &+ \sum_{a \in \mathbb{Z}} \prod_{p \nmid a} \frac{1}{\zeta_p(s)} \cdot \prod_{p|a} \hat{H}_p(x_p)^{-s} dx_p \cdot \int_{\mathbb{R}} (1+x^2)^{-s/2} \cdot e^{2\pi i a x} dx \end{aligned}$$

For $\Re(s) > 2 - \delta$, we have the upper bounds

eqn:ga-ineq

$$(6.12) \quad \left| \prod_{p|a} \hat{H}_p(x_p)^{-s} dx_p \right| \ll \left| \prod_{p|a} \int_{\mathbb{Q}_p} H_p(x_p)^{-s} dx_p \right| \ll |a|^\epsilon$$

$$(6.13) \quad \left| \int_{\mathbb{R}} (1+x^2)^{-s/2} \cdot e^{2\pi i a x} dx \right| \ll_N \frac{1}{(1+|a|)^N}, \quad \text{for any } N \in \mathbb{N},$$

where the second inequality is proved via repeated integration by parts.

This gives a meromorphic continuation of $Z(s)$ and its pole at $s = 2$ (corresponding to $-K_X = 2L \in \mathbb{Z} = \text{Pic}(\mathbb{P}^1)$). The leading coefficient at this pole is the Tamagawa number defined by Peyre.

Now we turn to the general case.

- $\text{Pic}(X) = \bigoplus_i \mathbb{Z} D_i$
- $-K_X = \sum_i \kappa_i D_i$, with $\kappa_i \geq 2$
- $\Lambda_{\text{eff}}(X) = \bigoplus_i \mathbb{R}_{\geq 0} D_i$

Local and global heights are given by

$$H_{D_i, v}(x) := \|\mathbf{f}_i(x)\|_v^{-1} \quad \text{and} \quad H_{D_i}(x) = \prod_v H_{D_i, v}(x),$$

where \mathbf{f}_i is the unique G -invariant section of $H^0(X, D_i)$. We get a *height pairing*:

$$\begin{aligned} H : G(\mathbb{A}_F) \times \text{Pic}(X)_{\mathbb{C}} &\rightarrow \mathbb{C} \\ (x, \sum_i s_i D_i) &\mapsto \prod_i H_{D_i}(x)^{s_i} \end{aligned}$$

Similarly, obtain characters of $\mathbb{G}_a^n(\mathbb{A}_F)$.

A character is determined by a “linear form” $\langle a, \cdot \rangle = f_{\mathbf{a}}$, on \mathbb{G}_a^n , which gives a rational function $f_{\mathbf{a}} \in F(X)^*$. We have

$$\operatorname{div}(f_{\mathbf{a}}) = E_{\mathbf{a}} - \sum_i d_i(f_{\mathbf{a}})D_i$$

with $d_i \geq 0$, for all i .

Define:

- $S(\mathbf{a}) \subset \operatorname{Val}(F)$
- $\mathcal{I}_0(\mathbf{a}) := \{i \mid d_i(f_{\mathbf{a}}) = 0\} \subsetneq \mathcal{I}$

We have

$$\hat{H}(\psi_{\mathbf{a}}, \mathbf{s}) = \prod_{i \in \mathcal{I}_0(\mathbf{a})} \zeta_F(s_i - \kappa_i + 1) \cdot \Phi_{\mathbf{a}}(\mathbf{s}) \cdot \int_{G(\mathbb{A}_{\infty})} H_{\infty}(x, \mathbf{s})^{-1} \psi_{\mathbf{a}, \infty}(x) dx$$

with $\Phi_{\mathbf{a}}(\mathbf{s})$ holomorphic for $\Re(s_i) > \kappa_i - \delta$ and bounded by $(1 + \|a\|)^{\epsilon}$.

$$\begin{aligned} Z(\mathbf{s}) &= \int_{G(\mathbb{A}_F)} H(x, \mathbf{s})^{-1} dx + \sum_{\mathcal{I}_0 \subsetneq \mathcal{I}} \sum_{\psi_{\mathbf{a}}: \mathcal{I}_0(\mathbf{a}) = \mathcal{I}} \hat{H}(\psi_{\mathbf{a}}, \mathbf{s}) \\ &= \prod_{i \in \mathcal{I}} \zeta_F(s_i - \kappa_i + 1) \cdot \Phi(\mathbf{s}) \\ &\quad + \sum_{I \subset \mathcal{I}} \prod_{i \in I} \zeta_F(s_i - \kappa_i + 1) \cdot \tilde{\Phi}_I(\mathbf{s}) \end{aligned}$$

sect:toric

6.5. Toric varieties. Analytic properties of height zeta functions of toric varieties have been established in [BT95], [BT98a], and [?].

An algebraic torus is a linear algebraic groups T over a field F such that

$$T_E \simeq \mathbb{G}_{m,E}^d$$

for some finite Galois extension E/F . Such an extension is called a *splitting field* of T . A torus is *split* if $T \simeq \mathbb{G}_{m,F}^d$. The group of algebraic characters

$$M := \mathfrak{X}^*(T) = \operatorname{Hom}(T, \bar{K}^*)$$

is a torsion free $\Gamma := \operatorname{Gal}(E/F)$ -module. The standard notation for its dual, the cocharacters is $N := \mathfrak{X}^*(T)$. There is an equivalence of categories:

$$\left\{ \begin{array}{l} d - \text{dimensional integral} \\ \Gamma - \text{representations,} \\ \text{up to equivalence} \end{array} \right\} \Leftrightarrow \left\{ \begin{array}{l} d - \text{dimensional} \\ \text{algebraic tori, split over } E, \\ \text{up to isomorphism} \end{array} \right\}$$

The local and global theory of tori can be summarized as follows: The local Galois groups $\Gamma_v := \operatorname{Gal}(E_w/F_v) \subset \Gamma$ act on $M_v := M^{\Gamma_v}$, the

characters of $T(F_v)$. Let N_v be the lattice of local cocharacters. Write $T(\mathfrak{o}_v) \subset T(F_v)$ for a maximal compact subgroup (after choosing an integral model, it is indeed the group of \mathfrak{o}_v -valued points of T , for almost all v). Then

$$T(F_v)/T(\mathfrak{o}_v) \hookrightarrow N_v = N^{\Gamma_v},$$

an isomorphism for v unramified in E/F . Adelically, we have

$$T(\mathbb{A}_F) \supset T^1(\mathbb{A}_F) = \{t \mid \prod_v |m(t_v)|_v = 1 \ \forall m \in M^\Gamma\}$$

and

$$T(F) \hookrightarrow T^1(\mathbb{A}_F).$$

Let $\mathbf{K}_T := \prod_v T(\mathfrak{o}_v)$ be the maximal compact subgroup of $T(\mathbb{A}_F)$.

thm:tori-adelic

Theorem 6.5.1. *We have*

- $T(\mathbb{A}_F)/T^1(\mathbb{A}_F) = N_{\mathbb{R}}^\Gamma$;
- $T^1(\mathbb{A}_F)/T(F)$ compact;
- $\mathbf{K}_T \cap T(F)$ finite;
- the map $(T(\mathbb{A}_F)/\mathbf{K}_T \cdot T(F))^* \rightarrow \bigoplus_{v|\infty} M_v \otimes \mathbb{R}$ has finite kernel (analogs of roots of 1) and maps the characters to a lattice $\bigoplus M_{\mathbb{R}}^\Gamma$.

Over algebraically closed fields, complete *toric varieties*, i.e., equivariant compactifications of algebraic tori, are described and classified by a combinatorial structure (M, N, Σ) , where

- M is the free abelian group of finite rank (the algebraic characters of the torus T),
- $N := \text{Hom}(M, \mathbb{Z})$ is the dual group of cocharacters, and
- $\Sigma = \{\sigma\}$ is a *fan*, i.e., a collection of strictly convex cones in $N_{\mathbb{R}}$ such that
 - (1) $0 \in \sigma$ for all $\sigma \in \Sigma$,
 - (2) $N_{\mathbb{R}} = \bigcup_{\sigma \in \Sigma} \sigma$,
 - (3) every face $\tau \subset \sigma$ is in Σ
 - (4) $\sigma \cap \sigma' \in \Sigma$ and is face of σ, σ' .

A fan Σ is called *regular* if the generators of every $\sigma \in \Sigma$ form part of a basis of N . In this case, the corresponding toric variety X_Σ is smooth. The toric variety is constructed as follows:

$$X_\Sigma := \bigcup_{\sigma} U_\sigma \quad \text{where} \quad U_\sigma := \text{Spec}(F[M \cap \check{\sigma}]),$$

and $\check{\sigma} \subset M_{\mathbb{R}}$ is the cone dual to $\sigma \subset N_{\mathbb{R}}$. The fan Σ encodes all geometric information about X_Σ . For example, 1-dimensional generators e_1, \dots, e_n of Σ correspond to boundary divisors D_1, \dots, D_n , i.e.,

the irreducible components of $X_\Sigma \setminus T$. There is an explicit criterion for projectivity and a description of the cohomology ring, cellular structure etc.

Example 6.5.2. The simplest toric variety is $X = \mathbb{P}^1$. We have three distinguished Zariski open subsets:

- $\mathbb{P}^1 \supset \mathbb{G}_m = \text{Spec}(F[x, x^{-1}]) = \text{Spec}(F[x^{\mathbb{Z}}])$
- $\mathbb{P}^1 \supset \mathbb{A}^1 = \text{Spec}(F[x]) = \text{Spec}(F[x^{\mathbb{Z}_{\geq 0}}])$,
- $\mathbb{P}^1 \supset \mathbb{A}^1 = \text{Spec}(F[x^{-1}]) = \text{Spec}(F[x^{\mathbb{Z}_{\leq 0}}])$

They correspond to the semi-groups:

- \mathbb{Z} dual to 0,
- $\mathbb{Z}_{\geq 0}$ dual to $\mathbb{Z}_{\geq 0}$,
- $\mathbb{Z}_{\leq 0}$ dual to $\mathbb{Z}_{\leq 0}$.

Over nonclosed ground fields F one has to account for the action of the Galois group of a splitting field E/F . The necessary modifications can be described as follows. The Galois group Γ acts on M, N, Σ . A fan Σ is called Γ -invariant if $\gamma \cdot \sigma \in \Sigma$, for all $\gamma \in \Gamma, \sigma \in \Sigma$. If Σ is a complete regular Γ -invariant fan such that, over the splitting field, the resulting toric variety $X_{\Sigma, E}$ is projective, then it can be descended to a complete algebraic variety $X_{\Sigma, F}$ over the groundfield F such that

$$X_{\Sigma, E} \simeq X_{\Sigma, F} \otimes_{\text{Spec}(F)} \text{Spec}(E),$$

as E -varieties with Γ -action.

Picard group

The split case

$PL(\Sigma)$ - piecewise linear \mathbb{Z} -valued functions φ on Σ determined by $M \supset \{m_{\sigma, \varphi}\}_{\sigma \in \Sigma}$, i.e., by its values $s_j := \varphi(e_j)$, $j = 1, \dots, n$.

$$0 \rightarrow M \rightarrow PL(\Sigma) \xrightarrow{\pi} \text{Pic}(X_\Sigma) \rightarrow 0$$

- every divisor is equivalent to a linear combination of boundary divisors D_1, \dots, D_n , and φ is determined by its values on e_1, \dots, e_n
- relations come from characters of T

The nonsplit case

$$0 \rightarrow M^\Gamma \rightarrow PL(\Sigma)^\Gamma \xrightarrow{\pi} \text{Pic}(X_\Sigma)^\Gamma \rightarrow H^1(\Gamma, M) \rightarrow 0$$

$$\begin{aligned}\Lambda_{\text{eff}}(X_\Sigma) &= \pi(\mathbb{R}_{\geq 0}D_1 + \dots + \mathbb{R}_{\geq 0}D_n) \\ -K_\Sigma &= \pi(D_1 + \dots + D_n)\end{aligned}$$

Example 6.5.3. $\mathbb{P}^1 = \{x = (x_0 : x_1)\} \supset \mathbb{G}_m$

$$H_v(x) := \begin{cases} \left| \frac{x_0}{x_1} \right|_v & \text{if } |x_0|_v \geq |x_1|_v \\ \left| \frac{x_1}{x_0} \right|_v & \text{otherwise} \end{cases}$$

$$H(x) := \prod_v H_v(x)$$

In general, $T(F_v)/T(\mathcal{O}_v) \hookrightarrow N_v$. For $\varphi \in PL(\Sigma)$ put

$$H_{\Sigma,v}(x, \varphi) := q_v^{\varphi(\bar{x}_v)} \quad H_\Sigma(x, \varphi) := \prod_v H_{\Sigma,v}(x, \varphi),$$

with $q_v = e$, for $v \mid \infty$. This height has the following properties:

- it gives a pairing $T(\mathbb{A}_F) \times PL(\Sigma)_\mathbb{C} \rightarrow \mathbb{C}$;
- the restriction to $T(F) \times PL(\Sigma)_\mathbb{C}$ descends to a well-defined pairing

$$T(F) \times \text{Pic}(X_\Sigma)_\mathbb{C} \rightarrow \mathbb{C};$$

- $T(\mathcal{O}_v)$ -invariance, for all v

Height zeta function:

$$Z_\Sigma(\mathbf{s}) := \sum_{x \in T(F)} H_\Sigma(x, \varphi_{\mathbf{s}})^{-1}$$

Poisson formula

$$\begin{aligned}Z_\Sigma(\mathbf{s}) &:= \int_{(T(\mathbb{A}_F)/\mathbf{K}_T \cdot T(F))^*} \hat{H}_\Sigma(\chi, \mathbf{s}) d\chi \\ \hat{H}_\Sigma(\chi, \mathbf{s}) &:= \int_{T(\mathbb{A}_F)} H_\Sigma(x, -\varphi_{\mathbf{s}}) \chi(x) dx\end{aligned}$$

- for χ nontrivial on \mathbf{K}_T we have $\hat{H}_\Sigma = 0$
- converges absolutely for $\Re(s_j) > 1$ (for all j)
- Haar measures on $T(F_v)$ normalized by $T(\mathcal{O}_v)$

Example 6.5.4. Consider the projective line \mathbb{P}^1 over \mathbb{Q} . We have

$$0 \rightarrow M \rightarrow PL(\Sigma) \rightarrow \text{Pic}(\mathbb{P}^1) \rightarrow 0$$

with $M = \mathbb{Z}$ and $PL(\Sigma) = \mathbb{Z}^2$. The Fourier transforms of local heights can be computed as follows:

$$\hat{H}_p(\chi_0, \mathbf{s}) = 1 + \sum_{n \geq 1} p^{-s_1 - im} + \sum_{n \geq 1} p^{-s_1 + im} = \frac{\zeta_p(s_1 + im)\zeta_p(s_2 - im)}{\zeta_p(s_1 + s_2)},$$

$$\hat{H}_\infty(\chi_0, \mathbf{s}) = \int_0^\infty e^{(-s_1 - im)x} dx + \int_0^\infty e^{(-s_2 + im)x} dx = \frac{1}{s_1 + im} + \frac{1}{s_2 - im}.$$

We obtain

$$Z_{\mathbb{P}^1}(s_1, s_2) = \int_{\mathbb{R}} \zeta(s_1 + im)\zeta(s_2 - im) \cdot \left(\frac{1}{s_1 + im} + \frac{1}{s_2 - im} \right) dm.$$

The integral converges for $\Re(s_1), \Re(s_2) > 1$, absolutely and uniformly on compact subsets. It remains to establish its meromorphic continuation. This can be achieved by shifting the contour of integration and computing the resulting residues.

It is helpful to compare this approach with the analysis of \mathbb{P}^1 as an additive variety in Example ??.

The Fourier transforms of local height functions in the case of \mathbb{G}_m^d over \mathbb{Q} are given by:

- $v \nmid \infty$:

$$\hat{H}_{\Sigma, v}(\chi_v, -\mathbf{s}) = \sum_{k=1}^d \sum_{\sigma \in \Sigma(k)} (-1)^k \prod_{e_j \in \sigma} \frac{1}{1 - q_v^{-(s_j + i\langle e_j, m \rangle)}}$$

- $v \mid \infty$

$$\hat{H}_{\Sigma, v}(\chi_v, -\mathbf{s}) = \sum_{\sigma \in \Sigma(d)} \prod_{e_j \in \sigma} \frac{1}{(s_j + i\langle e_j, m \rangle)}$$

where $\chi = \chi_m$ is the character corresponding to $m \in M_{\mathbb{R}}$. The general case of nonsplit tori over number fields requires more care. We have an exact sequence of Γ -modules:

$$0 \rightarrow M \rightarrow PL(\Sigma) \rightarrow \text{Pic}(X_\Sigma) \rightarrow 0,$$

with $PL(\Sigma)$ a permutation module. Duality gives a sequence of groups:

$$0 \rightarrow T_{\text{Pic}}(\mathbb{A}_F) \rightarrow T_{PL}(\mathbb{A}_F) \rightarrow T(\mathbb{A}_F)$$

with

$$T_{PL}(\mathbb{A}_F) = \prod_{j=1}^k R_{F_j/F} \mathbb{G}_m(\mathbb{A}_F) \quad (\text{restriction of scalars})$$

We get a map

$$\begin{array}{ccc} (T(\mathbb{A}_F)/\mathbf{K}_T \cdot T(F))^* & \rightarrow & \prod_{j=1}^k (\mathbb{G}_m(\mathbb{A}_{F_j})/\mathbb{G}_m(F_j))^* \\ \chi & \mapsto & (\chi_1, \dots, \chi_k) \end{array}$$

This map has finite kernel. Assembling local computations, we have

eqn:hat-h

$$(6.14) \quad \hat{H}_\Sigma(\chi, \mathbf{s}) = \frac{\prod_{j=1}^k L(s_j, \chi_j)}{Q_\Sigma(\chi, \mathbf{s})} \zeta_{\Sigma, \infty}(\mathbf{s}, \chi),$$

where $Q_\Sigma(\chi, \mathbf{s})$ bounded uniformly in χ , in compact subsets in $\Re(s_j) > 1/2 + \delta$, $\delta > 0$, and

$$|\zeta_{\Sigma, \infty}(\mathbf{s}, \chi)| \ll \frac{1}{(1 + \|m\|)^{d+1}} \cdot \frac{1}{(1 + \|\chi\|)^{d'+1}}.$$

This implies that

$$Z_\Sigma(\mathbf{s}) = \int_{M_{\mathbb{R}}^\Gamma} f_\Sigma(\mathbf{s} + im) dm,$$

where

$$f_\Sigma(\mathbf{s}) := \sum_{\chi \in (T^1(\mathbb{A}_F)/\mathbf{K}_T \cdot T(F))^*} \hat{H}_\Sigma(\chi, \mathbf{s})$$

We have

- (1) $(s_1 - 1) \cdot \dots \cdot (s_k - 1) f_\Sigma(\mathbf{s})$ is holomorphic for $\Re(s_j) > 1 - \delta$;
- (2) f_Σ satisfies growth conditions in vertical strips (this follows by applying the Phragmen-Lindelöf principle 6.1.2 to bound L -functions appearing in equation (6.14));
- (3) $\lim_{s_j \rightarrow 1} f_\Sigma(\mathbf{s}) = c(f_\Sigma) \neq 0$.

The Convexity principle 6.1.1 implies a meromorphic continuation of $Z(\mathbf{s})$ to a tubular neighborhood of the shifted cone $\Lambda_{\text{eff}}(X_\Sigma)$. The identification of the remaining factors β and τ requires another application of the Poisson formula. Other line bundles require a version of the technical theorem above.

sect:unipotent

6.6. Unipotent groups. Let $X \supset G$ be an equivariant compactification of a unipotent group over a number field F and

$$X \setminus G = D = \cup_{i \in \mathcal{I}} D_i.$$

Throughout, we will assume that G acts on X on both sides, i.e., that X is a compactification of $G \times G/G$, or a bi-equivariant compactification. We also assume that D is a divisor with normal crossings and its components D_i are geometrically irreducible. The main geometric invariants of X have been computed in Example 1.1.4: The Picard

group is freely generated by the classes of D_i , the effective cone is simplicial, and the anticanonical class is sum of boundary components with nonnegative coefficients.

Local and global heights have been defined in Example 4.10.6:

$$H_{D_i,v}(x) := \|f_i(x)\|_v^{-1} \quad \text{and} \quad H_{D_i}(x) = \prod_v H_{D_i,v}(x),$$

where f_i is the unique G -invariant section of $H^0(X, D_i)$. We get a *height pairing*:

$$H : G(\mathbb{A}_F) \times \text{Pic}(X)_{\mathbb{C}} \rightarrow \mathbb{C}$$

as in Section 6.3. The bi-equivariance of X implies that H is invariant under the action *on both sides* of a compact open subgroup \mathbf{K} of the finite adeles. Moreover, we can arrange that H_v is smooth in g_v for archimedean v .

The height zeta function

$$Z(\mathbf{s}, g) := \sum_{\gamma \in G(F)} H(\mathbf{s}, g)^{-1}$$

is holomorphic in \mathbf{s} , for $\Re(\mathbf{s}) \gg 0$. As a function of g it is continuous and in $L^2(G((F) \backslash G(\mathbb{A}_F)))$, for these \mathbf{s} . We proceed to analyze its spectral decomposition. We get a formal identity

eqn:formalz

$$(6.15) \quad Z(\mathbf{s}; g) = \sum_{\varrho} Z_{\varrho}(\mathbf{s}; g),$$

where the sum is over all irreducible unitary representations $(\varrho, \mathcal{H}_{\varrho})$ of $G(\mathbb{A}_F)$ occurring in the right regular representation of $G(\mathbb{A}_F)$ in $L^2(G(F) \backslash G(\mathbb{A}_F))$. They are parametrized by F -rational orbits $\mathcal{O} = \mathcal{O}_{\varrho}$ under the coadjoint action of G on the dual of its Lie algebra \mathfrak{g}^* . The relevant orbits are *integral* - there exists a lattice in $\mathfrak{g}^*(F)$ such that $Z_{\varrho}(\mathbf{s}; g) = 0$ unless the intersection of \mathcal{O} with this lattice is nonempty. The pole of highest order is contributed by the trivial representation and integrality insures that this representation is “isolated”.

Let ϱ be an integral representation as above. It has the following explicit realization: There exists an F -rational subgroup $M \subset G$ such that

$$\varrho = \text{Ind}_M^G(\psi),$$

where ψ is a certain character of $M(\mathbb{A}_F)$. In particular, for the trivial representation, $M = G$ and ψ is the trivial character. Further, there

exists a finite set of valuations $S = S_\varrho$ such that $\dim(\varrho_v) = 1$ for $v \notin S$ and consequently

$$\boxed{\text{eqn:zeta}} \quad (6.16) \quad Z_\varrho(\mathbf{s}; g') = Z^S(\mathbf{s}; g') \cdot Z_S(\mathbf{s}; g').$$

It turns out that

$$Z^S(\mathbf{s}; g') := \prod_{v \notin S} \int_{M(F_v)} H_v(\mathbf{s}; g_v g'_v)^{-1} \bar{\psi}(g_v) dg_v,$$

with an appropriately normalized Haar measure dg_v on $M(F_v)$. The function Z_S is the projection of Z to $\otimes_{v \in S} \varrho_v$.

The first key result is the explicit computation of *height integrals*:

$$\int_{M(F_v)} H_v(\mathbf{s}; g_v g'_v)^{-1} \bar{\psi}(g_v) dg_v$$

for almost all v . This has been done in [?] for equivariant compactifications of additive groups \mathbb{G}_a^n (see Section 6.4); the same approach works here too. The contribution from the trivial representation can be computed using the formula of Denef-Loeser, as in (6.10):

$$\int_{G(\mathbb{A}_F)} H(\mathbf{s}; g)^{-1} dg = \prod_i \zeta_F(s_i - \kappa_i + 1) \cdot \Phi(\mathbf{s}),$$

where $\Phi(\mathbf{s})$ is holomorphic for $\Re(\mathbf{s}) \in \mathbb{T}_{-K_X - \epsilon}$, for some $\epsilon > 0$, and $-K_X = \sum_i \kappa_i D_i$. As in the case of additive groups in Section 6.4, this term gives the “correct” pole at $-K_X$. The analysis of 1-dimensional representations, with $M = G$, is similar to the additive case. New difficulties arise from infinite-dimensional ϱ on the right side of the expansion (6.15).

Next we need to estimate $\dim(\varrho_v)$ and the local integrals for nonarchimedean $v \in S_\varrho$. The key result here is that the contribution to the Euler product from these places is a holomorphic function which can be bounded from above by a *polynomial* in the coordinates of ϱ , for $\Re(\mathbf{s}) \in \mathbb{T}_{-K_X - \epsilon}$. The uniform convergence of the spectral expansion comes from estimates at the archimedean places: for every (left or right) G -invariant differential operator ∂ (and $\mathbf{s} \in \mathbb{T}$) there exists a constant $c(\partial)$ such that

$$\boxed{\text{eqn:c}} \quad (6.17) \quad \int_{G(F_v)} |\partial H_v(\mathbf{s}; g_v)^{-1} dg_v|_v \leq c(\partial).$$

Let v be real. It is known that ϱ_v can be modeled in $L^2(\mathbb{R}^r)$, where $2r = \dim(\mathcal{O})$. More precisely, there exists an isometry

$$j : (\pi_v, L^2(\mathbb{R}^r)) \rightarrow (\varrho_v, \mathcal{H}_v)$$

(an analog of the Θ -distribution). Moreover, the universal enveloping algebra $\mathfrak{U}(\mathfrak{g})$ surjects onto the Weyl algebra of differential operators with polynomial coefficients acting on the smooth vectors $C^\infty(\mathbb{R}^r) \subset L^2(\mathbb{R}^r)$. In particular, we can find an operator Δ acting as the (r -dimensional) harmonic oscillator

$$\prod_{j=1}^r \left(\frac{\partial^2}{\partial x_j^2} - a_j x_j^2 \right),$$

with $a_j > 0$. We choose an orthonormal basis of $L^2(\mathbb{R}^r)$ consisting of Δ -eigenfunctions $\{\tilde{\omega}_\lambda\}$ (which are well known) and analyze

$$\int_{G(F_v)} H_v(\mathbf{s}; g_v)^{-1} \bar{\omega}_\lambda(g_v) dg_v,$$

where $\omega_\lambda = j^{-1}(\tilde{\omega}_\lambda)$. Using integration by parts we find that for $\mathbf{s} \in \mathbb{T}$ and any $N \in \mathbb{N}$ there is a constant $c(N, \Delta)$ such that this integral is bounded by

eqn: bound

 (6.18)
$$(1 + |\lambda|)^{-N} c(N, \Delta).$$

This estimate suffices to conclude that for *each* ϱ the function Z_{S_ϱ} is holomorphic in \mathbb{T} .

Now the issue is to prove the convergence of the sum in (6.15). Using any element $\partial \in \mathfrak{U}(\mathfrak{g})$ acting in \mathcal{H}_ϱ by a scalar $\lambda(\partial) \neq 0$ (for example, any element in the center of $\mathfrak{U}(\mathfrak{g})$) we can improve the bound (6.18) to

$$(1 + |\lambda|)^{-N_1} \lambda(\partial)^{-N_2} c(N_1, N_2, \Delta, \partial)$$

(for any $N_1, N_2 \in \mathbb{N}$). However, we have to insure the uniformity of such estimates over the set of all ϱ . This relies on a parametrization of coadjoint orbits. There is a finite set $\{\Sigma_{\mathbf{d}}\}$ of “packets” of coadjoint orbits, each parametrized by a locally closed subvariety $Z_{\mathbf{d}} \subset \mathfrak{g}^*$, and for each \mathbf{d} a finite set of F -rational polynomials $\{P_{\mathbf{d},r}\}$ on \mathfrak{g}^* such that the restriction of each $P_{\mathbf{d},r}$ to $Z_{\mathbf{d}}$ is invariant under the coadjoint action. Consequently, the corresponding derivatives

$$\partial_{\mathbf{d},r} \in \mathfrak{U}(\mathfrak{g})$$

act in \mathcal{H}_ϱ by multiplication by the scalar

$$\lambda_{\varrho,r} = P_{\mathbf{d},r}(\mathcal{O}).$$

There is a similar uniform construction of the “harmonic oscillator” $\Delta_{\mathbf{d}}$ for each \mathbf{d} . Combining the resulting estimates we obtain the uniform convergence of the right hand side in (6.15).

The last technical point is to prove that both expressions (6.6) and (6.15) for $Z(\mathbf{s}; g)$ define *continuous* functions on $G(F)\backslash G(\mathbb{A}_F)$. Then (6.15) gives the desired meromorphic continuation of $Z(\mathbf{s}; e)$.

Background material on representation theory of unipotent groups can be found in the books [CG66], [Dix96] and the papers [Moo65], [Kir99].

sect:semisimple

6.7. Semisimple groups.

sect:homogeneous

6.8. Homogeneous spaces.

REFERENCES

- `abramovich` [Abr08] D. ABRAMOVICH – “Birational geometry for number theorists”, 2008, this volume.
- `amerik-campana` [AC08] E. AMERIK and F. CAMPANA – “Fibrations méromorphes sur certaines variétés à fibré canonique trivial”, *Pure Appl. Math. Q.* **4** (2008), no. 2, part 1, p. 509–545.
- `araujo` [Ara05] C. ARAUJO – “The cone of effective divisors of log varieties after Batyrev”, 2005, [arXiv:math/0502174](https://arxiv.org/abs/math/0502174).
- `amerik-voisin` [AV07] K. AMERIK and C. VOISIN – “Potential density of rational points on the variety of lines of a cubic fourfold”, 2007, [arXiv:0707.3948](https://arxiv.org/abs/0707.3948).
- `baker-diagonal` [Bak89] R. C. BAKER – “Diagonal cubic equations. II”, *Acta Arith.* **53** (1989), no. 3, p. 217–250.
- `Ba` [Bat92] V. V. BATYREV – “The cone of effective divisors of threefolds”, *Proceedings of the International Conference on Algebra, Part 3 (Novosibirsk, 1989)* (Providence, RI), Contemp. Math., vol. 131, Amer. Math. Soc., 1992, p. 337–352.
- `bchm` [BCHM06] C. BIRKAR, P. CASCINI, C. D. HACON and J. MCKERNAN – “Existence of minimal models for varieties of log general type”, 2006, [arXiv.org:math/0610203](https://arxiv.org/abs/math/0610203).
- `browning-hb-sal` [BHBS06] T. D. BROWNING, D. R. HEATH-BROWN and P. SALBERGER – “Counting rational points on algebraic varieties”, *Duke Math. J.* **132** (2006), no. 3, p. 545–578.
- `birch62` [Bir62] B. J. BIRCH – “Forms in many variables”, *Proc. Roy. Soc. Ser. A* **265** (1961/1962), p. 245–263.
- `katsylo-b` [BK85] F. A. BOGOMOLOV and P. I. KATSYLO – “Rationality of some quotient varieties”, *Mat. Sb. (N.S.)* **126(168)** (1985), no. 4, p. 584–589.
- `batyrev-manin` [BM90] V. V. BATYREV and Y. I. MANIN – “Sur le nombre des points rationnels de hauteur borné des variétés algébriques”, *Math. Ann.* **286** (1990), no. 1-3, p. 27–43.

- bog-br** [Bog87] F. A. BOGOMOLOV – “The Brauer group of quotient spaces of linear representations”, *Izv. Akad. Nauk SSSR Ser. Mat.* **51** (1987), no. 3, p. 485–516, 688.
- bruedern-small** [Brü94] J. BRÜDERN – “Small solutions of additive cubic equations”, *J. London Math. Soc. (2)* **50** (1994), no. 1, p. 25–42.
- borevich-sch** [BS66] A. I. BOREVICH and I. R. SHAFAREVICH – *Number theory*, Translated from the Russian by Newcomb Greenleaf. Pure and Applied Mathematics, Vol. 20, Academic Press, New York, 1966.
- bt95** [BT95] V. V. BATYREV and Y. TSCHINKEL – “Rational points of bounded height on compactifications of anisotropic tori”, *Internat. Math. Res. Notices* (1995), no. 12, p. 591–635.
- batyrev-t98** [BT98a] ———, “Manin’s conjecture for toric varieties”, *J. Algebraic Geom.* **7** (1998), no. 1, p. 15–53.
- bt-tamagawa** [BT98b] ———, “Tamagawa numbers of polarized algebraic varieties”, *Astérisque* (1998), no. 251, p. 299–340, *Nombre et répartition de points de hauteur bornée* (Paris, 1996).
- bogomolov-t99** [BT99] F. A. BOGOMOLOV and Y. TSCHINKEL – “On the density of rational points on elliptic fibrations”, *J. Reine Angew. Math.* **511** (1999), p. 87–93.
- bogomolov-t00** [BT00] ———, “Density of rational points on elliptic $K3$ surfaces”, *Asian J. Math.* **4** (2000), no. 2, p. 351–368.
- bruce-wall** [BW79] J. W. BRUCE and C. T. C. WALL – “On the classification of cubic surfaces”, *J. London Math. Soc. (2)* **19** (1979), no. 2, p. 245–256.
- campana02** [Cam04] F. CAMPANA – “Orbifolds, special varieties and classification theory”, *Ann. Inst. Fourier (Grenoble)* **54** (2004), no. 3, p. 499–630.
- cantat** [Can01] S. CANTAT – “Dynamique des automorphismes des surfaces $K3$ ”, *Acta Math.* **187** (2001), no. 1, p. 1–57.
- cassels-bounds** [Cas55] J. W. S. CASSELS – “Bounds for the least solutions of homogeneous quadratic equations”, *Proc. Cambridge Philos. Soc.* **51** (1955), p. 262–264.
- cast** [Cas07] A.-M. CASTRAVET – “The Cox ring of $\bar{M}_{0,6}$ ”, 2007, to appear in *Trans. AMS*, [arXiv:07050070](https://arxiv.org/abs/07050070).
- cg** [CG66] J. W. S. CASSELS and M. J. T. GUY – “On the Hasse principle for cubic surfaces”, *Mathematika* **13** (1966), p. 111–120.
- clem-grif** [CG72] C. H. CLEMENS and P. A. GRIFFITHS – “The intermediate Jacobian of the cubic threefold”, *Ann. of Math. (2)* **95** (1972), p. 281–356.
- cheltsov** [Che05] I. A. CHELTSOV – “Birationally rigid Fano varieties”, *Uspekhi Mat. Nauk* **60** (2005), no. 5(365), p. 71–160.
- caporaso-h-m95** [CHM97a] L. CAPORASO, J. HARRIS and B. MAZUR – “How many rational points can a curve have?”, *The moduli space of curves* (Texel Island, 1994), *Progr. Math.*, no. 129, Birkhäuser, 1997, p. 13–31.
- caporaso-h-m97** [CHM97b] ———, “Uniformity of rational points”, *J. Amer. Math. Soc.* **10** (1997), no. 1, p. 1–35.
- clemens-k-m88** [CKM88] H. CLEMENS, J. KOLLÁR and S. MORI – “Higher-dimensional complex geometry”, *Astérisque* (1988), no. 166, p. 144 pp. (1989).

- `chambert-loir-t2000` [CLT99] A. CHAMBERT-LOIR and YU. TSCHINKEL – “Points of bounded height on equivariant compactifications of vector groups, I”, *Compositio Math.* **124** (1999), no. 1, p. 65–93.
- `chambert-loir-t2000b` [CLT00] ———, “Points of bounded height on equivariant compactifications of vector groups, II”, *J. Number Theory* **85** (2000), no. 2, p. 172–188.
- `chambert-loir-t2002` [CLT02] ———, “On the distribution of points of bounded height on equivariant compactifications of vector groups”, *Invent. Math.* **148** (2002), p. 421–452.
- `corti` [Cor96] A. CORTI – “Del Pezzo surfaces over Dedekind schemes”, *Ann. of Math. (2)* **144** (1996), no. 3, p. 641–683.
- `cox` [Cox95] D. A. COX – “The homogeneous coordinate ring of a toric variety”, *J. Algebraic Geom.* **4** (1995), no. 1, p. 17–50.
- `cs` [CS06] I. COSKUN and J. STARR – “Divisors on the space of maps to Grassmannians”, *Int. Math. Res. Not.* (2006), p. Art. ID 35273, 25.
- `ct-sal` [CTS89] J.-L. COLLIOT-THÉLÈNE and P. SALBERGER – “Arithmetic on some singular cubic hypersurfaces”, *Proc. London Math. Soc. (3)* **58** (1989), no. 3, p. 519–549.
- `ct-sd` [CTSSD87a] J.-L. COLLIOT-THÉLÈNE, J.-J. SANSUC and P. SWINNERTON-DYER – “Intersections of two quadrics and Châtelet surfaces. I”, *J. Reine Angew. Math.* **373** (1987), p. 37–107.
- `ct-s-swd-chat` [CTSSD87b] ———, “Intersections of two quadrics and Châtelet surfaces. II”, *J. Reine Angew. Math.* **374** (1987), p. 72–168.
- `colliot-ss97` [CTSSD97] J.-L. COLLIOT-THÉLÈNE, A. N. SKOROBOGATOV and P. SWINNERTON-DYER – “Double fibres and double covers: paucity of rational points”, *Acta Arith.* **79** (1997), no. 2, p. 113–135.
- `derenthal` [Der07] U. DERENTHAL – “Universal torsors of del Pezzo surfaces and homogeneous spaces”, *Adv. Math.* **213** (2007), no. 2, p. 849–864.
- `dietmann` [Die03] R. DIETMANN – “Small solutions of quadratic Diophantine equations”, *Proc. London Math. Soc. (3)* **86** (2003), no. 3, p. 545–582.
- `dixmier` [Dix96] J. DIXMIER – *Algèbres enveloppantes*, Les Grands Classiques Gauthier-Villars. [Gauthier-Villars Great Classics], Éditions Jacques Gabay, Paris, 1996, Reprint of the 1974 original.
- `derenthal-joyce` [DJT08] U. DERENTHAL, M. JOYCE and Z. TEITLER – “The nef cone volume of generalized del Pezzo surfaces”, *Algebra Number Theory* **2** (2008), no. 2, p. 157–182.
- `Br` [dlB02] R. DE LA BRETÈCHE – “Nombre de points de hauteur bornée sur les surfaces de del Pezzo de degré 5”, *Duke Math. J.* **113** (2002), no. 3, p. 421–464.
- `dolgachev-bull` [Dol08] I. V. DOLGACHEV – “Reflection groups in algebraic geometry”, *Bull. Amer. Math. Soc. (N.S.)* **45** (2008), no. 1, p. 1–60 (electronic).
- `seminaire79` [DP80] M. DEMAZURE and H. C. PINKHAM (eds.) – *Séminaire sur les Singularités des Surfaces*, Lecture Notes in Mathematics, vol. 777, Springer, Berlin, 1980, Held at the Centre de Mathématiques de l’École Polytechnique, Palaiseau, 1976–1977.
- `els-jah-cubic` [EJ] A.-S. ELSENHANS and J. JAHNEL – “On the smallest point on a diagonal cubic surface”, preprint 2008.

- elsenhans** [EJ06] ———, “The Diophantine equation $x^4 + 2y^4 = z^4 + 4w^4$ ”, *Math. Comp.* **75** (2006), no. 254, p. 935–940 (electronic).
- elkies** [Elk88] N. D. ELKIES – “On $A^4 + B^4 + C^4 = D^4$ ”, *Math. Comp.* **51** (1988), no. 184, p. 825–835.
- esnault** [Esn03] H. ESNAULT – “Varieties over a finite field with trivial Chow group of 0-cycles have a rational point”, *Invent. Math.* **151** (2003), no. 1, p. 187–191.
- faltings83** [Fal83] G. FALTINGS – “Endlichkeitsätze für abelsche Varietäten über Zahlkörpern”, *Invent. Math.* **73** (1983), no. 3, p. 349–366.
- faltings91** [Fal91] ———, “Diophantine approximation on abelian varieties”, *Ann. of Math.* **133** (1991), p. 549–576.
- f1** [Far06] G. FARKAS – “Syzygies of curves and the effective cone of \overline{M}_g ”, *Duke Math. J.* **135** (2006), no. 1, p. 53–98.
- f2** [FG03] G. FARKAS and A. GIBNEY – “The Mori cones of moduli spaces of pointed curves of small genus”, *Trans. Amer. Math. Soc.* **355** (2003), no. 3, p. 1183–1199 (electronic).
- fmt** [FMT89] J. FRANKE, Y. I. MANIN and Y. TSCHINKEL – “Rational points of bounded height on Fano varieties”, *Invent. Math.* **95** (1989), no. 2, p. 421–435.
- gibney** [GKM02] A. GIBNEY, S. KEEL and I. MORRISON – “Towards the ample cone of $\overline{M}_{g,n}$ ”, *J. Amer. Math. Soc.* **15** (2002), no. 2, p. 273–294 (electronic).
- grant-m97** [GM97] G. R. GRANT and E. MANDUCHI – “Root numbers and algebraic points on elliptic surfaces with base \mathbf{P}^1 ”, *Duke Math. J.* **89** (1997), no. 3, p. 413–422.
- gelfand-p69** [GP69] I. M. GELFAND and V. A. PONOMAREV – “Remarks on the classification of a pair of commuting linear transformations in a finite-dimensional space”, *Funkcional. Anal. i Priložen.* **3** (1969), no. 4, p. 81–82.
- hartshorne** [Har77] R. HARTSHORNE – *Algebraic geometry*, Springer-Verlag, New York, 1977, Graduate Texts in Mathematics, No. 52.
- harari-trans** [Har96] D. HARARI – “Obstructions de Manin transcendantes”, Number theory (Paris, 1993–1994), London Math. Soc. Lecture Note Ser., vol. 235, Cambridge Univ. Press, Cambridge, 1996, p. 75–87.
- harari-weak** [Har04] ———, “Weak approximation on algebraic varieties”, Arithmetic of higher-dimensional algebraic varieties (Palo Alto, CA, 2002), Progr. Math., vol. 226, Birkhäuser Boston, Boston, MA, 2004, p. 43–60.
- harari-here** [Har08] D. HARARI – “Non-abelian descent”, 2008, this volume.
- hassett-here** [Has08] B. HASSETT – “Surfaces over nonclosed fields”, 2008, this volume.
- heath-brown-cubic** [HB83] D. R. HEATH-BROWN – “Cubic forms in ten variables”, *Proc. London Math. Soc.* (3) **47** (1983), no. 2, p. 225–257.
- heath-brown-cayley** [HB03] ———, “The density of rational points on Cayley’s cubic surface”, *Proceedings of the Session in Analytic Number Theory and Diophantine Equations* (Bonn), Bonner Math. Schriften, vol. 360, Univ. Bonn, 2003, p. 33.

- `hb-count` [HB06] ———, “Counting rational points on algebraic varieties”, Analytic number theory, Lecture Notes in Math., vol. 1891, Springer, Berlin, 2006, p. 51–95.
- `heath-brown-moroz` [HBM04] D. R. HEATH-BROWN and B. Z. MOROZ – “On the representation of primes by cubic polynomials in two variables”, *Proc. London Math. Soc. (3)* **88** (2004), no. 2, p. 289–312.
- `hu-keel` [HK00] Y. HU and S. KEEL – “Mori dream spaces and GIT”, *Michigan Math. J.* **48** (2000), p. 331–348, Dedicated to William Fulton on the occasion of his 60th birthday.
- `harari-sko-comp` [HS02] D. HARARI and A. N. SKOROBOGATOV – “Non-abelian cohomology and rational points”, *Compositio Math.* **130** (2002), no. 3, p. 241–273.
- `harari-sko` [HS05] D. HARARI and A. SKOROBOGATOV – “Non-abelian descent and the arithmetic of Enriques surfaces”, *Int. Math. Res. Not.* (2005), no. 52, p. 3203–3228.
- `ht-hol-cubic` [HT] B. HASSETT and Y. TSCHINKEL – “Flops on holomorphic symplectic fourfolds and determinantal cubic hypersurfaces”, [arXiv:0805.4162](https://arxiv.org/abs/0805.4162).
- `hassett-t99` [HT99] ———, “Geometry of equivariant compactifications of \mathbf{G}_a^n ”, *Internat. Math. Res. Notices* (1999), no. 22, p. 1211–1230.
- `harris-t00` [HT00] J. HARRIS and Y. TSCHINKEL – “Rational points on quartics”, *Duke Math. J.* **104** (2000), no. 3, p. 477–500.
- `ht-m06` [HT02] B. HASSETT and Y. TSCHINKEL – “On the effective cone of the moduli space of pointed rational curves”, Topology and geometry: commemorating SISTAG, Contemp. Math., vol. 314, Amer. Math. Soc., Providence, RI, 2002, p. 83–96.
- `ht-stable` [HT03] ———, “Integral points and effective cones of moduli spaces of stable maps”, *Duke Math. J.* **120** (2003), no. 3, p. 577–599.
- `hassett-t03` [HT04] ———, “Universal torsors and Cox rings”, Arithmetic of higher-dimensional algebraic varieties (Palo Alto, CA, 2002), Progr. Math., vol. 226, Birkhäuser Boston, Boston, MA, 2004, p. 149–173.
- `hunt` [Hun96] B. HUNT – *The geometry of some special arithmetic quotients*, Lecture Notes in Mathematics, vol. 1637, Springer-Verlag, Berlin, 1996.
- `isk-manin` [IM71] V. A. ISKOVSKIĖ and Y. I. MANIN – “Three-dimensional quartics and counterexamples to the Lüroth problem”, *Mat. Sb. (N.S.)* **86(128)** (1971), p. 140–166.
- `inose` [Ino78] H. INOSE – “Defining equations of singular $K3$ surfaces and a notion of isogeny”, *Proceedings of the International Symposium on Algebraic Geometry (Kyoto Univ., Kyoto, 1977)* (Tokyo), Kinokuniya Book Store, 1978, p. 495–502.
- `iskovskikh99` [IP99a] V. A. ISKOVSKIĖ and Y. G. PROKHOROV – “Fano varieties”, Algebraic geometry, V, Encyclopaedia Math. Sci., vol. 47, Springer, Berlin, 1999, p. 1–247.
- `isk-pro` [IP99b] ———, “Fano varieties”, Algebraic geometry, V, Encyclopaedia Math. Sci., vol. 47, Springer, Berlin, 1999, p. 1–247.
- `isk71` [Isk71] V. A. ISKOVSKIĖ – “A counterexample to the Hasse principle for systems of two quadratic forms in five variables”, *Mat. Zametki* **10** (1971), p. 253–257.

- `iskovskih79` [Isk79] ———, “Anticanonical models of three-dimensional algebraic varieties”, *Current problems in mathematics*, Vol. 12 (Russian), VINITI, Moscow, 1979, p. 59–157, 239 (loose errata).
- `isk-bir` [Isk01] V. A. ISKOVSKIKH – “Birational rigidity of Fano hypersurfaces in the framework of Mori theory”, *Uspekhi Mat. Nauk* **56** (2001), no. 2(338), p. 3–86.
- `kasp` [Kas08] A. KASPRZYK – “Bounds on fake weighted projective space”, 2008, [arXiv.org:0805.1008](https://arxiv.org/abs/0805.1008).
- `katsylo` [Kat82] P. I. KATSYLO – “Sections of sheets in a reductive algebraic Lie algebra”, *Izv. Akad. Nauk SSSR Ser. Mat.* **46** (1982), no. 3, p. 477–486, 670.
- `katsura` [Kat87] T. KATSURA – “Generalized Kummer surfaces and their unirationality in characteristic p ”, *J. Fac. Sci. Univ. Tokyo Sect. IA Math.* **34** (1987), no. 1, p. 1–41.
- `kawaguchi` [Kaw08] S. KAWAGUCHI – “Projective surface automorphisms of positive topological entropy from an arithmetic viewpoint”, *Amer. J. Math.* **130** (2008), no. 1, p. 159–186.
- `kirillov0` [Kir99] A. A. KIRILLOV – “Merits and demerits of the orbit method”, *Bull. Amer. Math. Soc. (N.S.)* **36** (1999), no. 4, p. 433–488.
- `kleiman` [Kle66] S. L. KLEIMAN – “Toward a numerical theory of ampleness”, *Ann. of Math. (2)* **84** (1966), p. 293–344.
- `kollar-mori` [KM98] J. KOLLÁR and S. MORI – *Birational geometry of algebraic varieties*, Cambridge Tracts in Mathematics, vol. 134, Cambridge University Press, Cambridge, 1998, With the collaboration of C. H. Clemens and A. Corti, Translated from the 1998 Japanese original.
- `kawamata-mm87` [KMM87] Y. KAWAMATA, K. MATSUDA and K. MATSUKI – “Introduction to the minimal model problem”, *Algebraic geometry, Sendai, 1985*, *Adv. Stud. Pure Math.*, vol. 10, North-Holland, Amsterdam, 1987, p. 283–360.
- `kollar-mm92` [KMM92] J. KOLLÁR, Y. MIYAOKA and S. MORI – “Rational connectedness and boundedness of Fano manifolds”, *J. Differential Geom.* **36** (1992), no. 3, p. 765–779.
- `kresch-t` [KT04] A. KRESCH and Y. TSCHINKEL – “On the arithmetic of del Pezzo surfaces of degree 2”, *Proc. London Math. Soc. (3)* **89** (2004), no. 3, p. 545–569.
- `kresch-t-brauer` [KT08] ———, “Effectivity of Brauer-Manin obstructions”, *Adv. Math.* **218** (2008), no. 1, p. 1–27.
- `lehmann` [Leh08] B. LEHMANN – “A cone theorem for nef curves”, 2008, [arXiv:0807.2294](https://arxiv.org/abs/0807.2294).
- `manin-icm` [Man71] Y. I. MANIN – “Le groupe de Brauer-Grothendieck en géométrie diophantienne”, *Actes du Congrès International des Mathématiciens (Nice, 1970)*, Tome 1, Gauthier-Villars, Paris, 1971, p. 401–411.
- `manin86` [Man86] ———, *Cubic forms*, second ed., North-Holland Publishing Co., Amsterdam, 1986.
- `manduchi95` [Man95] E. MANDUCHI – “Root numbers of fibers of elliptic surfaces”, *Compositio Math.* **99** (1995), no. 1, p. 33–58.

- `masser-bounds` [Mas02] D. W. MASSER – “Search bounds for Diophantine equations”, A panorama of number theory or the view from Baker’s garden (Zürich, 1999), Cambridge Univ. Press, Cambridge, 2002, p. 247–259.
- `matsuki-book` [Mat02] K. MATSUKI – *Introduction to the Mori program*, Universitext, Springer-Verlag, New York, 2002.
- `mazur77` [Maz77] B. MAZUR – “Modular curves and the Eisenstein ideal”, *Inst. Hautes Études Sci. Publ. Math.* (1977), no. 47, p. 33–186 (1978).
- `mckinnon` [McK00] D. MCKINNON – “Counting rational points on $K3$ surfaces”, *J. Number Theory* **84** (2000), no. 1, p. 49–62.
- `mcmullen` [McM02] C. T. MCMULLEN – “Dynamics on $K3$ surfaces: Salem numbers and Siegel disks”, *J. Reine Angew. Math.* **545** (2002), p. 201–233.
- `merel96` [Mer96] L. MEREL – “Bornes pour la torsion des courbes elliptiques sur les corps de nombres”, *Invent. Math.* **124** (1996), no. 1-3, p. 437–449.
- `mori-mukai` [MM86] S. MORI and S. MUKAI – “Classification of Fano 3-folds with $B_2 \geq 2$. I”, Algebraic and topological theories (Kinosaki, 1984), Kinokuniya, Tokyo, 1986, p. 496–545.
- `mori-mukai-2` [MM03] ———, “Erratum: “Classification of Fano 3-folds with $B_2 \geq 2$ ” [Manuscripta Math. **36** (1981/82), no. 2, 147–162; MR0641971 (83f:14032)]”, *Manuscripta Math.* **110** (2003), no. 3, p. 407.
- `mori-m81` [MM82] ———, “Classification of Fano 3-folds with $B_2 \geq 2$ ”, *Manuscripta Math.* **36** (1981/82), no. 2, p. 147–162.
- `moore` [Moo65] C. C. MOORE – “Decomposition of unitary representations defined by discrete subgroups of nilpotent groups”, *Ann. of Math. (2)* **82** (1965), p. 146–182.
- `mori82` [Mor82] S. MORI – “Threefolds whose canonical bundles are not numerically effective”, *Ann. of Math. (2)* **116** (1982), no. 1, p. 133–176.
- `manin-ts` [MT86] Y. I. MANIN and M. A. TSFASMAN – “Rational varieties: algebra, geometry, arithmetic”, *Uspekhi Mat. Nauk* **41** (1986), no. 2(248), p. 43–94.
- `nikulin` [Nik81] V. V. NIKULIN – “Quotient-groups of groups of automorphisms of hyperbolic forms by subgroups generated by 2-reflections. Algebro-geometric applications”, Current problems in mathematics, Vol. 18, Akad. Nauk SSSR, Vsesoyuz. Inst. Nauchn. i Tekhn. Informatsii, Moscow, 1981, p. 3–114.
- `pitman-2` [NP89] T. NADESALINGAM and J. PITMAN – “Bounds for solutions of simultaneous diagonal equations of odd degree”, *Théorie des nombres* (Quebec, PQ, 1987), de Gruyter, Berlin, 1989, p. 703–734.
- `peyre95` [Pey95] E. PEYRE – “Hauteurs et mesures de Tamagawa sur les variétés de Fano”, *Duke Math. J.* **79** (1995), no. 1, p. 101–218.
- `peyre-counting` [Pey04] ———, “Counting points on varieties using universal torsors”, Arithmetic of higher-dimensional algebraic varieties (Palo Alto, CA, 2002), Progr. Math., vol. 226, Birkhäuser Boston, Boston, MA, 2004, p. 61–81.
- `peyre-obstr` [Pey05] ———, “Obstructions au principe de Hasse et à l’approximation faible”, *Astérisque* (2005), no. 299, p. Exp. No. 931, viii, 165–193, Séminaire Bourbaki. Vol. 2003/2004.

- pitman-1** [Pit71] J. PITMAN – “Bounds for solutions of diagonal equations”, *Acta Arith.* **19** (1971), p. 223–247. (loose errata).
- poonen-br** [Poo08] B. POONEN – “Insufficiency of the Brauer–Manin obstruction applied to étale covers”, 2008, preprint.
- peyre-t01** [PT01] E. PEYRE and Y. TSCHINKEL (eds.) – *Rational points on algebraic varieties*, Progress in Mathematics, vol. 199, Birkhäuser Verlag, Basel, 2001.
- pukh-icm** [Puk98] A. V. PUKHLIKOV – “Birational automorphisms of higher-dimensional algebraic varieties”, *Proceedings of the International Congress of Mathematicians, Vol. II (Berlin, 1998)*, no. Extra Vol. II, 1998, p. 97–107 (electronic).
- pukh-usp** [Puk07] ———, “Birationally rigid varieties. I. Fano varieties”, *Uspekhi Mat. Nauk* **62** (2007), no. 5(377), p. 15–106.
- saltman** [Sal84] D. J. SALTMAN – “Noether’s problem over an algebraically closed field”, *Invent. Math.* **77** (1984), no. 1, p. 71–84.
- salberger-tors** [Sal98] P. SALBERGER – “Tamagawa measures on universal torsors and points of bounded height on Fano varieties”, *Astérisque* (1998), no. 251, p. 91–258, *Nombre et répartition de points de hauteur bornée* (Paris, 1996).
- sheph2** [SB88] N. I. SHEPHERD-BARRON – “The rationality of some moduli spaces of plane curves”, *Compositio Math.* **67** (1988), no. 1, p. 51–88.
- sheph1** [SB89] ———, “Rationality of moduli spaces via invariant theory”, *Topological methods in algebraic transformation groups* (New Brunswick, NJ, 1988), *Progr. Math.*, vol. 80, Birkhäuser Boston, Boston, MA, 1989, p. 153–164.
- serre90** [Ser90] J.-P. SERRE – *Lectures on the Mordell–Weil theorem*, *Aspects of mathematics*, no. 15, Vieweg, 1990.
- siegel-einheiten** [Sie69] C. L. SIEGEL – “Abschätzung von Einheiten”, *Nachr. Akad. Wiss. Göttingen Math.-Phys. Kl. II* **1969** (1969), p. 71–86.
- silverman** [Sil91] J. H. SILVERMAN – “Rational points on $K3$ surfaces: a new canonical height”, *Invent. Math.* **105** (1991), no. 2, p. 347–373.
- skinner** [Ski97] C. M. SKINNER – “Forms over number fields and weak approximation”, *Compositio Math.* **106** (1997), no. 1, p. 11–29.
- skoro-beyond** [Sko99] A. N. SKOROBOGATOV – “Beyond the Manin obstruction”, *Invent. Math.* **135** (1999), no. 2, p. 399–424.
- skoro-book** [Sko01] ———, *Torsors and rational points*, Cambridge Tracts in Mathematics, vol. 144, Cambridge University Press, Cambridge, 2001.
- serganova-sko** [SS07] V. V. SERGANOVA and A. N. SKOROBOGATOV – “Del Pezzo surfaces and representation theory”, *Algebra Number Theory* **1** (2007), no. 4, p. 393–419.
- serganova-sko2** [SS08] V. SERGANOVA and A. SKOROBOGATOV – “On the equations for universal torsors over Del Pezzo surfaces”, 2008, [arXiv:0806.0089](https://arxiv.org/abs/0806.0089).
- stillman-2006** [STV06] M. STILLMAN, D. TESTA and M. VELASCO – “Groebner bases, monomial group actions, and the Cox rings of Del Pezzo surfaces”, 2006, [arXiv:math/0610261](https://arxiv.org/abs/math/0610261).
- testa-cox08** [TVAV08] D. TESTA, A. VARILLY-ALVARADO and M. VELASCO – “Cox rings of degree one Del Pezzo surfaces”, 2008, [arXiv:0803.0353](https://arxiv.org/abs/0803.0353).

- varillyalvarado-2008 [VAZ] A. VARILLY-ALVARADO and D. ZYWINA – “Arithmetic E_8 lattices with maximal Galois action”, [arXiv:0803.3063](#).
- luijk [vL07] R. VAN LUIJK – “K3 surfaces with Picard number one and infinitely many rational points”, *Algebra Number Theory* **1** (2007), no. 1, p. 1–15.
- vaughan-w95 [VW95] R. C. VAUGHAN and T. D. WOOLEY – “On a certain nonary cubic form and related equations”, *Duke Math. J.* **80** (1995), no. 3, p. 669–735.
- weyl [Wey16] H. WEYL – “Über die Gleichverteilung von Zahlen mod. Eins”, *Math. Ann.* **77** (1916), no. 3, p. 313–352.
- wittenberg03 [Wit04] O. WITTENBERG – “Transcendental Brauer-Manin obstruction on a pencil of elliptic curves”, Arithmetic of higher-dimensional algebraic varieties (Palo Alto, CA, 2002), *Progr. Math.*, vol. 226, Birkhäuser Boston, Boston, MA, 2004, p. 259–267.
- zarhin-deg1 [Zar08] Y. G. ZARHIN – “Del Pezzo surfaces of degree 1 and Jacobians”, *Math. Ann.* **340** (2008), no. 2, p. 407–435.

COURANT INSTITUTE, 251 MERCER STREET, NEW YORK, NY 10012, USA
AND MATHEMATISCHES INSTITUT, GEORG-AUGUST-UNIVERSITÄT GÖTTINGEN,
BUNSENSTRASSE 3-5, D-37073 GÖTTINGEN, GERMANY

E-mail address: tschinkel@cims.nyu.edu